

8535.99915 Euclid . K

E U C L I D E ' s
E L E M E N T S ;

The whole Fifteen BOOKS
Compendiously Demonstrated.

W I T H
A R C H I M E D E S *Theorems*
Of the Sphere and Cylinder, investi-
gated by the *Method of Indivisibles*.

By ISAAC BARROW, D. D. *Late Master*
of Trinity College in Cambridge.

To which is added in this Edition,
E U C L I D E ' s D A T A
with *Marinus's* Preface.

And a Brief T R E A T I S E of
R E G U L A R S O L I D S .

Καθαροὶ ψυχῆς λογικῆς εἰσὶν αἱ μαθηματικαὶ ἐπιστήμαι.

L O N D O N : Printed and Sold by *W. Redmayne*
in *Fewen-street*, *R. Mount* on *Tower-hill*, and *J. and*
B. Sprint in *Little-britain*. 1714.

THE
ELEMENTS

OF THE WHOLE ART OF ARCHITECTURE
AS PRACTISED IN GREECE

WITH
ARCHITECTURAL DECORATIONS

OF THE GREEK AND ROMAN ARCHITECTURE
AS PRACTISED BY THE ARTISTS OF THESE NATIONS

BY
JOHN BARROW, ESQ. OF LINCOLN'S INN

IN TWO VOLUMES
THE FIRST



OF THE
ART OF ARCHITECTURE

AND
THE ART OF BUILDING

IN TWO VOLUMES
THE SECOND

OF THE
ART OF ARCHITECTURE

To the READER.

IF you are desirous, Courteous Reader, to know what I have perform'd in this Edition of the Elements of Euclide, I shall here explain it to you in short, according to the nature of the Work. I have endeavor'd to attain two ends chiefly; the first, to be very perspicuous, and at the same time so very brief, that the Book may not swell to such a Bulk, as may be troublesome to carry about one, in which I think I have succeeded, unless in my absence the Printer's care should frustrate my Design. Some of a brighter Genius, and endued with greater Skill, may have demonstrated most of these Propositions with more nicety, but perhaps none with more succinctness than I have; especially since I alter'd nothing in the number and order of the Author's Propositions; nor presum'd either to take the liberty of rejecting, as less necessary, any of them, or of reducing some of the easier sort into the rank of Axioms, as several have done; and among others, that most expert Geometrician A. Tacquetus C. (whom I the more willingly name, because I think it is but civil to acknowledge that I have imitated him in some Points) after whose most accurate Edition I had no Thoughts of attempting any thing of this nature, till I consider'd that this most learned Man thought fit to publish only eight of Euclide's Books, which he took the pains to explain and embellish, having in a manner rejected and undervalued the other seven, as less appertaining to the Elements of Geometry. But my Province was

To the READER.

originally quite different, not that of writing the Elements of Geometry, after what method soever I pleas'd, but of demonstrating, in as few words as possible I cou'd, the whole Works of Euclide. As to four of the Books, viz. the seventh, eighth, ninth, and tenth, altho' they don't so nearly appertain to the Elements of plain and solid Geometry, as the six precedent and the two subsequent, yet none of the more skilful Geometricians can be so ignorant as not to know that they are very useful for Geometrical matters, not only by reason of the mighty near affinity that is between Arithmetick and Geometry, but also for the knowledge of both measurable and unmeasurable Magnitudes, so exceeding necessary for the Doctrine of both plain and solid figures. Now the noble Contemplation of the five Regular Bodies that is contain'd in the three last Books, cannot without great Injustice be pretermitted, since that for the sake thereof our σοιχειωτης, being a Philosopher of the Platonic Sect, is said to have compos'd this universal System of Elements; as Proclus lib. 2. witnesseth in these Words, *Ὅθεν δὴ καὶ τῆς συμπάσης σοιχειώσεως τέλος προσηύδατο τὴν ἑκκατέρωθεν πλατωνικῶν ἡμετέρων σύνασιν.* Besides, I easily perswaded my self to think, that it would not be unacceptable to any Lover of these Sciences to have in his Possession the whole Euclidean Work, as it is commonly cited and celebrated by all Men. Wherefore I resolv'd to omit no Book or Proposition of those that are found in P. Herigonius's Edition, whose Steps I was oblig'd closely to follow, by reason I took a Resolution to make use of most of the Schemes of the said Book, very well foreseeing that time would not allow me to form

new

TO the READER.

new ones, tho' sometimes I chose rather to do it. For the same reason I was willing to use for the most part Euclide's own Demonstrations, having only express'd them in a more succinct Form, unless perhaps in the second, thirteenth, and very few in the seventh, eighth, and ninth Book, in which it seem'd not worth my while to deviate in any particular from him. Therefore I am not without good hopes that as to this part I have in some measure satisfied both my own Intentions, and the Desire of the Studious. As for some certain Problems and Theorems that are added in the Scholions (or short Expositions) either appertaining (by reason of their frequent Use) to the nature of these Elements, or conducing to the ready Demonstration of those things that follow, or which do intimate the reasons of some principal Rules of practical Geometry, reducing them to their original Fountains, these I say, will not, I hope, make the Book swell to a Size beyond the design'd Proportion.

The other Butt, which I levell'd at, is to content the Desires of those who are delighted more with symbolical than verbal Demonstrations. In which kind, whereas most among us are accusom'd to the Symbols of Gulielmus Oughtredus, I therefore thought best to make use, for the most part, of his. None hitherto (as I know of) has attempted to interpret and publish Euclide after this manner, except P. Herigonius; whose Method (tho' indeed most excellent in many things, and very well accommodated for the particular purpose of that most ingenious Man) yet seems in my Opinion to labour under a double Defect. First, in regard that, altho' of two or more Propositions, produced

TO the READER.

ced for the Proof of any one Problem or Theorem, the former don't always depend of the latter, yet it don't readily enough appear either from the order of each, or by any other manner, when they agree together, and when not; wherefore for want of the Conjunctions and Adjectives, ergo, rursus, &c. many difficulties and occasions of doubt do often arise in reading, especially to those that are Novices. Besides it frequently happens, that the said Method cannot avoid superfluous Repetitions, by which the Demonstrations are oftentimes render'd tedious, and sometimes also more intricate; which Faults my Method doth easily remedy by the arbitrary mixture of both Words and Signs. Therefore let what has been said, touching the Intention and Method of this little Work, suffice. As to the rest, whoever covets to please himself with what may be said, either in Praise of the Mathematicks in general, or of Geometry in particular, or touching the History of these Sciences, and consequently of Euclide himself, (who digested those Elements) and others ~~ΕΓΓΕΝΕΤΑΙ~~ of that kind, may consult other Interpreters. Neither will I (as if I were afraid least these my Endeavors may fall short of being satisfactory to all Persons) alledge as an Excuse (tho' I may very lawfully do it) the want of due time which ought to be employ'd in this Work, nor the Interruption occasion'd by other Affairs, nor yet the want of requisite help for these Studies nor several other things of the like nature. But what I have here employ'd my Labour and Study in for the Use of the ingenuous Reader, I wholly submit to his Censure and Judgment, to approve if useful, or reject if otherwise.

J. B.

Ad amicissimum Virum, I. C. de EUCLIDE
contracto, Εὐκλείδης.

Factum bene! didicit Laconice loqui
Senex profundus, & aphorismos induit.
Immensa dudum margo commentarii
Diagramma circuit minutum; utque Insula
Problema breve natabat in vasto mari.
Sed unda jam detumuit; & glossa arctior
Stringit Theoremata: minoris anguli
Lateribus ecce totus Euclides jacet,
Inclusus olim velut Homerus in nuce;
Pluteoque farsina modo qui incubuit, levis
En sit manipulus. Pelle in exigua latet
Ingens Mathesis, matris utero Hercules,
In glande quercus, vel Ithaca Eurus in pila.
Nec mole dum decrescit, usu fit minor;
Quin auctior jam evadit, & cumulatius
Contracta prodest erudita pagina.
Sic uberè magis liquor è presso affluit;
Sic pleniori vasa inundat sanguinis
Torrente cordis Systole; sic fusus
Procurrit aquor ex Abyla angustiis.
Tantilli operis ars tanta referenda unice est
BAROVIANO nomini, ac solertia.
Sublimis euge mentis ingentium potens!
Cui invitum nil, arduum esse nil solet;
Sic usque pergas prospero contamine,
Radiusque multum debeat ac abacus tibi;
Sic crescat indies feracior seges,
Simili colonum germine assiduò beans.
Specimen futura messis hic fiet labor.
Magnæque fama illustra hac præludia.
Juvenis dedit qui tanta, quid dabit senex?

Car. Robotham, CANTAB.
Coll. Trin. Sen. Soc.

The

The Explication of the Signs or Characters.

=	}	Equal.
>	}	Greater.
<	}	Lesser.
+	}	More, or to be added.
-	}	Less, or to be subtracted.
	}	The Differences, or Excess; Also, that all the quantities which follow, are to be subtracted, the Signs not being changed.
×	}	Multiplication, or the Drawing one side of a Rectangle into another. The same is denoted by the Conjunction of letters; as $AB = A \times B$.
√	}	The Side or Root of a Square, or Cube, &c.
Q & q	}	A Square.
C & c	}	A Cube.
Q. Q.	}	The ratio of a square number to a square number.

Other Abbreviations of words, where-ever they accur, the Reader will without trouble understand of himself; saving some few, which, being of less general use, we refer to be explained in their places.

The




THE FIRST BOOK
OF
EUCLIDE'S
ELEMENTS.

Definitions.

- I. **A** Point is that which has no part.
- II. A Line is a longitude without latitude.
- III. The ends, or limits, of a line are Points.
- IV. A Right Line is that which lies equally betwixt its Points.
- V. A Superficies is that which has only longitude and latitude.
- VI. The extremes, or limits, of a Superficies are lines.
- VII. A plain Superficies is that which lies equally betwixt its lines.
- VIII. A plain Angle is the inclination of two lines the one to the other, the one touching the other in the same plain, yet not lying in the same strait line.
- IX. And if the lines which contain the Angle be right lines, it is called a right-lined Angle.
- A
- X. When

The first Book of



X. When a right line CG standing upon a right line AB, makes the angles on either side thereof, CGA, CGB, equal one to the other, then both those equal angles are right angles; and the right line CG, which standeth on the other, is termed a Perpendicular to that (AB) whereon it standeth.

Note, When several angles meet at the same point (as at G) each particular angle is described by three letters; whereof the middle letter sheweth the angular point, and the two other letters the lines that make that angle: As the angle which the right lines CG, AG make at G, is called CGA, or AGC.



XI. An obtuse angle is that which is greater than a right angle; as ACD.

XII. An acute angle is that which is less than a right angle; as ACB.


XIII. A Limit, or Term, is the end of any thing.

XIV. A Figure is that which is contained under one or more terms.

XV. A Circle is a plain figure contained under one line, which is called a Circumference; unto which all lines drawn from one point within the figure, and falling upon the circumference thereof, are equal the one to the other.

XVI. And that point is called the Center of the Circle.

XVII. A Diameter of a circle is a right line drawn through the center thereof, and ending at the circumference on either side, divi-



dividing the circle into two equal parts.

XVIII. A Semicircle is a figure which is contained under the diameter, and under that part of the circumference which is cut off by the diameter.

In the circle EABCD, E is the center, AC the diameter, ABC the semicircle.

XIX. Right-lined figures are such as are contained under right lines.


XX. Three-sided or Trilateral figures are such as are contained under three right lines.

XXI. Four-sided or Quadrilateral figures are such as are contained under four right lines.


XXII. Many-sided figures are such as are contained under more right lines than four.



XXIII. Of Trilateral figures, that is an Equilateral Triangle, which hath three equal sides; as the Triangle A.



XXIV. Isosceles is a Triangle which hath only two sides equal; as the Triangle B.



XXV. Scalenum is a Triangle whose three sides are all unequal; as C.

A.

XXVI. Of

The first Book of




XXVI. Of these Trilateral figures, a right-angled Triangle is that which has one right angle; as the Triangle A.

XXVII. An Amblygonium, or obtuse-angled Triangle, is that which has one angle obtuse; as B.



XXVIII. An Oxygonium, or acute-angled Triangle, is that which has three acute angles; as C.

An Equiangular, or equal-angled figure is that whereof all the angles are equal. Two figures are equiangular, if the several angles of the one figure be equal to the several angles of the other. The same is to be understood of Equilateral figures.



XXIX. Of Quadrilateral, or four-sided figures, a Square is that whose sides are equal, and angles right; as ABCD.



XXX. A figure on the one part longer, or a long square, is that which hath right angles, but not equal sides; as ABCD.


XXXI. A

EUCLIDE'S *Elements*.

5




XXXI. A Rhombus, or diamond-figure, is that which has four equal sides, but is not right-angled; as A.



XXXII. A Rhomboides, or diamond-like figure, is that whose opposite sides, and opposite angles, are equal; but has neither equal


nor right angles; as GLMH.



XXXIII. All other quadrilateral figures besides these are called *Trapezia*, or *Tables*; as GNDH.




XXXIV. Parallel, or equidistant right lines are such, which being in the same superficies, if infinitely produced, would never meet; as A and B.



XXXV. A Parallelogram is a quadrilateral figure, whose opposite sides are parallel, or equidistant; as GLHM.

A 3

XXXVI. In



XXXVI. In a Parallelogram ABCD, when a diameter AC, and two lines EF, HI parallel to the sides, cutting the diameter in one and the same point G, are drawn, so that the Parallelogram be divided by them into four parallelograms; those two DG, GB, through which the diameter passeth not, are called Complements; and the other two HE, FI, through which the diameter passeth, the Parallelograms standing about the diameter.

A Problem is, when something is proposed to be done or effected.

A Theoreme is, when something is proposed to be demonstrated.

A Corollary is a consectary, or some consequent truth gained from a preceding demonstration.

A Lemma is the demonstration of some premise, whereby the proof of the thing in hand becomes the shorter.

Postulates or Petitions.

1. **F**rom any point to any point to draw a right line.
2. To produce a right line finite, strait forth continually.
3. Upon any center, and at any distance, to describe a circle.

Axioms.

1. **T**Hings equal to the same third, are also equal one to the other.

As $A=B=C$. Therefore $A=C$. Or therefore all, A, B, C, are equal the one to the other.

Note, When several quantities are joined the one to the other continually with this mark $=$, the first quantity is by virtue of this axiome equal to the last, and every one to every one: In which case we often ab-

stain

It is not necessary to stain from citing the axiome, for brevity's sake; altho' the force of the consequence depend thereon.

2. If to equal things you add equal things, the wholes shall be equal.

3. If from equal things you take away equal things, the things remaining will be equal.

4. If to unequal things you add equal things, the wholes will be unequal.

5. If from unequal things you take away equal things, the remainders will be unequal.

6. Things which are double to the same third, or to equal things, are equal one to the other. Understand the same of triple, quadruple, &c.

7. Things which are half of one and the same thing, or of things equal, are equal the one to the other. Conceive the same of subtriple, subquadruple, &c.

8. Things which agree together, are equal one to the other.

The converse of this axiome is true in right lines and angles, but not in figures, unless they be like.


Moreover, magnitudes are said to agree, when the parts of the one being applied to the parts of the other, they fill up an equal or the same place.

9. Every whole is greater than its part.

10. Two right lines cannot have one and the same segment (or part) common to them both.

11. Two right lines meeting in the same point, if they be both produced, they shall necessarily cut one the other in that point.

12. All right angles are equal the one to the other.



13. If a right line BA falling on two right lines

A 4

AD, CB,

The first Book of

AD, CB, make the internal angles on the same side, BAD, ABC, less than two right angles, those two right lines produced shall meet on that side, where the angles are less than two right angles.

14. Two right lines do not contain a space.

15. If to equal things, you add things unequal, the excess of the wholes shall be equal to the excess of the additions.

16. If to unequal things equal be added, the excess of the wholes shall be equal to the excess of those which were at first.


17. If from equal things, unequal things be taken away, the excess of the remainders shall be equal to the excess of the wholes.

18. If from things unequal, things equal be taken away, the excess of the remainders shall be equal to the excess of the wholes.

19. Every whole is equal to all its parts taken together.

20. If one whole be double to another, and that which is taken away from the first to that which is taken away from the second, the remainder of the first shall be double to the remainder of the second.

The Citations are to be understood in this manner; When you meet with two numbers, the first shews the Proposition, the second the Book; as by 4, 1. you are to understand the fourth Proposition of the first Book; and so of the rest. Moreover, ax. denotes Axiome, post. Postulate, def. Definition, sch. Scholium, cor. Corollary.



PROPOSITION I.




Upon a finite right line given *AB*, to describe an equilateral triangle *ACB*.

From the centers *A* and *B*, at the distance of *AB*, or *BA*, *a* describe two circles *a* 3. *post.* cutting each other in the point *C*; from whence *b* draw two right lines *CA*, *CB*. Then is *AC* *c* *b* 1. *post.* *= AB* *c* *= BC* *d* *= AC*. *e* Wherefore the Tri-*c* 15. *def.* angle *ACB* is equilateral. Which was to be done. *d* 1. *ax.* *e* 23. *def.*

Scholium.

After the same manner upon the line *AB* may be described an Iſosceles triangle, if the distances of the equal circles be taken greater or less than the line *AB*.

PROP. II.



At a point given *A*, to make a right line *AG* equal to a right line given *BC*.

From the center *C*, at the distance of *CB*, *a* de-*a* 3. *post.* scribe the circle *CBE*. *b* join *AC*; upon which *b* 1. *post.* *c* raise the equilateral triangle *ADC*. *d* Produce *c* 1. 1. *DC* to *E*. From the center *D*, at the distance *d* 2. *post.* of

e 2. post. of DE, describe the circle DEH; and let DA e be produced to the point G in the circumference thereof. Then $AG = CB$.

f 15. def. For $DG = DE$, and $DA = DC$. Wherefore $AG = CE$ $k = BC$ $l = AG$, Which was to be done.

h 3. ax. The putting of the point A within or without the line BC varies the cases; but the construction, and the demonstration, are every where alike.


k 15. def.

l 1. ax.

Schol.

The line AG might be taken with a pair of compasses; but the so doing answers to no Postulate, as Proclus well intimates.

PROP. III.



Two right lines, A and BC, being given, from the greater BC to take away the right line BE equal to the lesser A.

At the point B a draw the right line $BD = A$.

The circle described from


the center B at the distance of BD shall cut off $BE = BD$ $c = A$ $d = BE$. Which was to be done.

b 15. def.

c constr.

d 1. ax.

PROP. IV.




If two triangles BAC, EDF, have two sides of the one BA, AC equal to two sides of the other ED, DF, each to its correspondent side (that is, $BA = ED$, and

and $AC=DF$) and have the angle A equal to the angle D contained under the equal right lines; they shall have the base BC equal to the base EF ; and the triangle BAC shall be equal to the triangle EDF ; and the remaining angles B, C , shall be equal to the remaining angles E, F , each to each, under which the equal sides are subtended.

If the point D be applied to the point A , and the right line DE plac'd upon the right line AB , the point E shall fall upon B , because $DE = AB$, a *hyp.* also the right line DF shall fall upon AC , because the angle $A = D$. moreover the point F shall fall upon the point C , because $AC = DF$. b 14. *ax.* Therefore the right lines EF, BC shall agree, because they have the same Terms, and consequently are equal. Wherefore the triangles BAC, DEF , and the angles B, E , as also the angles C, F , do agree, and are equal. Which was to be Demonstrated.

PROP. V.



The angles ABC, ACB , at the base of an Isosceles triangle ABC , are equal one to the other: And if the equal sides AB, AC be produc'd, the angles CBD, BCE , under the base, shall be equal one to the other.

a Take $AE=AD$; and b join a 3. 1.
CD, and BE. b 1. *post.*


Because, in the triangles ACD, ABE , are $AB = AC$, and $AE = AD$, and the angle A common to them both, d *constr.* e therefore is the angle $ABE = ACD$, and the angle e 4. 1.
 $AEB = ADC$, and the base $BE = CD$; also $EC = DB$. Therefore in the triangles BEC, BDC g shall be the angle $ECB = DBC$. Which g 4. 1.
was to be Dem. Also therefore the angle $EBC = DCB$. but the angle $ABE = ACD$; therefore h before.
the angle $ABC = ACB$. Which was to be Dem. k 3. *ax.*

Coroll.

Coroll.

Hence, every equilateral triangle is also equiangular.

PROP. VI.



If two angles ABC , ACB of a triangle ABC be equal the one to the other, the sides AC , AB subtended under the equal angles, shall also be equal one to the other.

If the sides be not equal, let one be bigger than the other, suppose $BA > CA$. *a* Make $BD = CA$, and *b* draw the line CD .


a 3. I.*b* 1. post.*c* suppos.*d* hyp.*e* 4. I.*f* 9. ax.

In the triangles DBC , ACB , because $BD = CA$, and the side BC is common, and the angle $DBC = ACB$, the triangles DBC , ACB shall be equal the one to the other, *a* part to the whole. *f* Which is impossible.

Coroll.

Hence, Every equiangular triangle is also equilateral.

PROP. VII.



Upon the same right line AB two right lines being drawn AC , BC , two other right lines equal to the former, AD , BD , each to each (viz. $AD = AC$, and $BD = BC$) cannot be drawn from the same points A , B , on the same side C , to several points, as C and D , but only to C .

1. Case. If the point D be set in the line AC , it is plain that AD is not equal to AC .

a 9. ax.

2. Case. If the point D be placed within the triangle ACB , then draw the line CD , and produce BDF , and BCE . Now you would have $AD = AC$.


then

then the angle $ADC = ACD$; as also, because b 5. 1.
 $BD = BC$, the angle $FDC = b$ ACD . there- c *suppos.*
 fore is the angle $FDC = d$ ACD , that is, the d 9. *ax.*
 angle $FDC = ADC$. d *Which is impossible.*

3. *Case.* If D falls without the triangle ACB ,
 let CD be joined.

Again the angle $ACD = ADC$, and the e 5. 1.
 angle $BCD = BDC$. f Therefore the angle f 9. *ax.*
 $ACD = BDC$, viz. the angle $ADC = BDC$.
Which is impossible. Therefore, &c.

PROP. VIII.



If two triangles
 ABC , DEF , have
 two sides AB , AC
 equal to two sides
 DE , DF , each to
 each, and the base
 BC equal to the base EF , then the angles con-
 tained under the equal right lines shall be equal,
 viz. A to D .

Because $BC = EF$, if the base BC be laid on a *hyp.*
 the base EF , b they will agree: therefore whereas b *ax.* 8.
 $AB = DE$, and $AC = DF$, the point A will fall c *hyp.*
 on D (for it cannot fall on any other point, by
 the precedent proposition) and so the sides of
 the angles A and D are coincident; d wherefore d 8. *ax.*
 those angles are equal. *Which was to be Dem.*

Coroll.

1. Hence, Triangles mutually equilateral, are
 also mutually equiangular.


2. Triangles mutually equilateral, are equal e 4. 1.
 one to the other.

PROP.

PROP. IX.

a 3. I.

b 1. I.

c constr.
d 8. I.

To bisect, or divide into two equal parts, a right-lined angle given BAC .

a Take $AD = AE$, and draw the line DE ; upon which b make an equilateral triangle DFE . draw the right line AF ; it shall bisect the angle.

For $AD = AE$, and the side AF is common, and the base $DF = FE$. d therefore the angle $DAF = EAF$. Which was to be done.

Coroll.


Hence it appears, how an angle may be cut into any equal parts, as 4, 8, 16, &c. to wit, by bisecting each part again.

The method of cutting angles into any equal parts required, by a Rule and Compass, is as yet unknown to Geometricians.

PROP. X.

a 1. I.

b 9. I.

c constr.
d 4. I.

To bisect a right line given AB .


Upon the line given AB a erect an equilateral triangle ABC ; and b bisect the angle C with the right line CD . That line shall also bisect the line given AB .

For $AC = BC$, and the side CD is common, and the angle $ACD = BCD$. therefore $AD = BD$. Which was to be done.

The practice of this and the precedent Proposition is easily shewn by the construction of the 1 Prop. of this Book.

PROP.

PROP. XI.




From a point C in a right line given AB to erect a right line CF at right angles.

a Take on either side of the point given CD ^{a 3. 1.} $= CE$. upon the right line DE ^{b 1. 1.} erect an equilateral triangle, draw the line FC , and it will be the perpendicular required.

For the triangles DFC , EFC are mutually *c e-c constr.* equilateral; *d* therefore the angle $DCF = ECF$. *ed 8. 1.* therefore FC is perpendicular. *Which was to be done. e 10. def.*

The practice of this and the following is easily performed by the help of a square.

PROP. XII.



Upon an infinite right line given AB , from a point given that is not in it, to let fall a perpendicular right line CG .

From the center C describe a circle cutting the right line given AB in the points E and F . Then bisect EF in ^{a 3. post.} G , and draw the right line CG , which will be ^{b 10. 1.} the perpendicular required.

Let the lines CE , CF be drawn. The triangles EGC , FGC are mutually *c equilateral. d there-c constr.* fore the angles EGC , FGC are equal, and by *d 8. 1.* *e* consequence right. *e* Wherefore GC is a per- *e 10. def.* pendicular. *Which was to be done.*

PROP. XIII.



When a right line AB standing upon a right line CD maketh angles ABC , ABD ; it maketh either two right angles, or two angles equal to two right.

IF

a def. 10.

b 11. 1.

c 19. ax.

d 3. ax.

e 2. ax.

If the angles ABC, ABD be equal, a then they make two right angles; if unequal, then from the point B b let there be erected a perpendicular BE . Because the angle ABC $c =$ to a right $+ ABE$, and the angles ABD $d =$ to a right $- ABE$, therefore shall be $ABC + ABD$ $e =$ to two right angles $+ ABE - ABE =$ two right angles. Which was to be demonstrated.

Corollaries.


1. Hence, if one angle ABD be right, the other ABC is also right; if one acute, the other is obtuse, and so on the contrary.

2. If more right lines than one stand upon the same right line at the same point, the angles shall be equal to two right.

3. Two right lines cutting each other make angles equal to four right.

4. All the angles made about one point, make four right; as appears by Coroll. 2.

P R O P. XIV.



If to any right line AB , and a point therein B , two right lines, not drawn from the same side, do make the angles ABC, ABD on each side equal to two right, the lines CB, BD shall make one strait line.


a 13. 1.

b hyp.

c 9. ax.

If you deny it, let CB, BE make one right line, then shall be the angle $ABC + ABE$ $a =$ two right angles $b = ABC + ABD$. Which is absurd.

P R O P. XV.



If two right lines AB, CD cut thro' one another, then are the two angles which are opposite, viz. CEB, AED , equal one to the other.


a 13. 1.

b 3. ax.

For the angle $AEC + CEB$ $a =$ to two right angles $= AEC + AED$; b therefore $CEB = AED$. Which was to be demonstrated.

Schol.


Schol.



If to any right line GH, and in it a point A, two right lines being drawn EA, AF, and not taken on the same side, make the vertical (or opposite) angles D and B equal, those right lines EA, AF, do meet directly and make one strait line.

For two right angles are a equal to the angle $D + A$ $a = B + A$. b therefore, EA, AF, are in a strait line. Which was to be demonstrated. a 13. 1. b 14. 1.


Schol.



If four right lines EA, EB, EC, ED, proceeding from one point E, make the angles vertically opposite equal the one to the other, each two lines, AE, EB, and CE, ED, are placed in one strait line.

For because the angle AEC + AED + CEB + DEB $a =$ to 4 right angles, therefore the angle AEC + AED $b =$ CEB + DEB $=$ to two right angles, c therefore CED and AEB are strait lines. Which was to be demonstrated. a 4. c. 13. 1. b hyp. and c 14. 1.

PROP. XVI.




One side BC of any triangle ABC being produc'd, the outward angle ACD will be greater than either of the inward and opposite angles CAB, CBA.

Let the right lines AH, BE a bisect the sides AC, BC; a 10. 1. b produce EF = BE, and HI, $b = b$ 3. 1. c 10. 1. d 1. post.

$b = AH$. and join FC , and IC ; and produce ACG .

c *confr.* Because $CE = EA$, and $EF = EB$, and the
d 15. 1. angle $FEC = BEA$, the angle ECF shall be
e 4. 1. equal to EAB . By the like argument is the an-
f 15. 1. gle $ICH = ABH$. Therefore the whole angle
g 9. *ax.* ACD (*f* BCG) *g* is greater than either the angle
 CAB or ABC . Which was to be demonstrated.

PROP. XVII.



Two angles of any triangle ABC , which way soever they be taken, are less than two right angles.

Let the side BC be produced. Because the angle $ACD + ACB = 2$ right angles, and the angle $ACD < A$, c therefore $A + ACB >$ then two right angles. After the same manner is the angle $B + ACB >$ then two right. Lastly, the side AB being produced, the angle $A + B$ will be also less than two right angles. Which was to be demonstrated.

Coroll.

1. Hence it follows that in every triangle, wherein one angle is either right or obtuse, the two others are acute angles.



2. If a right line AE make unequal angles with another right line D , one acute AED , the other obtuse AEC , a perpendicular AD let fall from any point A to the other line CD , shall fall on that side the acute is of.


For if AC , drawn on the side of the obtuse angle, be a perpendicular, then in the triangle AEC shall $AEC + ACE$ be * greater than two right angles. Which is contrary to the precedent Prop.

3. All the angles of an equilateral triangle, and the two angles of an Isosceles triangle that are upon the base, are acute.

PROP.

PROP. XVIII

The greatest side AC of every triangle ABC subtends the greatest angle ABC .




From AC take away $AD = AB$, and join BD . Therefore is the angle $ADB = ABD$.

But $ADB < C$; therefore is $ABD < C$; therefore the whole angle $ABC < C$. After the same manner, shall be $ABC < A$. Which was to be dem.

PROP. XIX.

In every triangle ABC , under the greatest angle A is subtended the greatest side BC .




For if AB be supposed equal to BC , then will be the angle $A = C$, which is contrary to the

Hypothesis: and if $AB < BC$, then shall be the angle $C < A$, which is against the Hypothesis. Wherefore rather $BC < AB$; and after the same manner $BC < AC$. Which was to be dem.

PROP. XX.


Of every triangle ABC two sides BA, AC , any way taken, are greater than the side that remains BC .



Produce the line BA , take $AD = AC$, and draw the line DC , then shall the angle D be equal to ACD , therefore is the whole angle $BCD < D$; therefore $BD < BC$. Which was to be demonstrated.

PROP. XXI.

If from the utmost points of one side BC of a triangle ABC two right lines BD, CD be drawn to any point within the triangle, then are both those two lines shorter than the two other sides of the




B 2

the

the triangle, BA, CA ; but do contain a greater angle, BDC .

- Let BD be produced to E . Then is $CE + ED$
 a 20. 1. $a = CD$, and BD common to both b then shall be
 $BD + DE + EC = CD + BD$. Again, $BA = AE$
 b 4. ax. $a = BE$. b therefore $BA + AC = BE + EC$. Where-
 fore 1. $BA + AC = BD + DC$. 2. The angle BDC
 c 16. 1. $c = DEC = A$. Therefore the angle $BDC = A$.
 Which was to be demonstrated.

PROP. XXII.



To make a triangle FKG of three right lines FK, FG, GK , which shall be equal to three right lines given A, B, C . Of which it is necessary that any two taken together be longer than the third.

- From the infinite line DE take DF, FG, GH
 a 3. 1. equal to the lines given, A, B, C . Then if from
 b 3. post. the b centers F and G by the distances of FD
 and GH , two circles be drawn cutting each
 c 15. def. other in K , and the right lines KF, KG be joined,
 the triangle FKG shall be made, c whose
 d 1. ax. sides FK, FG, GK are equal to the three lines
 DF, FG, GH d that is to the three lines given
 A, B, C . Which was to be done.

PROP. XXIII.




a 1. post.

At a point A in a right line given AB , to make a right-lined angle A equal to a right-lined angle given D .

Draw the right line

line CF cutting the sides of the angle given any ways; *b* make $AG = CD$; upon AG *c* raise a trian- *b* 3. 1.
gle equilateral to the former CDF , so that AH be *c* 21. 1.
equal to DF , and GH to CF . then shall you have
the angle A *d* $= D$. Which was to be done. *d* 8. 1.

PROP. XXIV.



If two triangles ABC , DEF have two sides of the one triangle AB , AC equal to two sides of the other triangle DE , DF , each to other, and have the angle A greater than the angle EDF contained under the equal right lines, they shall also have the base BC greater than the base EF .

a Let the angle EDG be made equal to A , *a* 23. 1.
and the side DG *b* $= DF$ *c* $= AC$; and let EG , *b* 3. 1.
and FG be joined. *c* hyp.

1. Case. If EG fall above EF ; because AB *d* $= DE$, and AC *e* $= DG$, and the angle A *e* $= EDG$, *e* constr.
f therefore is $BC = EG$. But because DF *e* $= DG$, *f* 4. 1.
g therefore is the angle $DFG = DGF$; *h* therefore *g* 5. 1.
is the angle $DFG = FFG$, and by consequence *h* 9. ax.
the angle EFG *k* $= EGF$, *k* wherefore EG (BC) *k* 19. 1.
 $= EF$.

B 3

2. Case.

19. ax. 2. Case. If the base EF falls in the same place with EG, it is evident that $EG (BC) \sqsubset EF$.
- m 21. 1. 3. Case. If EG fall below EF, then because DG $\perp$ GE $m \sqsubset$ DF + FE, if from both DG, DF be taken away, which are equal, EG (BC) remains
- n 5. ax. $n \sqsubset EF$. Which was to be demonstrated.

PROP. XXV.




If two triangles ABC , DEF have two sides AB , AC equal to two sides DE , DF , each to other, and have the base BC greater than the base EF , they shall also have the angle A contained under the equal right lines greater than the angle D .

a 4. 1.

b 24. 1.

For if the angle A be said to be equal to D , then is the base $BC = EF$, which is against the Hypothesis. If it be said the angle $A \supset D$, then b will be $BC \supset EF$, which is also against the Hyp. Therefore $BC \sqsubset EF$. Which was to be dem.

PROP. XXVI.



If two triangles BAC , EDG have two angles of the one B, C equal to the two angles of the other E, DGE , each to his correspondent angle, and have also one side of the one equal to one side of the other, either that side which lyeth betwixt the equal angles, or that which is subtended under one of the equal angles; the other sides also

EUCLIDE's Elements.

25


also of the one shall be equal to the other sides of the other each to his correspondent side, and the other angle of the one shall be equal to the other angle of the other.

1. Hyp. Let BC be equal to EG , which are the sides that lie between the equal angles. Then I say $BA = ED$, and $AC = DG$, and the angle $A = EDG$. For if it be said that $ED < BA$, then a let EH^a 3. 1. be made equal to BA , and let the line GH be drawn.

Because $AB^b = HE$, and $BC^c = EG$, and the b suppos. angle $B^c = E$, therefore shall be the angle EGH^c hyp. $d = C^e = DGE$. f Which is absurd. After the d 4. 1. same manner let AC be equal to DG , d then will e hyp. the angle A be equal to EDG . f 9. ax.

2. Hyp. Let AB be equal to ED . Then I say $BC = EG$, and $AC = DG$, and the angle $A = EDG$. For if EG be greater than BC , make $EI = BC$, and join the line DI . Now because $AB^g = ED$, and g hyp. $BC^h = EI$, and the angle $G^g = E$; therefore h suppos. will be the angle $EID^k = C^l = EGD$. m Which k 4. 1. is absurd. Therefore is $BC = EG$, and so as be- l hyp. fore $AC = DG$, and the angle $A = EDG$. Which m 16. 1. was to be demonstrated.

PROP. XXVII.




If a right line EF falling upon two right lines AB, CD , make the alternate angles AEF, DFE , equal the one

to the other, then are the right lines AB, CD parallel.

If AB, CD be said not to be parallel, produce them till they meet in G . which being supposed, the outward angle AEF will be a greater than a 16. 1. the inward angle DFE , to which it was equal by Hypothesis. Which is repugnant.

PROP. XXVIII.



If a right line EF falling upon two right lines, AB, CD make the outward angle AGE of the one line equal to CHG the inward and opposite angle B 4 of

of the other on the same side, or make the inward angles on the same side AGH, CHG equal to two right angles, then are the right lines AB, CD parallel.

Hyp. 1. Because by Hypothesis the angle $AGE = CHG$, *a* therefore are BGH, CHG alternate angles and equal: And *b* therefore are AB and CD parallel.

Hyp. 2. Because by Hypothesis the angle $AGH + CHG =$ to two right, *a* $= AGH + BGH$, *b* thence is $CHG = BGH$; and *c* therefore AB, CD are parallel. Which was to be demonstrated.

PROP. XXIX.



If a right line EF fall upon two parallels, AB, CD , it will both make the alternate angles DHG, AGH equal each to other, and the outward angle BGE equal to the inward and opposite angle on the same side DHE , as also the inward angles on the same side AGH, CHG equal to two right angles.

It is evident that $AGH + CHG = 2$ right angles; *a* otherwise AB, CD would not be parallel, which is contrary to the Hyp. But moreover the angle $DHG + CHG = 2$ right; therefore is $DHG + AHG = BGE$. Which was to be dem.

Coroll.



Hence it follows that every parallelogram AC having one angle right A , the rest are also right.

For $A + B = 2$ right angles. Therefore whereas A is right, *b* must B be also right. By the same argument are C and D right angles.

PROP.

PROP. XXX.

Right lines (AB, CD) parallel to one and the same right line EF , are also parallel the one to the other.

Let GI cut the three right lines given any ways, then because AB, EF are parallel, will be the angle $AGI = EHI$. also because CD and EF are parallel, will be the angle $EHI = DIG$. Therefore the angle $AGI = DIG$. whence AB and CD are parallel. Which was to be dem.

PROP. XXXI.

From a point given A to draw a right line AE parallel to a right line given BC .

From the point A draw a right line AD to any point of the given right line; with which at the point thereof A make an angle $DAE = ADC$. then will AE and BC be parallel. Which was to be done.

PROP. XXXII.

Of any triangle ABC one side BC , being drawn out, the outward angle ACD shall be equal to the two inward opposite angles A, B , and the three inward angles of the triangle, A, B, ACB , shall be equal to two right angles.

From C draw CE parallel to BA . Then is the angle $A = ACE$, and the angle $B = ECD$. Therefore $A + B = ACE + ECD = ACD$. Which was to be demonstrated.

I affirm $ACD + ACB =$ two right angles; therefore $A + B + ACB = 2$ right angles. Which was to be demonstrated.

Coroll.

1. The three angles of any triangle taken together

ther are equal to three angles of any other triangle taken together. From whence it follows,

2. That if in one triangle, two angles (taken severally, or together) be equal to two angles of another triangle (taken severally, or together) then is the remaining angle of the one equal to the remaining angle of the other. In like manner, if two triangles have one angle of the one equal to one of the other, then is the sum of the remaining angles of the one triangle equal to the sum of the remaining angles of the other.

3. If one angle in a triangle be right, the other two are equal to a right. Likewise, that angle in a triangle which is equal to the other two, is it self a right angle.

4. When in an Isosceles the angle made by the equal sides is right, the other two upon the base are each of them half a right angle.

5. An angle of an equilateral triangle makes two third parts of a right angle. For $\frac{1}{3}$ of two right angles is equal to $\frac{2}{3}$ of one.

Schol.

By the help of this Proposition you may know how many right angles the inward and outward angles of a right-lined figure make; as may appear by these two following Theorems.

THEOREM I.



All the angles of a right-lined figure do together make twice as many right angles, bating four, as there are sides of the figure.

From any point within the figure, let right lines

lines be drawn all thro' the angles of the figure, which shall resolve the figure into as many triangles as there are sides of the figure. Wherefore, whereas every triangle affords two right angles, all the triangles taken together will make up twice as many right angles as there are sides. But the angles about the said point within the figure make up four right; therefore, if from the angles of all the triangles you take away all the angles which are about the said point, the remaining angles, which make up the angles of the figure, will make twice as many right angles, bating four, as there are sides of the figure. *Which was to be dem.*

Coroll.

Hence, All right-lined figures of the same species have the sums of their angles equal.

THEOREM II.

All the outward angles of any right-lined figure, taken together, make up four right angles.

For all the several inward angles of a figure with the several outward angles of the same make two right angles; therefore all the inward angles, together with the outward, make twice as many right angles as there are sides of the figure; but (as it was now shewn) all the inward angles with four right, make twice as many right as there are sides of the figure; therefore the outward angles are equal to four right angles. *Which was to be dem.*

Coroll.


All right-lined figures of whatsoever species have the sums of their outward angles equal.

PROP. XXXIII.

If two equal and parallel lines AB, CD be joined together with two other right lines AC, BD, then are those lines also equal and parallel.

Draw a line from C to B. Now because AB and CD are parallel, and the angle $ABC = BCD$; a 29. 1. and also by Hypothesis $AB = CD$, and the side CB

com-



- b 4. 1. common, therefore is $AC = BD$, and the angle $ACB = BD$. c whence also AC, BD are parallel.

P R O P. XXXIV.


 In parallelograms, as $ABCD$, the opposite sides AB, CD , and AC, BD , are equal each to other; and the opposite angles A, D , and ABD, ACD are also equal; and the diameter BC bisects the same.

- a hyp. Because AB, CD are parallel, b therefore is the angle $ABC = BCD$. Also because AC, BD are a parallel, b therefore is the angle $ACB = CBD$; c therefore the whole angle $ACD = ABD$. After the same manner is $A = D$. Moreover because the angles ABC, ACB lie at each end of the side CB , and are equal to BCD, CBD , d therefore is $AC = BD$, and $AB = CD$, and so the triangle $ABC = CBD$. Which was to be dem.

Schol.

Every four sided figure $ABDC$ having the opposite sides equal, is a parallelogram.

- a 27. 1. For by 8. 1. the angle $ABC = BCD$; a wherefore AB, CD are parallel. In likemanner is the angle $BCA = CBD$; a wherefore AC, BD are also parallel. b Therefore $ABCD$ is a parallelogram. Which was to be demonstrated.




AB , thro' the point assigned C .

Take in the line AB any point, as E . From the centers E and C at any distance draw two equal circles EF, CD . From the center F by the space of EC draw a circle FD , which shall cut the former circle CD in the point D . Then shall the line drawn CD be parallel to AB . for as it was before demonstrated, $CEFD$ is a parallelogram.

P R O P.

PROP. XXXV.



Parallelograms, $BCDA$, $BCEF$, which stand upon the same base BC , and between the same parallels AF , BC , are equal one to the other.

For $AD = BC = a$ 34. 1.
 EF , add DE common to both, b then is $AE = b$ 2. ax.
 DF . But also $AB = DC$, and the angle $A = C$ 29. 1.
 CDF . d Therefore is the triangle $ABE = DCF$. d 4. 1.
take away DGE common to both triangles, e 3. ax.
then is the Trapezium $ABGD = EGCF$. add
 BGC common to both, f then is the parallelo- f 2. ax.
gram $ABCD = EBCF$. Which was to be dem.

The demonstration of any other cases is not unlike, but much more plain and easy.

Schol.




If the side AB of a right-angled parallelogram $ABCD$ be conceived to be carried along perpendicularly thro' the whole line BC , or BC thro' the whole line AB , the Area or content of the Rectangle $ABCD$ shall be produc'd by that motion. Hence a rectangle is said to be made by the drawing or multiplication of two contiguous sides. For example's sake; let AB be supposed four foot, and BC be three: draw 3 into 4, there will be produced 12 square feet for the Area of the Rectangle.

This being supposed, the dimension of any parallelogram (*EBCF) is found out by this Theorem. For the Area thereof is produced from the figure of altitude BA drawn into the base BC . So the Area Prop. 35. of the parallelogram $AC = EBCF$, is made by the drawing of BA into BC , therefore, &c.

PROP.


PROP. XXXVI.



Parallelograms, $BCDA$, $GHFE$, standing upon equal bases BC , GH , and betwixt the same parallels AF , BH , are equal to the other.

- a hyp. Draw BE and CE . Because $BC = GH$ $b = EF$, c therefore is $BCEF$ a parallelogram. Whence the parallelogram $BCDA = BCEF = GHFE$.
 b 34. i. $d = BCEF$
 c 33. i. $d = GHFE$.
 d 35. i. Which was to be dem.


PROP. XXXVII.



Triangles, BCA , BCD , standing upon the same base BC , and between the same parallels BC , EF , are equal one to the other.

- a 31. i. Draw BE parallel to CA , a and CF parallel
 b 34. i. to BD . Then is the triangle $BCA = \frac{1}{2}$ of the
 c 35. i. and parallelogram $BCAE = \frac{1}{2}$ $BDFC = BCD$.
 7. ax. Which was to be dem.

PROP. XXXVIII.




Triangles, BCA , EDF , set upon equal bases BC , EF , and between the same parallels GH , BF , are equal the one to the other.

- Draw BG parallel to CA and FH parallel to ED . Then is the triangle $BCA = \frac{1}{2}$ Pgr. $BCAG = \frac{1}{2}$ $EDHF = EDF$.
 a 34. i. c
 b 36. i. and Which was to be dem.
 7. ax. Schol.

- c 34. i. If the base BC be greater than EF , then is the triangle $BAC > EDF$, and so on the contrary.

PROP.

PROP. XXXIX.



Equal triangles BCA , BCD , standing on the same base BC , and on the same side, they are also between the same parallels AD , BC .


If you deny it, let another line AF be parallel to BC ; and let CF be drawn. Then is the triangle CBF $a = CBA$ $b = CBD$. c Which is absurd.

a 37. I.

b hyp.

c 9. ax.

PROP. XL.



Equal triangles BCA , EFD , standing upon equal bases BC , EF , and on the same side, they are betwixt the same parallels.


If you deny it, let another line AH be parallel to BF , and let FH be drawn. Then is the triangle EFH $a = BCA$ $b = EFD$. c Which is absurd.

a 38. I.

b hyp.

c 9. ax.

PROP. XLI.



If a Pgr. $ABCD$ have the same base BC with the triangle BCE , and be between the same parallels AE , BC then is the Pgr. $ABCD$ double to the triangle BCE .

Let the line AC be drawn. Then is the triangle BCA $a = BCE$. therefore is the Pgr. $ABCD$. $b = 2 BCA$ $c = 2 BCE$. Which was to be dem.

a 37. I.

b 34. I.

c 6. ax.


Schol.

From hence may the Area of any triangle BCE be found, for whereas the Area of the Pgr. $ABCD$ is produced by the altitude drawn into the base, therefore shall the Area of a triangle be produced by half the altitude drawn into the base, or half the

The first Book of

the base drawn into the altitude, as if so be the base BC be 8, and the altitude 7, then is the Area of the triangle BCE, 28.

PROP. XLII.




To make a Pgr. ECGF equal to a triangle given ABC in an angle equal to a right-lined angle given A.

a 31. I. Through A draw AG parallel to BC b make
b 23. I. the angle BCG = D. c bisect the base BC in E,
c 10. I. and draw EF parallel to CG. then is the problem resolved.

For AE being drawn, the angle ECG is equal
d 31. I. to D by construction, and the triangle BAC d
e 41. I. = 2 AEC = Pgr. ECGF. Which was to be done.

PROP. XLIII.




In every Pgr. ABCD, the complements DG, GE of those Pgrs HE, FI, which stand about the diameter, are equal one to the other.

a 34. I. For the triangle ACD
b 3. ax. a = ACB, and the triangle AGH a = AGE, and the triangle GCF a = GCI. b Therefore the Pgr. DG = BG. Which was to be dem.

PROP.

PROP. XLIV.




Upon a right line given *A*, to make a Parallelogram *FL* at a right-lined angle given *C*, equal to a triangle given *B*.

a Make a Pgr. $FD =$ to the triangle *B*, so a 42. r. that the angle GFE may be equal to *C*. Produce GF till FH be equal to the line given *A*. thro' *H* draw IL parallel to EF , which let b 31. r. DE produced meet in the point *I*. let DG drawn forth meet with a right line drawn from *I* in the point *K*. thro' *K* draw KL parallel to GH , with which let EF drawn out meet at *M*, and IH at *L*. Then shall FL be the Pgr. required.

For the Pgr. $FL = FD = B$, and the angle c 43. r. $MFH = GFE = C$. Which was to be done. d 15. r.

PROP. XLV.




Upon a right line given FG to make a Pgr. FL equal to a right-lined figure given $ABCD$, at a right-lined angle given *E*.

C

Res


- Resolve the right-lined figure given into triangles BAD, BCD. then *a* make a Pgr. FH = BAD, so that the angle F may be equal to E, FI being produced, *a* make on HI the Pgr. IL = BCD. Then is the Pgr. FL *b* = FH + IL *c* = *c* constr. ABCD. Which was to be done.

Schol.



Hence is easily found the excess, HE, whereby any right-lined figure, A, exceeds a less right-lined figure, B; namely, if to some right line, CD, both be applied, Pgr. DF = A, and DH = B.

PROP. XLVI.



Upon a right line given AD to describe a square AC.

a Erect two perpendiculars AB, DC, *b* equal to the line given AD; then join BC, and the thing required is done.

a 11. 3.

b 3. 1.

c constr.

d 28. 1.

e constr.

f 34. 1.

g sch. 29. 1.


h 29. def.

For, whereas the angle A + D = 2 right, therefore are AB, DC parallel. But they are also equal; therefore AD, BC are both parallel and equal; therefore the figure AC is a Pgr. and equilateral. Moreover the angles are all right, *g* because one, A, is right; *h* therefore AC is a square. Which was to be done.

After the same manner you may easily describe a Rectangle contained under two right lines given.

PROP.

PROP. XLVII.



In right angled Triangles, BAC , the square BE , which is made of the side BC that subtends the right angle BAC is equal to both the squares BG , CH , which are made of the sides AB , AC containing the right angle.

Join AE , and AD ; and draw

AM parallel to CE .

Because the angle $DBC = FBA$, add the angle ABC common to them both; then is the angle $ABD = FBC$. Moreover $AB = FB$, and $BD = BC$; therefore is the triangle $ABD = FBC$. But the Pgr. $BM = 2 ABD$, and the Pgr. $BG = 2 FBC$ (for GAC is one right line by Hyp. and 14. 1.) therefore is the Pgr. $BM = BG$. By the same way of argument is the Pgr. $CM = CH$. Therefore is the whole $BE = BG + CH$. Which was to be dem.

Schol.

This most excellent and useful Theorem has deserved the title of *Pythagoras's Theorem*, because he was the Inventer of it. By the help of which the addition and subtraction of squares are performed; to which purpose serve the two following Problems.

C 2

PRO


PROBLEM I.

*Andr.
Tacq.*

2 It. I.

b 47. I.

€ 2. 4x.




To make one square equal
to any number of squares
given.

Let three squares be given whereof the sides are AB, BC, CE. *a* Make the right angle FBZ having the sides infinite; and on them transfer BA and BC; join AC, then is $AC^2 = AB^2 + BC^2$. Then transfer AC from B to X, and CE the third side given from B to E; join

EX. b Then is $EXq = EBq (CEq) + BXq (ACq) = CEq + ABq + BCq$. Which was to be done.

PROBLEM II.



Two unequal right lines
being given AB, BC , to make
a square equal to the diffe-
rence of the two squares of the
given lines, AB, BC .


A B C From the center B, at the distance of BA, describe a circle; and from the point C erect a perpendicular CE meeting with the circumference in E; and draw BE. *a* Then is BEq (BAq) = BCq + CEq. *b* Therefore BAq - BCq = CEq. Which was to be done.

a 47. I.

b 3. ax.

PRO-

PROBLEM III.




Any two sides of a right-angled triangle ABC being known, to find out the third.

Let the sides AB , AC encompassing the right angle, be, the one 6 foot, the other 8. Therefore, whereas $ACq + ABq = 64 + 36 = 100 = BCq$, thence is $BC = \sqrt{100} = 10$. 47. 1.

Otherwise, let the sides AB , BC be known, the one six foot, the other 10. Therefore since $BCq - ABq = 100 - 36 = 64 = ACq$, thence 47. 1, is $AC = \sqrt{64} = 8$. Which was to be dem.

PROP. XLVIII.



If the square made upon one side BC of a triangle be equal to the squares made on the other sides of the triangle AB , AC , then the angle BAC comprehended under those two other sides of the triangle AB , AC , is a right angle.

Draw to the point A in AC a perpendicular line $DA = AB$, and join CD .

Now is $CDq = ADq + ACq = ABq + ACq$ a 47. 1. $= BCq$. * Therefore is $CD = BC$. And therefore * See the the triangles CAB , CAD are mutually equilateral. Wherefore the angle $CAB = CAD$ Theor. $=$ a right angle, Which was to be dem. b 8. 1. c constr.


Schol.

We assumed in the demonstration of the last Proposition, $CD = BC$, because CDq was equal to BCq : our assumption we prove by the following theorem.

C 3

THEO-

THEOREME.



The squares AF , CG of equal right lines AB , CD are equal one to the other : And the sides IK , LM of equal squares NK , PM , are equal one to the other.

1. Hyp. Draw the diameters EB , HD . Then it is evident that AF is *a* equal to the triangle EAB twice taken, and *b* equal to the triangle HCD twice taken, and equal to *a* CG . Which was to be done.

2. Hyp. If it may be, let LM be greater than IK . Make $LT = IK$, and let LS be *a* equal to LT . Therefore is LS *b* $= NK$ *c* $= LQ$. Which is absurd.

a 34. 1.
b 4. 1. &
6. ax.
a 46. 1.
b 1. part
c hyp.
d 9. ax.

Coroll.


After the same manner any rectangles equilateral one to another, are demonstrated also to be equal.

The End of the first Book.

THE


THE SECOND BOOK OF EUCLIDE'S ELEMENTS.

Definitions.



- I. **E** Very right-angled Parallelogram AB CD is said to be contained under two right lines AB, AD comprehending a right angle.

Therefore when you meet with such as these, the rectangle under BA, AD, or for shortness sake the rectangle BAD, or $BA \times AD$ (or ZA , for $Z \times A$) the rectangle meant is that which is contained under the right lines BA, AD set at right angles.



- II. In every Pgr. FHIK, any one of those parallelograms which are about the diameter, together with its two complements is called a Gnomon. *As the Pgr. $FB + BI + GA$ (EHM) is a Gnomon; and likewise the Pgr. $FB + BI + EM$ (GKA) is a Gnomon.*

PROP. I.



If two right lines AF , AB , be given, and one of them AB divided into as many parts or segments as you please; the rectangle comprehended under the two whole right lines AB , AF , shall be equal to all the rectangles contained under the whole line AF and the several segments, AD , DE , EB .

a II. I.

b 19. ax. I.

c 34. I.

a Set AF perpendicular to AB . Thro' F draw an infinite line FG perpendicular to AF . From the points, D , E , B erect perpendiculars DH , EI , BG . Then is AG a rectangle comprehended under AF , AB , and is b equal to the rectangles AH , DI , EG , that is (because DH , EI , AF are equal) to the rectangles under AF , AD , under AF , DE , under AF , EB . Which was to be dem.

Schol.

If two right lines given be both divided into how many parts soever, the product of the whole multiplied into itself shall be the same with that of the parts multiplied into themselves.

a I. 2.

b 2. ax.

For let Z be $= A + B + C$, and $Y = D + E$; then, because $DZ = DA + DB + DC$, and $EZ = EA + EB + EC$, and $YZ = DZ + EZ$, b shall ZY be $= DA + DB + DC + EA + EB + EC$. Which was to be dem.

From hence is understood the manner of multiplying compounded right lines into compounded. For you must take all the Rectangles of the parts, and they will present you with the Rectangle of the wholes.


But whensoever in the multiplication of lines into themselves you meet with these signs — intermingled with these +, you must also have particular regard to the signs. For of + multiplied into — ariseth —; but of — into — ariseth +, ex. gr. let + A be to be multiplied into $B - C$; then because + A is not affirmed of all B , but only of a part of it, whereby it exceeds C , therefore AC must remain denied; so that the product will be

be $AB - AC$. Or thus; because B consists of the parts C and $B - C$, * thence $AB = AC + A \times B - C$. * 1. 3. take away AC from either, then $AB - AC = A \times B - C$. In like manner if A be to be multiplied into $B - C$, then seeing by reason of the figure — that A is not denied of all B . but only of so much as it exceeds C , therefore AC must remain affirmed. whence the product will be $AB + AC$. Or thus; because $AB = AC + A \times B - C$; take away all throughout, and there will be $AB = AC - A \times -B + C$; add AC to either, and there will be $AB + AC = A \times B - C$.

This being sufficiently understood, the nine following Propositions, and innumerable others of that kind, arising from the comparing of lines multiplied into themselves (which you may find done to your hand in *Vieta* and other Analytical Writers) are demonstrated with great facility, by reducing the matter for the most part to almost a simple work.

Furthermore, * it appears that the product of * 19. ax. any magnitude multiplied into the parts of any number, is equal to the product of the same multiplied into the whole number: As; $A + 7 A = 12 A$, and $4 A \times 5 A + 4 A \times 7 A = 4 A \times 12 A$. Wherefore what is here deliver'd of the multiplying of right lines into themselves, the same may be understood of the multiplying of numbers into themselves, so that whatsoever is affirmed concerning lines in the nine following Theorems, holds good also in numbers; seeing they all immediately depend and are deriv'd from this first,

P R O P. II.




If a right line Z be divided any-wise into two parts, the rectangles comprehended under the whole line Z and each of the segments A , E , are equal to the square made of the whole line Z .

Q.E.D.

21.2.

I say that $ZA + ZE = Zq$. For take $B = Z$; then is $BA + BE = BZ$, that is (because $B = Z$) $ZA + ZE = Zq$. Which was to be dem.

PROP. III.




If a right line Z be divided any-wise into two parts, the rectangle comprehended under

21.2.

the whole line Z and one of the segments E is equal to the rectangle made of the segments A , E , and the square described on the said segment E .

I say $ZE = AE + Eq$. For $EZ = EA + Eq$.

PROP. IV.



If a right line Z be cut any-wise into two parts, the square made of the whole line


23.2.

Z is equal both to the squares made of the segments A , E , and to twice a rectangle made of the parts A , E .

22.2.

21.2.

I say that $Zq = Aq + Eq + 2AE$. For $ZA = Aq + AE$, and $ZE = Eq + EA$. Therefore whereas $ZA + ZE = Zq$, thence is $Zq = Aq + Eq + 2AE$. Which was to be dem.



Otherwise thus; Upon the right line AB make the square AD , and draw the diameter EB ; through C , the point wherein the line AB is divided, draw the perpendicular CF ; and thro' the point G draw HI parallel to AB .

Because the angle $EHG = A$ is a right angle, and AEB is half a right, therefore is the remaining angle HGE half a right angle. Therefore is $HE = HG = EF = AC$, so that HF is the square of the right line AC . After the same manner is CI proved to be CBq . Therefore AG , GD are rectangles under AC , CB . wherefore the whole square $AD = ACq + CBq + 2ACB$. Which was to be dem.

Coroll.

Coroll.

1. Hence it appears that the Parallelograms which are about the diameter of a square are also squares themselves.

2. That the diameter of any square bisects its angles.

3. That if $A = \frac{1}{2} Z$, then $Zq = 4 Aq$, and $Aq = \frac{1}{4} Zq$. As on the contrary, if it be so that $Zq = 4 Aq$, then is $A = \frac{1}{2} Z$.

PROP. V.

A ——— I ——— I ——— B If a right line AB
C D be cut into equal
into unequal parts, AD, DB, the rectangle compre-
hended under the unequal parts AD, BD, together
with the square that is made of the intermediate
part CD, is equal to the square that is made of the
half line CB.

I say that $CBq = ADB + CDq$.

For these are $\left\{ \begin{array}{l} CBq. \\ a CDq + CDB + DBq + CDB. \\ \text{all equal.} \end{array} \right. \left\{ \begin{array}{l} CDq + b CBD (c AC \times BD) + CDB. \\ CDq + d ADB. \end{array} \right.$

a 4. 2.
b 3. 2.
c hyp.
d 1. 2.

This Theorem is somewhat differently ex-
press'd, and more easily demonstrated thus; A
Rectangle made of the sum and the difference of two
lines A, E, is equal to the difference of their squares.

For if $A + E$ be multiplied into $A - E$, there
arise $Aq - AE + EA - Eq = Aq - Eq$. Which
was to be dem.

Schol.

A ——— C ——— E ——— D ——— B If the line AB be di-
vided otherwise, (viz.)
nearer to the point of bisection, in E; Then is
 $AEB \sqsubset ADB$.

For if $AEBa = CBq - CEq$, and $ADB a = CBq a$ 5. 2. and
 $- CDq$. Therefore, whereas $CDq \sqsubset CEq$, thence 3. ar.
is $AEB \sqsubset ADB$. Which was to be dem.


Coroll.

Coroll.

1. Hence is $ADq + DBq \sqsubset AEq + EBq$. For
 p 4. 2. $ADq + DBq + 2 ADB b = ABq b = AEq +$
 $EBq + 2 AEB$. Therefore because $2 AEB \sqsubset 2$
 ADB . thence is $ADq + DBq \sqsubset AEq + EBq$.
Which was to be dem.

c 3. ax. 2. Hence is $ADq + DBq - AEq + c + EBq$
 $= 2 AEB - 2 ADB$.

P R O P. VI.



If a right line A be di-
 vided into two equal parts,
 and another right line E
 added to the same directly in one right line, then the
 rectangle comprehended under the whole and the line
 added, (viz. $A + E$) and the line added E , together
 with the square which is made of $\frac{1}{2}$ the line A , is
 equal to the square of $\frac{1}{2} A + E$ taken as one line.

e 4. & 3. I say that $\frac{1}{4} Aq (a Q: \frac{1}{2} A) + AE + Eq = Q$.
 Cor. 4. 2. $\frac{1}{2} A + E$. a For, $Q: \frac{1}{2} A + E = \frac{1}{4} Aq + Eq + AE$.

Coroll.

Hence it follows that if 3 right lines $E, E + \frac{1}{2} A$,
 $E + A$ be in Arithmetical proportion, then the
 Rectangle contained under the extreme terms E ,
 $E + A$, together with the square of the difference
 $\frac{1}{2} A$. is equal to the square of the middle term
 $E + \frac{1}{2} A$.

P R O P. VII.



If a right line Z be divi-
 ded any-wise into two parts,
 the square of the whole line
 Z together with the square made of one of the seg-
 ments E , is equal to a double rectangle comprehended
 under the whole line Z and the said segment E , to-
 gether with the square made of the other segment A .

a 4. 2. I say that $Zq + Eq = 2 ZE + Aq$. For Zq
 b 3. 2. $a = Aq + Eq + 2 AE$. and $2 ZE b = 2 Eq +$
 $2 AE$. Which was to be dem.

Coroll.

Coroll.

Hence it follows that the square of the difference of any two lines Z , E , is equal to the squares of both the lines less by a double rectangle comprehended under the said lines.

$\therefore$ For $Zq + Eq - 2 ZE = Aq = Q: Z - E$.

c 7. 2. and
3. ax.

P R O P. VIII.

If a right line Z be divided any-wise into two parts, the rectangle comprehended under the whole line Z and one of the segments E four times, together with the square of the other segment A , is equal to the square of the whole line Z and the segment E taken as one line $Z + E$.

I say that $4 ZE + Aq = Q: Z + E$. For $2 ZE + Aq = Zq + Eq + Aq$. Therefore $4 ZE + Aq = Zq + Eq + 2 ZE = Q: Z + E$. Which was to be dem.

a 7. 2. and
3. ax.
b 4. 1.

P R O P. IX.

If a right line AB be divided into equal parts AC , CB , and into unequal parts, AD , DB , then are the squares of the unequal parts AD , DB , together, double to the square of the half line AC , and to the square of the difference CD .

I say that $ADq + DBq = 2 ACq + CDq$. For $ADq + DBq = ACq + CDq + 2 ACD + DBq$. But $2 ACD = 2 BCD$. $\therefore ADq + DBq = ACq + CDq + 2 BCD + DBq = ACq + CDq + 2 CBq = 2 ACq + CDq$. Therefore $ADq + DBq = 2 ACq + CDq$. Which was to be dem.

a 4. 1.
b hyp.
c 7. 2.
d 2. ax.

This may be otherwise deliver'd and more easily demonstrated thus; The aggregate of the squares made of the sum and the difference of two right lines A , E , is equal to the double of the squares made from those lines.

For $Q: A + E = Aq + Eq + 2 AE$. and $Q: A - E = Aq + Eq - 2 AE$. These added together make $2 Aq + 2 Eq$. Which was to be dem.

a 4. 2.
b sch. 7. 2.

PROP.


-The second Book of
PROP. X.



If a right line *A* be divided into two equal parts; and another line be added in a right line with the same, then is the square of the whole line, together with the added line, (as being one line,) together with the square of the added line *E*, double to the square of $\frac{1}{2} A$ and the added line *E*, taken as one line.

- a 4. 2. I say that $Eq + Q: A + E$, i. e. $a Aq + 2 Eq + 2$
 b cor. 4. 2. $AE = 2 Q: \frac{1}{2} A + 2 Q: \frac{1}{2} A + E$. For $2 Q: \frac{1}{2}$
 c 4. 2. $Ab = \frac{1}{2} Aq$. And $2 Q: \frac{1}{2} A + E$ $c = \frac{1}{2} Aq +$
 $2 Eq + 2 AE$. Which was to be dem.

PROP. XI.



To cut a right line given *AB* in point *G*, so that the rectangle comprehended under the whole line *AB*, and one of the segments *BG* shall be equal to the square that is made of the other segments *AG*.

- a 46. 1. Upon *AB* describe the square *AC*. b Bisect
 b 10. 1. the side *AD* in *E*, and draw the line *EB*; from the line *EA* produced take *EF = EB*. On *AF* make the square *AH*. Then is $AH = AB \times BG$.

- c 6. 2. For *HG* being drawn out to *I*; the rectangle *DH* + $EAq = EFq = EBq = BAq + EAq$.
 d constr. Therefore is $DH = BAq =$ to the square *AC*.
 e 47. 1. Take away *AI* common to both, then remains the square *AH = GC*, that is, $AGq = AB \times BG$.
 f 3. ax. Which was to be done.

Schol.

- ¶ 6. 13. This Proposition cannot be performed by numbers; * for there is no number that can be so divided, that the product of the whole into one part shall be equal to the square of the other part.

PROP.

EUCLIDE'S Elements.

47

PROP. XII.

In obtuse-angled triangles, ABC, the square that is made of the side AC subtending the obtuse angle ABC is greater than the squares of the sides BC, AB that contain the obtuse angle ABC, by a double rectangle contained under one of the sides BC, which are about the obtuse angle ABC, on which side produced the perpendicular AD falls, and under the line BD, taken without the triangle from the point on which the perpendicular AD falls to the obtuse angle ABC.

I say that $ACq = CBq + ABq + 2 CB \times BD$.

For these are all equal.


$\left\{ \begin{array}{l} a \text{ } ACq. \\ b \text{ } CBq + 2 CBD + BDq + ADq. \\ c \text{ } CBq + 2 CBD + ABq. \end{array} \right.$	$a \text{ } CDq + ADq.$	$a \text{ } 47. \text{ I.}$
	$b \text{ } CBq + 2 CBD + BDq + ADq.$	$b \text{ } 4. \text{ I.}$
	$c \text{ } CBq + 2 CBD + ABq.$	$c \text{ } 47. \text{ I.}$

Scholium.

Hence, the sides of any obtuse angled triangle ABC being known, the segment BD intercepted betwixt the perpendicular AD and the obtuse angle ABC, as also the perpendicular itself AD shall be easily found out.

Thus. Let AC be 10, AB 7, CB 5. Then is ACq 100, ABq 49, CBq 25. And $ABq + CBq = 74$. Take that out of 100, then will 26 remain for 2 CBD. Wherefore CBD shall be 13; divide this by CB 5, there will $2 \frac{3}{5}$ be found for BD. Whence AD will be found out by the 47. I.

PROP. XIII.



In acute-angled triangles ABC, the square made of the side AB subtending the acute angle ACB, is less than the squares made of the sides AC, CB comprehending the acute angle ACB by a double rectangle contained under one of the sides BC, which are about the acute angle ACB on which the perpendicular AD falls, and under the line DC taken within the triangle from the perpendicular AD to the acute angle ACB.

I

I say that $ACq + BCq = ABq + 2 BCD$.

a 47. 1.
b 7. 2.
c 47. 1.


For these are } $ACq + BCq$.
equal. } $a ADq + DCq + BCq$.
 } $b ADq + BDq + 2 BCD$.
 } $c ABq + 2 BCD$.

Coroll.

Hence, *The sides of an acute-angled triangle ABC being known, you may find out the segment DC intercepted betwixt the perpendicular AD and the acute angle ACB, as also the perpendicular it self AD.*

Let AB be 13, AC 15, BC 14. Take ABq (169) from $ACq + BCq$, that is, from $225 + 196 = 421$. Then remains 252 for $2 BCD$. wherefore BCD will be 126. divide this by BC 14, then will 9 be found out for DC. From whence it follows $AD = \sqrt{225 - 81} = 12$.

P R O P. XIV.



To find a square ML equal to a right-lined figure given A.


- a 45. 1. a Make the rectangle $DB = A$, and produce the greater side thereof DC to F, so that $CF = CB$.
b 10. 1. b bisect DE in G, about which as the center at the distance of GF describe the circle FHD, and draw out CB till it touch the circumference in H. Then shall be $CHq = * ML = A$.
* 46. 1. c constr. For let GH be drawn. Then is $A c = DB c$
d 5. 2. and $= DCF d = GFq - GCq e = HCq e = ML$.
3. ax. Which was to be done.
e 47. 1. and

The End of the second Book,


THE

THE THIRD BOOK OF EUCLIDE'S ELEMENTS.

Definitions.




- I. **E** Qual circles (GABC, HDEF) are such whose diameters are equal; or, from whose centers right lines drawn GA, HD, are equal.



- II. A right line AB is said to touch a circle FED, when touching the same, and being produced, it cutteth it not.

The right line FG cuts the circle FED.



- III. Circles DAC, ABE (and also FBG, ABE) are


The third Book of

are said to touch one to the other, which touch, but cut not one the other.

The circle BFG cuts the circle FCH



IV. In a circle GABD, right lines FE, KL are said to be equally distant from the center, when perpendiculars GH, GN drawn from the center G to them are equal. And that line BC is said to be furthest distant from it, on whom the greater perpendicular GI falls.



V. A segment of a circle (ABC) is a figure contained under a right line AC, and a portion of the circumference of a circle ABC.



VI. An angle of a segment CAB, is that angle which is contained under a right line CA and an arch of a circle AB.

VII. An angle ABC is said to be in a segment ABC, when in the circumference thereof some point B is taken, and from it right lines AB, CB, drawn to the ends of the right line AC, which is the base of the segment; then the angle ABC contained under the adjoined lines AB, CB, is said to be angle in a segment.

VIII. But when the right lines AB, BC comprehending the angle ABC, do receive any periphery of the circle ADC, then the angle ABC is said to stand upon that periphery.



IX. A sector of a circle (ADB) is when an angle ADB is set at the center D of that circle; namely, that figure ADB, comprehended under the right lines AD, BD containing the angle, and the part of the circumference received by them AB.



X. Like segments of a circle (ABC, DEF) are those which include equal angles (ABC, DEF); or, in whom the angles ABC, DEF are equal.

PROP. I.



To find the center F of a circle given ABC.


Draw a right line AC any-wise in the circle, which bisect in E, thro' E draw a perpendicular DB, and bisect the same in F, the point F shall be the center.

If you deny it, let G a point without the line DB be the center, (for it cannot be in the line BD, since that cannot be divided equally in any point but F;) let the lines GA, GC GE drawn. Now if G be the center, *a* then is $GA = GC$, and $AE = EC$ by construction, and the side GE common. *b* 8. 1. Therefore are the angles GEA, GEC equal, and *c* 10. def. 1. consequently right. *d* Therefore the angled *e* $GEC = FEC$: *e* Which is absurd.

D 2

Coroll.


Hence, if a right line BD bisect any right line AC in a circle at right angles, the center shall be in the line BD that cuts the other.



Andr.
Tacq.

The center of a circle is easily found out by applying the top of a square Q to the circumference thereof. For if the right line DE that joins the points D, E, in which the sides of the square QD, QE cut the circumference, be bisected in A, the point A shall be the center. The demonstration whereof depends upon Prop. 31. of this Book.

PROP. II.



If in the circumference of a circle CAB any two points A, B be taken, the right line AB which joins those two points shall fall within the circle.


Take in the right line AB any point D; from the center C draw CA, CD, CB. Because CA = CB, therefore is the angle ACD = BCD. But the angle CDB < ACD, therefore is CDB < BCD, therefore CB < CD. But CB only reaches the circumference, therefore CD comes not so far; wherefore the point D is within the circle. The same may be proved of any other point in the line AB. And therefore the whole line AB falls within the circle. Which was to be dem.

Coroll.

Coroll.

Hence, if a right line touch a circle, so that it cut it not, it touches but in one point.

PROP. III.



If in a circle *EABC* a right line *BD* drawn thro' the center, bisect any other line *AC* not drawn through the center, it shall also cut it at right angles in *F*: And if it cuts it at right angles, it shall also bisect the same.

From the center *E* let the lines *EA*, *EC* be drawn.


1. *Hyp.* Because $AF = FC$, and $EA = EC$, and the side *EF* common; the angles *EFA*, *EFC* shall be equal, and consequently right. *Which was to be dem.*

Hyp. 2. Because $EFA = EFC$, and the angle $EAF = ECF$, and the side *EF* common; therefore is $AF = FC$. Therefore is *AC* cut into two equal parts. *Which was to be dem.*

Coroll.

Hence, in any equilateral or Isosceles triangle, if a line drawn from the vertical angle bisect the base, that line is perpendicular to it. And on the contrary, a perpendicular drawn from the vertical angle bisects the base.

PROP. IV.



If in a circle *ACD* two right lines *AB*, *CD* cut thro' one another, yet neither of them pass thro' the center *E*, then neither of those lines are divided into equal parts.

For if one line pass thro' the


D 3

the center, it appears that it cannot be bisected by the other; because by Hypothesis, the other does not pass thro' the center.

If neither of them pass thro' the center, then from the center E draw EF: now if AB, CD were both bisected in F, then *a* would the angles EFB, EFD be both right, and consequently equal. *b* Which is absurd,

a 3. 3.
b 9. ax.

PROP. V.




a 15. def. 1.
b 9. ax.

If two circles BAC, BDC cut one the other, they shall not have the same center E.

For otherwise the lines EB, EDA drawn from E the common center, would DE be *a* = EB *a* = EA. *b* Which is absurd.

PROP. VI.




a 15. def. 1.
b 9. ax.

If two circles BAC, BDE, inwardly touch one the other (in B) they have not one and the same center F.

For otherwise the right lines FB, FDA drawn from the center F, would be FD *a* = FB *a* = FA. *b* Which is absurd.

PROP.

PROP. VII.



If in AB the diameter of a circle, some point G be taken, which is not the center of the circle, and from that point certain right lines GC , GD , GE fall on the circle, the greatest line shall be that (GA) in which is the center F ; the least, the remainder of the sameline (GB)

And of all the other lines, the line GC nearest to that which was drawn thro' the center is always greater than any line farther removed GD ; and only two lines are equal GE , GH , which fall upon the circle from the same point, on each side of the least GB or of the greatest GA .

From the center F draw the right lines FC , FD , FE ; and make the angle $BFH = BFE$. a 23. 1.


1. $GF + FC$ (that is GA) a $\square$ GC . Which a 20. 1. was to be dem.

2. The side FG is common, and $FC = FD$, b 15. def. 1. and the angle $GFC = GFD$; c wherefore the c 9. ax. base $GC = GD$. d 24. 1.

3. FB (FE) e $\supset$ $GE + GF$. f Therefore FG , e 20. which is common, being taken away from both, f 5. ax. $BG \supset EG$.

4. The side FG is common, and $FE = FH$, and the angle $BFH = BFE$; b Therefore is $GE = g$ constr GH . But that no other line GD from the point h 4. 1. G can be equal to GE , or GH , is already proved. Which was to be dem.

The third Book of
PROP. VIII.



If some point A be taken without a circle, and from that point be drawn certain right lines AI, AH, AG, AF to the circle, and of those one AI be drawn through the center K , and the others any-wise; of all those lines that fall on the concave of the circumference, that is the greatest AI which is drawn through the center; and of the others, that which is nearest (AH) to the line that passes through the center is greater than that which is more distant AG . But of all those lines that fall on the convex part of the circle, the least is that AB which is drawn from the point A to the diameter IB ; and of the others, that (AC) which is nearest to the least, is less than that which is farther distant AD . And from that point there can be only two equal right lines AC, AL drawn, which shall fall on the circumference on each side of the least line AB or of the greatest AI .


From the center K draw the right lines KH, KG, KF, KC, KD, KE . and make the angle $AKL = AKC$.

- a 20. I. 1. $AI (AK + KH) a \sqsubset AH$.
 b 24. I. 2. The side AK is common, and $KH = KG$, and the angle $AKH \sqsubset AKG$; b therefore the base $AH \sqsubset AG$.
 c 20. I. 3. $KA c \sqsupset KC + CA$. From hence take away KC, KB that are equal; then will remain $AB d \sqsupset AC$.
 d 5. ax. 4. $AC + CK e \sqsupset AD + DK$. From thence take away CK, DK that are equal; then remains $AC f \sqsupset AD$.
 e 21. I.
 f 5. ax.

5. The

5. The side KA is common, and $KL = KC$, and the angle $AKL = AKC$; *b* therefore g *constr.* $LA = CA$. But that no other line could be drawn equal to these, was proved above. Therefore, &c.


PROP. IX.



If in a circle BCK a point A be taken, and from that point more than two equal right lines AB, AC, AK , drawn to the circumference, then is that point A the center of the circle.

For a from no point without the center can more than two right lines equal be drawn to the circumference. Therefore A is the center. Which was to be dem.

PROP. X.




A circle $IAKBL$ cannot cut another circle $IEKFL$ in more than two points.

Let one circle, if it may be, cut the other in three points, I, K, L , and IK, KL being join'd, let them be bisected in M and N .

a Both circles have their centers in their perpendiculars MC, NH , and in the intersection of those perpendiculars, which is O . *b* Therefore the circles that cut each other have the same center. Which is false, by Prop. 5. 3.

PROP.

PROP. XI.




If two circles *GAD*, *FBC* touch one the other inwardly, and their centers be taken *G*, *F*; a right line *FG* joining their centers, and produced, shall cut the circumference in *A* the point of contact of the circles.

If it can be, let the right line *FG* produced cut the circles in some other point than *A*. so that not *FGA*, but *FGDB* shall be a right line.

a 15. def. 1. Let the line *GA* be drawn. Now, because *GD* *a* $\perp$ *GA*, and *GB* *b* $\perp$ *GA* (since the right line *FGB* passes through *F* the center of the greater circle) therefore is *GB* $\parallel$ *GD*. *c* Which is absurd.

b 7. 3.
c 9. ax.

PROP. XII.



If two circles *ACD*, *BCE* touch one the other outwardly, the right line *AB* which joins their centers *A*, *B*, shall pass thro' the point of contact *C*.


If it may be, let *ADEB* be a right line cutting the circles not in the point of contact *C*, but in the points *D*, *E*; draw *AC*, *CB*. then is *AD* + *EB* (*AC* + *CB*) *a* $\neq$ *ADEB*. *b* Which is absurd.

a 20. 1.

b 9. ax.

PROP.

PROP, XIII.




A circle CAF cannot touch a circle BAH in more points than one A, whether it be inwardly or outwardly.

1. Let one circle (if it can be) touch another in two points A, H, *a* 11. 3. Then will the right line CB that joins the

centers, if produced, fall as well in A, as H. Now because $CH = CA$, and $BH = CH$, therefore is $BA (= BH) = CA$. *d* Which is absurd. *c* 15. def. 1.

2. If it be said to touch outwardly in the points E and F, then draw the line EF, *e* which will be in both circles, Therefore those circles cut one the other; Which is against the Hyp.

PROP. XIV.




In a circle EABC equal right lines AC, BD are equally distant from the center E: and right lines AC, BD which are equally distant from the center, are equal among themselves.

From the center E draw the perpendiculars EF, EG. *a* which will bisect the lines AC, BD. join EA, EB. *a* 3. 3. 1. Hyp. $AC = BD$. therefore $AF = BG$. But $AE = BE$. *b* 7. ax. *f* 9

so $EA = EB$. therefore $EFq\ c = EAq - AFq =$
 $c\ 47.1. \& EBq - BGq\ c = EGq. \& \text{Therefore } FE = EG.$
 $3. ax. \quad 2. Hyp. EF = EG. \text{Therefore } AFq\ c = EAq - EFq$
 $d\ sch. 48.1. = EBq - EGq = BGq. \text{Therefore } AF\ d = GB, \text{ and}$
 $e\ 6. ax. \quad e\ \text{consequently } AC = BD. \text{ Which was to be dem.}$

PROP. XV.



a 15. def. 1.
b 20. 1.


In a circle $GABC$ the greatest line is AD the diameter; and of all other lines, that line FE , which is nearest to the center G is greater than any line BC farther distant from it.

1. Draw GB and GC . The diameter AD (a $GB + GC$) $b = BC$.

2. Let the distance GI be $c = GH$. Take $GN = GH$. Thro' the point N draw KL perpendicularly to GI : Join GK, GL . Because $GK = GB$, and $GL = GC$, and the angle $KGL = BGC$; c therefore is KL (FE) $c = BC$. Which was to be dem.

c 24. 1.

PROP. XVI.



A line CD drawn from the extreme point of the diameter HA of a circle $BALH$, perpendicular to the said diameter, shall fall without the circle; and between the same right line and the circumference cannot be drawn another line AL . And the angle of

of the semicircle BAI , is greater than any right-lined acute angle BAL ; and the remaining angle without the circumference DAI is less than any right-lined angle.

1. From the center B to any point F in the right line AC , draw the right line BF . The side BF subtending the right angle BAF is *a* greater 19. 1. than the side BA which is opposite to the acute angle BEA . Therefore whereas BA (BG) reaches to the circumference, BF shall reach further; and so the point F . and for the same reason is any other point of the line AC placed without the circle.

2. Draw BE perpendicular to AL . The side BA opposite to the right angle BEA is *b* greater 19. 1. than the side BE which subtends the acute angle BAE ; therefore the point E , and so the whole line EA falls within the circle.

3. Hence it follows that any acute angle, to wit, EAD , is greater than the angle of contact DAI , and that any acute angle BAL is less than the angle of a semicircle BAI . Which was to be dem.


Coroll.

Hence, A right line drawn from the extremity of the diameter of a circle, and at right angles, is a tangent to the said circle.

From this Proposition are gathered many Paradox and wonderful Confectaries, which you may meet with in the Interpreters.

P R O P. XVII.

From a point given A to draw a right line AC which shall touch a circle given DBC .




From D the center of the circle given let a line DA cutting the circumference in B , be drawn to the point given A ; from the center D describe another circle thro' the point A ; and from B draw a perpendicular to AD , which shall meet with the circle AE in the

The third Book of

the point E; and draw ED meeting with the circle BC in the point C. Then the line drawn from A to C shall touch the circle DBC.

- a 15. def. 1. For DB $a = DC$, and DE $a = DA$, and the
 b 4. 1. angle D is common; b therefore the angle ACD
 c cor. 16. 3. EBD and right. c Therefore AC touches the circle in C. Which was to be done.

PROP. XVIII.




If any right line AB touch a circle FEDC, and from the center to the point of contact E a right line FE be drawn; that line FE shall be perpendicular to the tangent AB.

If you deny it, let some other line FG be drawn from the center F perpendicular to the tangent, and

- a 2. def. 3. a cutting the circle in D. Therefore, whereas
 c 16. 3. the angle FGE is said to be right, b thence is
 b cor. 17. 1. the angle FEG acute; c so that FE (FD) $\neq$
 c 19. 1. FG. Which is absurd.
 d 9. ax.

PROP. XIX.



If any right line AB touch a circle, and from the point of contact C a right line CE be erected at right angles to the tangent, the center of the circle shall be in the line CE so erected.

If you deny it, let the center be without the line

- * 18. 3. CE in the point F; and from F to the point of
 a 11. ax. contact let FC be drawn. Therefore the angle
 b 9. ax. *FCB is right, and a consequently equal to the
 angle ECB, which was right by Hyp. b Which
 is absurd.

PROP.


PROP. XX.



In a circle $DABC$, the angle BDC at the center is double of the angle BAC at the circumference, when the same arch of the circle BC is the base of the angles.

Draw the diameter ADE . The outward angle BDE $a = DAB + DBA$ $b = 2 DAB$. Likewise a 32. 1. the angle $EDC = 2 DAC$. Therefore in the first b 5. 1. case the whole angle $c BDC = 2 BAC$. and in c 2. ax . the third case the d remaining angle $BDC = 2 d$ 20. ax . BAC . Which was to be dem.

PROP. XXI.




In a circle $EDAC$, the angles DAC and DBC which are in the same segment, are equal one to the other.

1. Case. If the segment $DABC$ be greater than a semicircle, from the center E draw ED , EC . Then is twice the angle A $a = E a = 2 B$. $W.W.$ to be dem. a 20. 3.

2. Case. If the segment be less than a semicircle, then is the sum of the angles of the triangle ADF equal to the sum of the angles of the triangle BCF . from each let AFD be taken away b equal to b 15. 1. BFC ,

c By the 1. BFC, and ADB $c = ACB$ be likewise taken away, then remains $DAC = DBC$. *W. W.* to be dem.

PROP. XXII.



The angles ADC , ABC of a quadrilateral figure $ABCD$ described in a circle, which are opposite one to the other, are equal to two right angles.

Draw AC , BD . The angle $ABC + BCA + BAC = 2$ right. But $BDA =$

a 32. 1.

b 21. 3.

c 1. *ax.*

BCA , and $BDC = BAC$. *c* Therefore $ABC + ADC = 2$ right angles. Which was to be dem.


Coroll.

* See the following Diagr.

1. Hence, If one side AB of a quadrilateral described in a circle be produced, the external angle EBC is equal to the internal angle ADC , which is opposite to that ABC which is adjacent to EBC , as appears by 13. 1. and 3. *ax.*

2. A circle cannot be described about a Rhombus; because its opposite angles are greater or less than two right angles.

Schol.



If in a quadrilateral $ABCD$ the angles A and C , which are opposite, be equal to two right, then a circle may be described about that quadrilateral.

For a circle will pass thro' any three angles (as shall appear by 5. 4.) I say that shall pass thro' A the fourth also of such

a quadrilateral: For if you deny it, let the circle pass thro' F. Therefore the right lines BF, FD, BD being drawn, the angle $C + F = 2a$ 22. 3. right $b = C + A$ wherefore, A is equal to F . b hyp. *Which is absurd.*

c 3. ax.
d 21. 1.

PROP. XXIII.




Two like and unequal segments of circles ABC , ADC cannot be set on the same right line AC , and the same side thereof.

For if they are said to be like, draw the line CB cutting the circumference in D and B , join AB and AD . Because the segments are supposed like, a therefore is the angle $ADC = a$ 10. def. 3. ABC . b Which is absurd.

b 16. 1.

PROP. XXIV.



Like segments of circles ABC , DEF upon equal right lines AC , DF , are equal one to the other.

The base AC being laid on the base DF will agree with it, because $AC = DF$. Therefore the segment ABC shall agree with the segment DEF (for otherwise it shall fall either within or without, and if so a then the segments are not like, a 23. 3. which is contrary to the Hypothesis, and at least it shall fall partly within and partly without, and so cut in three points, b which is absurd. c b 10. 3. Therefore the segments $ABC = DEF$. Which c 8. ax. was to be dem.

E

PROP.

P R O P. XXV.




A segment of a circle ABC being given, to describe the whole circle whereof that is a segment.

Let two right lines be drawn AB, BC, which bisect in the points D and E. From D and E draw the perpendiculars DF, EF meeting in the point F. I say this point shall be the center of the circle.

a cor. 1. 3. For the center shall be as well in a DF as; FE, therefore it must be in the common point F. Which was to be done.

P R O P. XXVI.



In equal circles GABC, HDEF, equal angles stand upon equal parts of the circumference, AC, DF; whether those angles be made at the centers, G, H, or at the circumferences, B, E.

Because the circles are equal, therefore is $GA = HD$, and $GC = HF$; also by Hypothesis the angle $G = H$; a therefore $AC = DF$.

b 20. 3. Moreover the angle $Bb = \frac{1}{2} Gc = \frac{1}{2} Hb = E$.
 c hyp. d Therefore the segments ABC, DEF are like,
 d 10. def. 3. and e consequently equal, f whence the remain-
 e 24. 3. ing segments also AC, DF are equal. Which
 f 3. ax. was to be dem.

Coroll.

Coroll.



In a circle $ABCD$ let an arch AB be equal to DC ; then shall AD be parallel to BC . For the right line AC being drawn, the angle $ACB = CAD$; wherefore by 27. 1. the said sides are parallel.

PROP. XXVII.




In equal circles $GABC$, $HDEF$, the angles standing upon equal parts of the circumference AC , DF , are equal between themselves,

whether they be made at the centers G , H , or at the circumferences, B , E .

For if it be possible, let one of the angles AGC be $\neq$ DHF , and make $AGI = DHF$. thence is the arch $AI = DF = AC$. Which is absurd.

a 26. 3:
b hyp.
c 9. ax.

Schol.



A right line EF , which being drawn from A the middle point of any periphery BC , toucheth the circle, is parallel to the right line BC subtending the said periphery.

From the center D draw a right line DA to the point of contact A , and join DB , DC .


The side DG is common, and $DB = DC$, and the angle $BDA = CDA$, because the arches BA , CA

E 2

CA

b hyp. CA are *b* equal) therefore the angles at the base
 c 4. 1. DGB, DGC are *c* equal, and *d* consequently
 d 10. def. 1. right; But the inward angles GAE, GAF are
 e hyp. also *e* right, *f* therefore BC, EF are parallel.
 f 28. 1. Which was to be dem.

PROP. XXVIII.




In equal circles
 GABC, HDEF,
 equal right lines AC,
 DF cut off equal parts
 of the circumference,
 the greatest ABC e-
 qual to the greatest

DEF, and the least AIC to the least DKF.

From the centers G, H draw GA, GC, & HD, HF.
 a hyp. Because GA = HD, and GC = HF; and AC =
 b 8. 1. DF, *b* therefore is the angle G = H; *c* whence
 c 26. 3. the arch AIC = DKF; *d* and so the remaining
 d 3. ax. arch ABC = DEF. Which was to be dem.

But if the subtended line AC be $\square$ or $\rhd$ than
 DF, then in like manner will the arch AC be
 $\square$ or $\rhd$ than DF.

PROP. XXIX.



In equal circles
 GABC, HDEF, equal
 right lines AC, DF
 subtend equal periph-
 eries ABC, DEF.

Draw the lines
 GA, GC, and HD, HF. Because GA = HD,
 and GC = HF, and (because the arches AC, DF
 are *a* equal) the angle G *b* = H. *c* therefore is
 the base AC = DF. Which was to be dem.

a hyp.
 b 27. 3.
 c 4. 1.

This and the three precedent Propositions may
 be understood also of the same circle.

PROP.

PROP. XXX.

To cut a periphery given ABC into two equal parts.




Draw the right line AC , and bisect it in D ; from D draw a perpendicular DB meeting with the arch in B , it shall bisect the same.

For join AB , and CB . The side DB is common, and $AD = DC$, and the angle $ADB = CDB$. c therefore $AB = BC$; d whence the arch $AB = BC$. Which was to be done.

a constr.
 b 12. ax.
 c 4. 1.
 d 28. 3.

PROP. XXXI.



In a circle, the angle ABC , which is in the semicircle, is a right angle; but the angle, which is in the greater segment BAC is less than a right angle; and the angle which is in the lesser segment BFC is greater than a right angle.


Moreover, the angle of the greater segment is greater than a right angle, and the angle of the lesser segment is less than a right angle.

From the center D draw DB , because $DB = DA$, therefore is the angle $A = DBA$, and the angle $DCB = DBC$, therefore the angle $ADC = A + ACB = EBC$, so that ABC and EBC are right angles. $W. W.$ to be dem. Therefore BAC is an acute angle. $W. W.$ to be dem. And BFC is an obtuse angle. Lasty, the angle contained under the right line CB , and the arch BAC is greater than the right angle ABC ; but the angle made by the right line CB and the periphery of the lesser segment BFC is less than the right angle ABC . Which was to be dem.

Schol.

In a right-angled triangle ABC , if the Hypotenuse (or subtended line) AC be bisected in D , a circle drawn from the center D thro' the point A shall also pass thro' the point B : As you may easily demonstrate from this Prop. and 21. 1.

PROP. XXXII.




If a right line AB touch a circle, and from the point of contact be drawn a right line CE cutting the circle, the angles ECB , ECA which it makes with the tangent line are equal to those angles EDC , EFC which are made in the alternate segments of the circle.

Let CD , the side of the angle EDC , be perpendicular to AB (a for it's to the same purpose) b therefore CD is the diameter. c therefore the angle CED in a semicircle is a right angle, d and therefore the angle $D + DCE =$ to a right angle $e = ECB + DCE$. f Therefore the angle $D = ECB$. Which was to be dem.

Now whereas the angle $ECB + ECA g = 2$ right $h = D + F$, from both of these take away ECB and D , which are equal, k then remains $ECA = F$. Which was to be dem.

PROP. XXXIII.




Upon a right line AB to describe a segment of a circle $AIEB$ which shall contain an angle AIB equal to a right lined angle given C .

a Make the angle BAD

223. 1.

BAD = C. Thro' the point A draw the line AE perpendicular to HD. At one end of the line given AB make an angle ABF = BAF, one side whereof let it cut the line AE in F; from the center F thro' the point A, describe a circle which shall pass thro' B (because the angles FBA b = FAB. and c therefore FB = FA,) AIB b constr. is the segment sought. For because HD is per- c 6. 1. pendicular to the diameter AE, it therefore d cor. 16. 3. touches the circle HD which AB cuts. And e 32. 3. therefore the angle AIB e = BAD f = C, Which f constr. was to be done.

PROP. XXXIV.




From a circle given ABC to cut off a segment ABC containing an angle B equal to a right-lined angle given D.

a Draw a right line a 17. 3. EF which shall touch the b 23. 1. circle given in A. b let c 32. 1.

AC be drawn also making an angle FAC = D. d constr. This line shall cut off ABC containing an angle B c = CAF d = D. Which was to be done,

PROP. XXXV.



If in a circle FB CA two right lines AB, DC cut each other, the rectangle comprehended under the segments AE, EB of the one, shall be equal to the

The third Book of

the rectangle comprehended under the segments CE, ED of the other.

1. *Case.* If the right lines cut one the other in the center, the thing is evident.

2. *Case.* If one line AB pass thro' the center F, and bisect the other line CD, then draw FD.

a 5. 2. Now the rectangle AEB + FEq a FBq b = FDq c
b sch. 48. 1. = EDq + FEq d = CED + FEq. e Therefore the
c 47. 1. rectangle AEB = CED. Which was to be dem.

d hyp. 3. *Case.* If one of the lines AB be the diameter, and cut the other line CD unequally, bisect CD by FG a perpendicular from the center.

e 3. ax. The rectangle AEB + FEq.
f FBq (FDq).
f 5. 2. These
g 47. 1. are e-
h 5. 2. qual. g FGq + GDq.
k 47. 1. EGq + b GEq + Rectang. CED.
l 3. ax. b FEq + CED.

Therefore the Rectangle AEB = CED.


4. *Case.* If neither of the right lines AB, CD pass thro' the center, then thro' the point of intersection E, draw the diameter GH. By that which hath been already demonstrated, it appears that the rectangle AEB = GEH = CED. Which was to be dem.

More easily, and generally, thus; join AC and BD, then because the angles a CEA, DEB, and b also C, B (upon the same arch AD) are equal, thence are the triangles CEA BED c equiangular. d Wherefore CE, EA :: EB, ED. and e consequently CE x ED =

AE x EB. Which was to be dem.

The citations out of the sixth Book, both here and in the following Prop. have no dependance upon the same; so that it was free to use them.

PROP.



a 15. 1.


b 21. 3.

c cor. 31. 1.

d 4. 6.

e 16. 6.

PROP. XXXVI.



If any point *D* be taken without a circle *EB C*, and from that point two right lines *DA*, *DB* fall upon the circle, whereof one *DA* cut the circle, the other *DB* touches it, the rectangle comprehended under the whole line *DA* that cuts the circle, and under *DC* that part which is taken from the point given *D* to the convex of the periphery, shall be equal to the square made of the tangent line.

1. Case. If the secant *AD* pass thro' the center, then join *EB*, this *a* will make a right angle with the line *DB*, wherefore *DBq + d EBq b* 47. 1. *(ECq) b = EDq c = AD × DC + ECq*. Therefore *AD × DC = DBq*. Which was to be dem. d 3. ax.

2. Case. But if *AD* pass not thro' the center then draw *EC*, *EB*, *ED*, and *EF* perpendicular to *AD*, *a* wherefore *AC* is bisected in *F*. a 3. 3.

Because *BDq + EBq b = DEq b = EFq + b* 47. 1. *FDq c = EFq + ADC + FCq d = ADC + c* 6. 2. *CEq (EBq.) e* Therefore is *BDq = ADC*. Which d 47. 1. was to be dem. e 3. ax.


More

a 32. 3.

b 32. 1.

c 4. 6.

d 17. 6.



More easily, and generally, thus; Draw AB and BC. Then because the angles A, and DBC *a* are equal, and the angle D common to both, thence are the triangles BDC, ADB *b* equiangular, *c* Wherefore $AD \cdot DB :: DB \cdot CD$. and *d* consequently $AD \times DC = DB^2$. Which was to be dem.

Coroll.

1. Hence, If from any point A taken without a circle, there be several lines AB, AC drawn which cut the circle, the rectangles comprehended under the whole lines AB, AC, and the outward parts AE, AF are equal between themselves.

For if the tangent AD be drawn, then is $CAF = AD^2 = BAE$.


2. It appears also from hence, that if two lines AB, AC drawn from the same point do touch a circle, those two lines are equal one to the other.

For if AE be drawn cutting the circle, then is $AB^2 = EAF = AC^2$.


3. It is also evident that from a point A taken without a circle, there can be drawn but two lines AB, AC that shall touch the circle.

For if a third line AD be said to touch the circle, thence is $AD^2 = AB^2 = AC^2$. Which is absurd.

4. And



a 36. 3.



a 36. 3.

b 36. 3.

c 2. cor.


d 8. 3.

4. And on the contrary, it is plain, that if two equal right lines AB, AC fall from any point A upon the convex periphery of a circle, and that if one of these equal lines AB touch the circle, then the other AC touches the circle also.

For if possible, let not AC, but another line AD, touch the circle; therefore is $AD = AC$ *e 2. cor.*
 $f AB.$ *g Which is absurd.* *f hyp.*

P R O P. XXXVIII.

g 8. 3.



If without a circle EBF any point D be taken, and from that point two right lines DA, DB fall on the circle, whereof one line DA cuts the circle the other DB falls upon it; and if also the rectangle comprehended under the whole line that cuts the circle, and under that part of it DC which is taken betwixt the point D and the convex periphery, be equal to that square which is made of the line DB falling on the circle, I say that the line DB so falling shall touch the circle given.

From that point D a let a tangent DF be *a 17. 3.*
drawn, and from the center E draw ED, EB, EF.
Now because $DB = ADC = DF$, there- *b hyp.*
fore is $DB = DF$: But $EB = EF$, and the side *c 36. 3.*
ED common; *e* therefore the angle $EBD = EFD$. *d 1. ax. &*
but EFD is a right angle, and *f* therefore EBD *sch. 4. 1.*
is right also, and *g* therefore DB touches the *e 8. 1.*
circle. *Which was to be dem.*

g cor. 16. 3.
h 8. 1.

Coroll.

From hence it follows that the *h* angle $EDB = EDF$.


The End of the third Book.

THE

THE FOURTH BOOK OF EUCLIDE'S ELEMENTS.

Definitions.


I. **A** Right-lined figure is said to be inscribed in a right-lined figure, when every one of the angles of the inscribed figure touch every one of the sides of the figure wherein it is inscribed.



So the triangle DEF is inscribed in the triangle ABC.

II. In like manner a figure is said to be described about a figure, when every one of the sides of the figure circumscribed touch every one of the angles of the figure about which it is circumscribed.

So the triangle ABC is described about the triangle DEF.



III. A right-lined figure is said to be inscribed in a circle, when all the angles of that figure which is inscribed do touch the circumference of the circle.

IV. A right-lined figure is said to be described about a circle, when all the sides of the figure which

which is circumscribed touch the periphery of the circle.


V. After the like manner a circle is said to be inscribed in a right-lined figure, when the periphery of the circle touches all the sides of the figure in which it is inscribed.

VI. A circle is said to be described about a figure, when the periphery of the circle touches all the angles of the figure, which it circumscribes.



VII. A right line is said to be co-opted or applied in a circle when the extremes thereof fall upon the circumference; as the right line AB .

PROP. I. *Probl. 1.*



In a circle given ABC to apply a right line AB equal to a right line given D , which doth not exceed AC the diameter of the circle.


From the center A by the space $AE = D$

a describe a circle meeting with the circle given a 3. post. & in B , draw AB . Then is $AB = AE = D$. 3. 1.

Which was to be done.

b 15. def. 1.
c constr.

PROP. II. *Probl. 2.*




In a circle given ABC to describe a triangle ABC , equiangular to a triangle given D . EF .

Let the right line

a 17. 3. line GH *a* touch the circle given in A; *b* make
 b 23. 1. the angle $HAC = E$, *b* and the angle $GAB = F$. then join BC; and the thing is done.

c 32. 3. For the angle $Bc = HAC d = E$, and the
 d *constr.* angle $Cc = GAB d = F$; *e* whence also the
 e 32. 1. angle $BAC = D$. Therefore the triangle BAC
 inscribed in the circle is equiangular to DEF.
Which was to be done.

PROP. III.




About a circle given IABC to describe a triangle LNM equiangular to a triangle given DEF.

Produce the side EF on both sides; at the
 a 23. 1. center I *a* make an angle $AIB = DEG$, and an
 b 17. 3. angle $BIC = DFH$. Then in the points A, B, C
 let three right lines LN, LM, NM *b* touch
 the circle, and the thing is done.

For it's evident that the right lines LN, LM,
 c 13. 1. MN will meet and make a triangle, *b* because
 d 11. 3. the angles LAI, LBI are right; so that the *d*
 right line AB produced will make the angles
 LAB, LBA, less than two right angles.

Since, therefore the angle $AIB + Le = 2$
 a *sch.* 32. 1. right angles *f* $= DEG + DEF$, and $AIB g =$
 f 13. 1. DEG ; *b* therefore is the angle $L = DEF$. By
 g *constr.* the like way of argument the angle $M = DEF$.
 h 3. *ax.* *k* therefore also the angle $N = D$. And there-
 k 32. 1. fore the triangle LNM described about the cir-
 cle is equiangular to EDF the triangle given.
Which was to be done.

PROP. IV.



In a triangle given *ABC*, to describe a circle *EFG*.

a Bisect the angles *B* and *C* with the right lines *BD*, *CD* meeting in the point *D*, *b* and draw the perpendiculars *DE*, *DF*, *DG*. A circle de-

scribed from the center *D* thro' *E*, will pass thro' *G* and *F*, and touch the three sides of the triangle.

For the angle *BDE* $\angle = \angle DBF$; and the angle *c* *constr.* *DEB* $\angle = \angle DFB$; and the side *DB* common. *ed* 12. *ax.* therefore *DE* $=$ *DF*. By the like argument *e* 26. 1. *DG* $=$ *DF*. The circle therefore described from the center *D* passes thro' the three points *E*, *F*, *G*. and whereas the angles at *E*, *F*, *G* are right, therefore it touches all the sides of the triangle. Which was to be done.


Schol.

Hence, The sides of a triangle being known, their segments which are made by the touchings of the circle inscribed shall be found, thus;

Let *AB* be 12, *AC* 18, *BC* 16, then is *AB* + *BC* $=$ 28. Out of which subduct 18 $=$ *AC* $=$ *AE* + *FC*, then remains 10 $=$ *BE* + *BF*. Therefore *BE*, or *BF* $=$ 5; and consequently *FC*, or *CG*, $=$ 11. Wherefore *GA*, or *AE*, $=$ 7.

PROP.

PROP. V.



About a triangle given ABC to describe a circle $FABC$.

a 10. and *a* Bisect any two sides BA, CA with a perpendicular DF, EF meeting in the point F . I say this shall be the center of the circle.

b constr. For, let the right lines FA, FB, FC be drawn. Now because $AD = DB$ and the side DF common, and the angles $FDA = FDB$, therefore is $FA = FB$. After the same manner is $FC = FA$. Therefore a circle described from the center F shall pass thro' the angles of the triangle given (*viz.*) B, A, C . Which was to be done.

Coroll.


*** 31. 3.** * Hence, if a triangle be acute-angled, the center shall fall within the triangle; if right-angled, in the side opposite to the right angle, and if obtuse-angled, without the triangle.

Schol.

By the same method may a circle be described, that shall pass thro' three points given, not being in the same strait line.

PROP.

PROP. VI.




In a circle given
E ABCD to inscribe a
square ABCD.

a Draw the diameters a 11. 1.
AC, BD cutting each
other at right angles in
the center E. Join the
extremes of these diame-
ters with the right lines
AB, BC, CD, DA. And
the thing is done.

Now because the four angles at E are right, the
arcs and c subtended lines AB, BC, CD, DA b 26. 3.
are equal; therefore is the figure ABCD equi c 29. 3.
lateral, and all the angles in semicircles, and so d 31. 3.
right. e Therefore ABCD is a square inscri e 29. 3. def.
ed in a circle given. Which was to be done. 1.

PROP. VII.



About a circle given
EABCD to describe a
square FHIG.


Draw the diameters
AC, BD cutting one
the other at right an-
gles; thro' the extremes
of these diameters a a 17. 3.
draw tangents meeting

in F, H, I, G, then I say it's done.
For because b the angles A and C are right, c b 18. 3.
therefore is FG parallel to h i. After the same c 18. 1.
manner is FH parallel to GI. and therefore FHIG
is a Pgr. and also right-angled. It is equilateral
because FG d = HI d - D b e CA d FH d = d 34. 1.
GI. Wherefore FHIG is a f square described e 15. def. 1.
about the circle given. Which was to be done. f 29. def. 1.

F.


Schol.

Schol.



A square ABCD described about a circle is double of the square EFGH inscribed in the same circle. For the rectangle HB = DHEF, and $HD = 2 HGF$ by the 41. 1.

PROP. VIII.



In a square given ABCD to inscribe a circle IEFHG.

Bisect the sides of the square in the points H, E, F, G, cutting one the other in I. A circle drawn from the center I through H shall

be inscribed in the square.

For because AH and BF are *a* equal and *b* parallel, *c* therefore is AB parallel to HF parallel to DC. After the same manner is AD parallel to EG, parallel to BC; therefore IA, ID, IB, IC are parallelograms. Therefore $AH = EI = FI = IG$. The circle therefore described from the center I thro' H shall pass thro' H, E, F, G, and touch the sides of the square, being the angles H, E, F, G, are right. Which was to be done.

a 7. ax.

b hyp.

c 33. 1.


d 7. ax.

e 34. 1.

PROP.

AC
circ
AC,
requ

PROP. IX.




About a square given
ABCD to describe a
circle EABCD

Draw the diameters
AC, BD cutting one
the other in E. From
the center E thro' A
describe a circle, then
I say that circle is
described about the
square.

For the angles ABD and BAC are a half of a 4. Cor.
right angles, b therefore $EA = EB$. After the 33. r.
same manner is $EA = ED = EC$. The circle b 6. 4.
therefore described from the center E passes
thro' A, B, C, D the angles of the square given.
Which was to be done.

PROP. X.



To make an
Isosceles triangle
ABD, having
each angle at the
base B and ADB
double to the
remaining angle
A.

Take any right
line AB, and di-
vide it in C, a so a 11. 2.
that $AB \times BC$
may be equal to

ACq. From the center A thro' B, describe the
circle ABD; and in this circle b apply $BD = b$ 1. 4.
AC, and join AD; I say ABD is the triangle
required.

F 2


For

For, draw DC, and thro' the points C, D, A.
 e 5. 4. c draw a circle. Now because $AB \times BC = ACq$, d
 d 37. 3. it is evident that FD touches the circle ACD
 e 32. 3. which CD cutteth; e therefore is the angle BDC
 f 2. ax. $= A$, and therefore the angle $BDC + CDA f =$
 g 32. 1. $A + CDA g = BCD$. But $BDC + CDA = BDA$
 h 5. 1. $b = CBD$. k therefore the angle $BCD = CBD$,
 k 1. ax. and therefore $DC l = DB = m AC$, n wherefore
 l 6. 1. the angle $CDA = A = BDC$. therefore ADB
 m constr. $= 2 A = ABD$. Which was to be done.
 n 5. 1.

This construction is Analytically found out
 thus; take the thing for done, and let the right
 line DC bisect the angle BDA; a therefore DA,
 a 3. 6. DB :: CA, CB. also because the angle $CDA b =$
 b constr. $\frac{1}{2} ADB c = A$, c therefore $CA = DC$. and be-
 c hyp. cause the angle $DCB b = A + CDA = 2 A e =$
 d 6. 1. B, d thence will be $DB = DC$. f from whence
 e 31. 1. also $DB = CA$. and so $DA(BA.)CA :: CA.CB$. g
 f 2. ax. whence $BA \times CB = CAq$.
 g 17. 6. Coroll.

Whereas all the angles A, B, D b make up
 h 32. 1. two right angles, it's evident that A is $\frac{1}{2}$ of two
 right angles.

PROP. XI.



In a circle given $ABCDE$ to describe a Pentagone
 figure $ABCDE$ equilateral and equiangular.

a 10. 4. a Describe an Isosceles triangle FGH , having
 b 2. 4. each angle at the base double to the other; b in-
 scribe

scribe a triangle CAD equiangular to the said triangle FGH. *c* Bisect the angles at the base *c* 9. 1. ACD and ADC with the right lines DB, CE meeting with the circumference in B and E. join the right line CB, BA, AE, ED. Then I say it is done.

For it is evident by construction that the angles CAD, CDB, BDA, DCE, ECA, are equal; wherefore the *d* arches and *e* subtended lines DC, *d* 26. 3. CB, BA, AE, DE are equal. Therefore the Pen- *e* 29. 3. tagone is equilateral, and equiangular *f* because *f* 27. 3. the angles of it BAE, AED, &c. stand on equal *g* arches BCDE, ABCD, &c.


A more easy Practice of this Problem shall be deliver'd at 10, 13. *g* 2. ax.

Coroll.

Hence, each angle of an equilateral and equiangular Pentagone is equal to $\frac{3}{5}$ of two right angles, or $\frac{6}{5}$ of one right angle.


Schol.

Generally all figures of odd number of sides are *Pet. Herig.* inscribed in circles by the help of Isosceles triangles, whose angles at the base are multiples of those at the top: and figures of even number of sides are inscribed in a circle by the help of Isosceles triangles, whose angles at the base are multiples sesquialter of those at the top.



As in the Isosceles triangle CAB, if the angle $A = \frac{1}{3} C = B$, then will AB be the side of a Heptagone. If $A = \frac{1}{4} C$, then is AB the side of Enneagone. But if $A = \frac{1}{5} C$, then is AB the side of a square. And if $A = \frac{1}{6} C$ CAB will subtend the sixth part of a circumference: and likewise if $A = \frac{1}{7} C$, then will AB be the side of Octagone.

The fourth Book of
PROP. XII.



About a circle given $FABCDE$, to describe an equilateral and an equiangular pentagone $HIKLG$.

a II. 4.

b cor. 16.

13.

c 2. cor. 36.

3.

d 8. 1.

e 27. 3.

f 7. ax.

g 12. ax.

h 26. 1.


k 3. ax.

a Inscribe a pentagone $ABCDE$ in the circle given; and from the center draw the right lines FA, FB, FC, FD, FE ; and to those lines draw so many perpendiculars GAH, HBI, ICK, KDL, LEG meeting in the points H, I, K, L, G . then I say it is done. For because GA, GE from the same point G touch the circle, therefore is $GA = GE$, and therefore the angle $GFA = GFE$, therefore the angle $AFE = 2 GFA$. After the same manner is the angle $AFH = FHB$, and consequently the angle $AFB = 2 AFH$. But the angle $AFE = AFB$, therefore the angle $GFA = AFH$. But also the angle $FAH = FAG$, and the side FA is common, therefore $HA = AG = GE = EL$, &c. Therefore HG, GL, LK, KI, IH the sides of the pentagone are equal, the angles also are, because double of the equal angles AGF, AHF . therefore, &c.

Coroll.

After the same manner, if any equilateral and equiangular figure be described in a circle, and at the extreme points of the semi-diameters drawn from the center to the angles, be drawn perpendicular lines to the said diameters, I say that these

perpen
many
about




F dra
Then
G w
D
the f
d the
but t
fore
man
Now
angl
ther
righ
a cir
thro
of
poin

F
tera
tha
the
ing
figu

perpendiculars shall make another figure of as many equal sides and equal angles, described about the circle.

PROP. XIII.



In an equilateral and equiangular pentagon given $ABCDE$, to inscribe a circle $FGHK$.

a Bisect two angles a 9. 1. of the pentagon A and B with the right lines AF , BF meeting in the point F . From F draw the perpendiculars FG , FH , FI , FK , FL . Then a circle described from the center F thro' G will touch all the sides of the pentagon.

Draw FC , FD , FE . Because BA *b* $= BC$, and *b* *hyp.* the side BF common, and the angle FBA *c* $= FBC$, *c* *constr.* therefore is $AF = FC$ and the angle $FAB = FCB$, *d* 4. 1. but the angle FAB *e* $= \frac{1}{2} BAE = \frac{1}{2} BCD$. There- *e* *hyp.* fore the angle $FCB = \frac{1}{2} BCD$. After the same manner are all the whole angles C , D , E bisected. Now whereas the angle FGB *f* $= FHB$, and the *f* 12. *a* angles $FBH = FBG$ and the side FB is common. *g* 26. 1. therefore is $FG = FH$. In like manner are all the right lines FH , FI , FK , FL , FG equal. Therefore a circle described from the center F thro' G passes thro' the points H , I , K , L and *h* touches the sides *h* *cor.* 16. 3. of the pentagon, because the angles at those points are right. Which was to be done.

Coroll.

Hence, If any two nearest angles of an equilateral and equiangular figure be bisected, and from that point in which the lines meet that bisect the angles be drawn right lines to the remaining angles of the figure, all the angles of the figure shall be bisected.


E 4

Schol.

Schol.

By the same method shall a circle be inscribed in any equilateral or equiangular figure.

PROP. XIV.



About a pentagone given *ABCDE* equilateral and equiangular to describe a circle *FABCDE*.

Bisect any two angles of the pentagone with the right lines *AF*, *BF* meeting in the point *F*; the circle described from the center *F* thro' *A* shall be described about the pentagone.

a cor. 13. 4.
b 6. 1.


For let *FC*, *FD*, *FE* be drawn. *a* Then the angles *C*, *D*, *E* are bisected; *b* and therefore *FA*, *FB*, *FC*, *FD*, *FE* are equal; therefore the circle described from the center *F* passes thro' *A*, *B*, *C*, *D*, *E* all the angles of the pentagone, Which was to be dem.

Schol.

By the same art is a circle described about any figure which is equilateral and equiangular.

PROP.

PROP. XV.



In a circle given *GABCD-
EF* to inscribe an Hexagone
(or six-sided figure) equilateral
and equiangular *ABC DEF*.

Draw the diameter *AD* ;
from the center *D* through
the center *G* describe a cir-
cle cutting the circle given
in the points *C* and *E*.
Draw the diameters *CF*,
EB ; and join *AB*, *BC*, *CD*,
DE, *EF*, *FA*. Then I say
it's done.

For the angle *CGD* $a =$ a 32. 1.
of 2 right $a =$ *DGE* $b =$ *AGF* $b =$ *AGB*. b 15. 1.
Therefore *BGC* $=$ of 2 right *FGE* ; c cor. 13. 1.
therefore the d arches and e subtenses *AB*, *BC*, d 26. 3.
CD, *DE*, *EF* are equal. Therefore the Hexa- e 29. 3.
gone is equilateral, but it is equiangled also, ff 27. 3.
because all the angles of it stand upon equal
arches.

Coroll.

1. Hence, The side of an Hexagone inscribed
in a circle is equal to the semidiameter.
2. Hereby an equilateral triangle *ACE* may
very easily be described in a circle given.


Schol.

To make a true Hexagone upon a right line
given *CD*.

a Make an equilateral triangle *CGD* upon the *Andr*.
line given *CD* ; from the center *G* thro' *C* and *Tacq*.
D describe a circle. That circle shall contain the a 1. 1.
Hexagone made upon the given line *CD*.

PROP.

PROP. XVI.



In a circle given *AEBC* to inscribe a quindecagone (or fifteen sided figure) equilateral and equiangular.

a 11. 4.

b 3. 4.

a Inscribe an equilateral pentagone *AEFGH* in the circle given, and *b* also an equilateral triangle *ABC*. then I say *BF* is the side of the quindecagone required.

c constr.

For the arch *AB* is $\frac{1}{3}$ or $\frac{1}{15}$ of that periphery whereof *AF* is $\frac{2}{3}$ or $\frac{10}{15}$. therefore the remaining part *BF* is $\frac{1}{15}$ of the periphery; and therefore the quindecagone, whose side is *BF*, is equilateral; but it is equiangular also, *d* because all the angles insit on equal arches of a circle, where every one $\frac{1}{15}$ of the whole circumference. Therefore, &c.

d 27. 3.

Schol.

A circle is geometrically divided into parts. $\left\{ \begin{array}{l} 4, 8, 16, \&c. \text{ by } 6, 4, \text{ and } 9, 1. \\ 3, 6, 12, \&c. \text{ by } 15, 4, \text{ and } 9, 1. \\ 5, 10, 20, \&c. \text{ by } 11, 4, \text{ and } 9, 1. \\ 15, 30, 60, \&c. \text{ by } 16, 4, \text{ and } 9, 1. \end{array} \right.$

Any other way of dividing the circumference into any parts given, is as yet unknown, wherefore in the construction of ordinate figures, we are forced to have recourse to mechanick artifices, concerning which you may consult the Writers of practical Geometry.

THE

THE FIFTH BOOK OF EUCLIDE'S ELEMENTS.

Definitions.

I. **A** Part, is a magnitude of a magnitude, a less of a greater, when the less measures the greater.

II. Multiple is a greater magnitude in respect of a lesser, when the lesser measures the greater.

III. Ratio (or rate) is the mutual habitude or respect of two magnitudes of the same kind each to other, according to quantity.

In every ratio, that quantity which is referred to another quantity is called the antecedent of the ratio, and that to which the other is referred is called the consequent of the ratio, as in the ratio of 6 to 4, 6 is the antecedent, and 4 the consequent.

Note, The quantity of any ratio is known by dividing the antecedent by the consequent; as the ratio of 12 to 5 is expressed by $\frac{12}{5}$, or the quantity of the ratio of A

to B is $\frac{A}{B}$. Wherefore, often for brevity sake, we denote

the quantities of ratios thus; $\frac{A}{B} >$, or $=$, or $< \frac{C}{D}$.

that is, the ratio of A to B is greater, equal, or less than the ratio of C to D. And this note must be diligently observed in the understanding of the following Book.

Concerning the divers species of ratio's, you may please to consult Interpreters.

IV. Proportion is a similitude of ratio's.

That which is here termed Proportion, is more rightly called Proportionality or Analogy; for
pro

proportion commonly denotes no more than the ratio betwixt two magnitudes.

V. Those numbers are said to have a ratio betwixt them, which being multiplied may exceed one the other.

E, 12 | A, 5. B, 6. | G, 24. VI. Magnitudes are said to be in the same ratio, the first A to the

second C, and the third C to the fourth D, when the equimultiples E and F of the first A, and the third C compared with the equimultiples G, H of the second B and the fourth D, according to any multiplication whatsoever, either both together E, F are less than G, H both together, or equal taken together, or exceed one the other together, if those be taken E, G and F, H, which answer one to the other.

The note hereof is :: ; as A. B :: C. D. That is, as A is to B, so is C to D. which signifies that A to B, and C to D are in the same ratio. We sometimes thus express

$\frac{A}{B} = \frac{C}{D}$ that is A. B :: C. D.

VII. Magnitudes that have the same ratio (A. B :: C. D. are called proportional.

E, 30. | A, 6. B, 4. | G, 28. VIII. When of equimultiples, E the multiple of the first magnitude A exceeds G the multiple of the second B, but F the multiple of the third C exceeds not H the multiple of the fourth D, then the first A to the second B has a greater ratio than the third C to the fourth D.

If $\frac{A}{B} < \frac{C}{D}$, it is not necessary from this definition that E should always exceed G, when F is less than H; but it is granted that this may be.

IX. Proportionality consists in three terms at least. Whereof the second supplies the place of two.

X. When 3 magnitudes A, B, C are proportional,

nal, the
third C
a magn
A shall
what i
der one

Dup

is, the

Triple

the rat

or 2, 6

XI.

to ant

As if

logous

XII

antec

confe

or by p

In

given

by Ma

pends

med i

XI

taken

antec

theref

XI

and c

to th

fore b

XV

the a

pare

divisi

nal, the first A shall have a duplicate ratio to the third C of that it has to the second B: But when magnitudes A, B, C, D are proportional, the first A shall have a triplicate ratio to the fourth D of what it had to the second B; and so always in order one more, as the proportion shall be extended.

Duplicate ratio is thus expressed $\frac{A}{C} = \frac{A}{B}$ twice, that is, the ratio of A to C is double of the ratio of A to B.

Triple ratio is thus expressed; $\frac{A}{D} = \frac{A}{B}$ thrice. That is the ratio of A to D is triple of the ratio of A to B. $\therefore$ denotes continual proportionals; as A, B, C, D; or 2, 6, 18, 54, are $\therefore$.

XI. Magnitudes of a like ratio, are antecedents to antecedents, and consequents to consequents; As if $A:B :: C:D$. A and C; and B and D are homologous or magnitudes of a like ratio.

XII. Alternate proportion is the comparing of antecedent to antecedent, and consequent to consequent. As if $A:B :: C:D$. therefore alternately, or by permutation, $A:C :: B:D$. by the 16. of 5.

In this definition, and the 5 following, names are given to the six ways of arguing which are often used by Mathematicians: the force of which inferences depends on the Propositions of this Book, which are named in their Explications.

XIII. Inverse ratio is when the consequent is taken as the antecedent, and so compared to the antecedent as the consequent; As $A:B :: C:D$. therefore generally $B:A :: D:C$. by cor. 4. 5.

XIV. Compounded ratio is when the antecedent and consequent taken both as one are compared to the consequent it self. As $A:B :: C:D$. therefore by composition $A+B:B :: C+D:D$ by 18. 5.

XV. Divided ratio is when the excess wherein the antecedent exceeds the consequent, is compared to the consequent. As $A:B :: C:D$. therefore by division $A-B:B :: C-D:D$. by 17. 5.

XVI.

XVI. Converse ratio is when the antecedent is compared to the excess wherein the antecedent exceeds the consequent. *As* $A.B :: C.D$. therefore by converse ratio $A.A-B :: C.C-D$. by the coroll. of the 19. of the 5.

XVII. Proportion of equality is where there are taken more magnitudes than two in one order, and also as many magnitudes in another order, comparing two to two being in the same ratio; it comes to pass that as in the first order of magnitudes, the first is to the last, so in the second order of magnitudes is the first to the last. Or otherwise: it is a comparison of the extremes together, the mean magnitudes being taken away.

XVIII. Ordinate proportionality is, when as the antecedent is to the consequent, so is the antecedent to the consequent, and as the consequent is to any other, so is the consequent to any other. *As* $A.B :: D.E$. also $B.C :: E.F$. it shall be true also $A.C :: D.F$. by the 22. of the 5.

XIX. Inordinate proportion is, when three magnitudes being put, and others also, which are equal to these in multitude, as in the first magnitudes the antecedent is to the consequent, so in the second magnitude is the antecedent to the consequent; and as in the first magnitudes the consequent is to any other, so in the second magnitudes any other thing to the antecedent. *As* $A.B :: F.G$. also $B.C :: E.F$. it shall be true in inordinate proportion. $A.C :: E.G$. by the 23. of the 5.

XX. Any number of magnitudes being put; the proportion of the first to the last is compounded out of the proportions of the first to the second, the second to the third, and the third to the fourth, and so forwards till the proportion arise.

Let there be any number of magnitudes $A, B, C,$


D . by this definition $\frac{A}{D} = \frac{A}{B} + \frac{B}{C} + \frac{C}{D}$.

Axioms.

Axioms.

Magnitudes equimultiples to the same multiple, are also equimultiples betwixt themselves.

PROP. I.




If there be a number of magnitudes how many soever, AB , CD equimultiples to a like number of magnitudes E , F each to other; how multiple one magnitude AB is one E , so multiples are all the magnitudes $AB + CD$ to all the other magnitudes $E + F$.

Let AG , GH , HB the parts of the quantity AB , be equal to E , and also let CI , IK , KD the parts of the quantity CD be equal to F . The number of these are put equal to those. Now whereas $AG + CI = E + F$; a and $GH + IK = E + F$; a and $HB + KD = E + F$; it is evident that $AB + CD$ does so often contain $E + F$ as one AB contains E . Which was to be done. a 2. ax.

PROP. II.


If the first AB be equimultiple to the second C , as the third DE is to the fourth F , and if the fifth BG be equimultiple to the second C as the sixth EH is to the fourth F ; then shall the first compounded with the fifth (AG) be equimultiple to the second C , as the third compounded with the sixth (DH) is to the fourth F .

The number of parts in AB equal each to C is put equal to the number of parts in DE . whereof each part is equal to F . Likewise the number of parts in BG is equal to the number of parts in EH . Therefore the number of parts in $AB + BG$ is equal to the number of parts in $DE + EH$.



a 2. ax. FH. a That is, the whole line AG is as equimultiple of C, as the whole line DH is of F. Which was to be dem.

PROP. III.



If the first A be equimultiple of the second B, and the third C of the fourth D, and there be taken EI, FM equimultiples of the first and third, then will each of the magnitudes taken be alike equimultiple of both, the one EI to the second B, the other FM to the fourth D.

Let EG, GH, HI the parts of the multiple EI be equal to A, also let FK, KL, LM the parts of the multiple FM be equal to F, & the number of these is equal to the number of those. Moreover A (that is) EG or GH, or HI is put as

a hyp.

b 205.

c 2. 5,

equimultiple of B, as C, or FK, &c. of D. b Therefore EG + GH is equimultiple of the second B, as FK + KL is of the fourth D. c By the same way of argument is EI (EH + HI) as multiple of B, as FM (FL + LM) is of D. Which was to be done.

PROP.

PROP. IV.

If the first *A* have the same ratio to the second *B*, as the third *C* to the fourth *D*; then also *E* and *F* the equimultiples of the first *A* and the third *C*, shall have the same ratio to *G* and *H* the equimultiples of the second *B* and the fourth *D*, according to any multiplication, if so taken as they answer each to other (*E. G* :: *F. H.*)

Take *I* and *K* the equimultiples of *E* and *F*; and also *L* and *M* the equimultiples of *G* and *H*.
a Then is *I* as multiple of *A*, as *a* *K* of *C*; *a* and also *L* is as multiple of *B*, as *M* of *D*. Therefore whereas it is *A. B* :: *C. D*; *b* *hyp.*
 according to the sixth definition, if *I* be $\sqsubset, =, \sqsupset$ *L*, then consequently after the same manner is *K* $\sqsubset, =, \sqsupset$ *M*. Therefore when *I* and *K* are taken as multiples of *E* and *F*, as *L* and *M* of *G*, and *H*, then will it be by the seventh definition *E. G* :: *F. H.* Which was to be dem.

Coroll.


From hence is wont to be demonstrated the proof of the inverse ratio.

For because *A. B* :: *C. D*, therefore if *E* $\sqsubset, =, \sqsupset$ *G*, then is *c* likewise *F* $\sqsubset, =, \sqsupset$ *H*. *c* 2. def. 5.
 Therefore it is evident that if *G* $\sqsubset, =, \sqsupset$ *E*, then is *H* $\sqsubset, =, \sqsupset$ *F*; *d* therefore *B. A* :: *D. C.* *d* 6. def. 5.
 Which was to be dem.

G

PROP.

PROP. V.



If a magnitude AB be as multiple of magnitude CD , as part taken from the one AE of a part taken from the other CF ; the residue of the one shall be as multiple of the residue of the other FD as the whole AB is of the whole CD .


a 1. 5.

b 6. ax.

c 3. ax.

Take any other GA , which shall be as multiple to FD the residue, as AB is of the whole CD , or as the part taken away AE is of the part taken away CF . a Therefore the whole $GA - AE$ is as multiple of the whole $CF + FD$, as the one AE is of the one CF , that is as AB is of CD . therefore $GE = AB$; and c so GA that was common being taken away, there remains $GA = EB$.

PROP. VI.




If two magnitudes AB, CD be equimultiples of two magnitudes E, F , and some magnitudes AG and CH be taken away; then the residues GB, HD are either equal to those magnitudes E, F , or else equimultiples of them.

2 3. ax. 1.

For because the number of parts in AB , whereof each is equal to E , is put equal to the number of parts in CD , whereof each is equal to F , and also the number of parts AG equal to the number of parts in CH ; If from one you take AG , and from the other CH , a then remains the number of parts in the remainder GB equal to the number of parts in HD . therefore if GB be once E , then HD once F . if GB be many times F , then is HD so of F . Which was to be dem.

PROP.

PROP. VII.


 Equal magnitudes *A* and *B* have to the same magnitude *C* the same proportion or ratio. And one and the same magnitude *C* has the same ratio to equal magnitudes *A* and *B*.

Take *D* and *E* equimultiples of the equal magnitudes *A* and *B*, and *F* any-wise multiple of *C*; then is $D :: E$. Wherefore if $D :: E$, a 6. ax. $\neg F$, then also *E* will be $::, =, \neg F$. *b* there-*b* 6. def. 5. fore $A :: B$. *C* and *c* by inversion $C :: c$. *c* cor. 4. *d* *B*. Which was to be dem.

Schol.

If instead of the multiple *F*, two equimultiples be taken, it shall be the same way prov'd that equal magnitudes have the same ratio to other magnitudes that are equal between themselves.

PROP. VIII.

 Of unequal magnitudes *AB*, *AC*, the greater *AB* has a greater ratio to the same third line *D*, than the lesser *AC*; and the same third line *D* hath a greater ratio to the lesser *AC*, than to the greater *AB*.

Take *EF*, *EG* equimultiples of the said *AB*, *AC*, so that *EH* being multiple of *D* be greater than *EG*, but lesser than *EF*. (which will easily happen, if both *EG* and *GF* be taken greater than *D*.) It is manifest from 8. def. 5. that $\frac{AB}{D} < \frac{AC}{D}$ and $\frac{D}{AB} > \frac{D}{AC}$. Which was to be dem.

G 2

PROP.

PROP. IX.

Magnitudes which to one and the same magnitude have the same ratio, are equal the one to the other. And if a magnitude have the same ratio to other magnitudes, those magnitudes are equal one to the other.

a 8. 5.

A B C 1. Hyp. If $A.C :: B.C$; I say that $A=B$. For let A be greater or less than C, a then is $\frac{A}{C} < \text{or } > \frac{B}{C}$. Which is contrary to the Hypothesis.

b 8. 5.

2. Hyp. If $C.B :: C.A$. I say that $A=B$. For let A be $< B$, b then $\frac{C}{B} < \frac{C}{A}$. Which is against the Hypothesis.

PROP. X.

Of magnitudes having ratio to the same magnitude, that which has the greater ratio, is the greater magnitude; and that magnitude to which the same carries a greater ratio is the lesser magnitude.

a 7. 5.

b 8. 5.

A B C 1. Hyp. If $\frac{A}{C} < \frac{B}{C}$. I say that $A < B$. For if it be said that $A=B$, a then $A.C :: B.C$. Which is contrary to the Hyp. If $A > B$, b then is $\frac{A}{C} > \frac{B}{C}$. Which is also against the Hyp.

c 7. 5.

d 8. 1.

2. Hyp. If $\frac{C}{B} < \frac{C}{A}$ I say that $B > A$. for if you say $B=A$, it's against the Hypothesis, for it will follow that $C.B :: C.A$. If you say $B < A$, d then is $\frac{C}{A} < \frac{C}{B}$. Which is also against the Hyp.

PROP.

P R O P. XI.

G	_____	H	_____	I	_____
A	_____	C	_____	E	_____
B	_____	D	_____	F	_____
K	_____	L	_____	M	_____

Proportions which are one and the same to any third, are also the same one to another.

Let $A.B :: E.F$, and $C.D :: E.F$. I say that $A.B :: C.D$. Take G, H, I the equimultiples of A, C, E ; and K, L, M the equimultiples of B, D, F . Now *a* because $A.B :: E.F$, if $G \sqsubset, =, \supset$, *a* *hyp.*
 $\supset K$, *b* then after the same manner $I \sqsubset, =, \supset$, *b* *6. def. 3.*
 $\supset M$. And likewise *a* because $E.F :: C.D$. if $I \sqsubset, =, \supset$, $\supset M$, *b* then is H likewise $\sqsubset, =, \supset L$.
c wherefore $A.B :: C.D$. Which was to be dem. *c* *6. def. 5.*

Schol.

Proportions that are one and the same to the same proportions, are the same betwixt themselves.

P R O P. XII.

G	_____	H	_____	I	_____
A	_____	C	_____	E	_____
B	_____	D	_____	F	_____
K	_____	L	_____	M	_____

If any number of magnitudes $A, B; C, D; E$ and F be proportionals; as one of the Antecedents A is to one of the Consequents B ; so are all the Antecedents A, C, E to all the Consequents B, D, F .

Take the equimultiples of the antecedents G, H, I , and of the consequents K, L, M . Because that as multiple as one G is of one A , *a* so multi-*a* *1. 5.*
 ples are all G, H, I , of all A, C, E ; and likewise as multiple as one K is of one B , so multiples are all K, L, M , of all B, D, F . moreover because $A.B :: C.D :: E.F$. if G be $\sqsubset, =, \supset$, or $\supset K$. then *b* *hyp.*
 will H likewise be $\sqsubset, =, \supset L$, and $I \sqsubset, =, \supset M$. and so if $G \sqsubset, =, \supset K$. in like manner will $G + H + I$ be $\sqsubset, =, \supset K + L + M$. *c* wherefore $A.B :: C.D :: E.F$. *c* *6. def. 5.*
 $A + C + E.B + D + F$. Which was to be dem.

Coroll.

From hence, if like proportionals be added to like proportionals, the wholes shall be proportional.

PROP. XIII.

G	_____	H	_____	I	_____
A	_____	C	_____	E	_____
B	_____	D	_____	F	_____
K	_____	L	_____	M	_____

If the first *A* have the same ratio to the second *B*, that the third *C* has to the fourth *D*; and if the third *C* have a greater proportion to the fourth *D*, than the fifth *E* to the sixth *F*; then also shall the first *A* have a greater proportion to the second *B*, than the fifth *E* to the sixth *F*.

Take *G*, *H*, *I* equimultiples of *A*, *C*, *E*, and *K*, *L*, *M*, equimultiples of *B*, *D*, *F*. Now because a 6. def. 1, that $A : B :: C : D$. if $H \sqsubset L$. a then is $G \sqsubset K$.

b 8. def. 5. but because $\frac{C}{D} \sqsubset \frac{E}{F}$, b it may be that $H \sqsubset$

c 8. def. 5. L , and yet I not $\sqsubset M$. c Therefore $\frac{A}{B} \sqsubset \frac{E}{F}$.

Which was to be dem.

Schol.

But if $\frac{C}{D} \supset \frac{E}{F}$, then also is $\frac{A}{B} \supset \frac{E}{F}$. Also, if $\frac{A}{B} \sqsubset \frac{C}{D} \sqsubset \frac{E}{F}$, then is $\frac{A}{B} \sqsubset \frac{E}{F}$. And if $\frac{A}{B} \supset \frac{C}{D} \supset \frac{E}{F}$, then is $\frac{A}{B} \supset \frac{E}{F}$.

PROP.

PROP. XIV.

If the first *A* have the same ratio to the second *B*, that the third *C* has to the fourth *D*; and if the first *A* be greater than the third *C*; then shall the second *B* be greater than the fourth *D*. But if the first *A* be equal to the third *C*, then the second *B* shall be equal to the fourth *D*. but if *A* be lesser, the is *B* also lesser.

Let $A \sqsubset C$ a then $\frac{A}{B} \sqsubset \frac{C}{B}$. *b* but a 8. 5.
 $\frac{A}{B} = \frac{C}{D}$. c therefore $\frac{C}{D} \sqsubset \frac{C}{B}$. c theref. c 13. 5.
b hyp.

$B \sqsubset D$. By the like way of argument, if $A \supset C$,
 d then is $B \supset D$. But if A be put equal to C , d 10. 5.
 then $C : B :: e A : B$ f :: $C : D$. g therefore $B = D$. e 7. 5.
 Which was to be dem. f hyp.

Schol.

g 11. 5. &

By an argument *a fortiori*, if $\frac{A}{B} \supset \frac{C}{D}$, and A 9. 5.
 $\sqsubset C$. then is $B \sqsubset D$. Likewise if $A = B$, then
 is $B = D$. and if $A \sqsubset$, or $\supset B$, then also is C
 $\sqsubset$ or $\supset D$.

PROP. XV.

Parts *C* and *F* are in the same ratio, with
 their like multiples *AB* and *DE*, if taken
 correspondently. ($AB : DE :: C : F$.)

Let *AG*, *GB* parts of the multiple
AB be equal to *C*; and let *DH*, *HE*
 parts of the multiple *DE* be equal to *F*.

a The number of these parts is equal to a hyp.
 the number of those. Therefore whereas b 7. 5.

b $AG : C :: DH : F$, and $GB : C :: HE : F$. c 12. 5.
 therefore is c $AG : GB (AB) : DH : HE$
 (DE) : $C : F$. Which was to be dem.

G 4

PROP.

PROP. XVI.




If four magnitudes A, B, C, D be proportional, they also shall be alternately proportional ($A : C :: B : D$.)

Take E and F equimultiples of A and B ; take also G and H equimultiples of C and D . There-
 a 15. 5. fore $E : F :: A : B$ b 11. 5. $G : H :: C : D$. Where-
 b hyp. fore if $E < G$, $=$, $> G$, then likewise is $F <$,
 c 11. 5. $=$, $> H$. d Therefore $A : C :: B : D$. Which was
 14. 5. to be dem.
 d 6. def. 5. Schol.

Alternate ratio has place only then when the quantities are of the same kind. For heterogeneous quantities are not compared together.

PROP. XVII.



a 1. 5.
b constr.

c 1. 5.

d 2. 5.

e 6. def.

If magnitudes compounded be proportional $AB : CB :: DE : FE$, they shall be proportional also when divided ($AC : CB :: DF : FE$.)

Take GH, HL, IK, KM , in order the equimultiples of AC, CB, DF, FE ; and also LN, MO , the equimultiples of CB, FE . The whole GL is a as multiple of the whole AB , as one GH of one AC , b that is as IK of DF , c or as the whole IM of the whole DE . Also HN ($HL + LN$) is as d multiple of CB , as KO ($KM + MO$) is of FE . Therefore, whereas by Hyp. $AB : BC :: DE : EF$, if GL be $<$, $=$, $>$ HN , then likewise e will IM be $<$, $=$, $>$ KO . Take from these HL, KM that are equal;

equal; and if the remainder GH be $\square$, $=$, $\sqsupset$
LN, f then will IK $\square$, $=$, $\sqsupset$ MO, g whence f 5. ax.
AC. CB :: DF. EF. Which was to be dem. g 6. def. 5.

PROP. XVIII.

If magnitudes divided be proportional
(AB. BC :: DE. EF) the same also
being compounded shall be proportional
(AC. CB :: DF. FE).

For if it can be, let AB. CB :: DF.
FG $\square$ FE. a Then by division will a 17. 5.
AB. BC :: DG. GF. b that is, DG, b hyp. &
GF :: DE. EF. and being DG $\square$ FE. c
DE, c therefore is GF $\square$ EF. d Which c 14. 5.
is absurd. The like absurdity will follow if it d 9. ax.
be said AB. CB :: DE. GF $\square$ FE.

PROP. XIX.

If the whole AB be
to the whole DE as the
part taken away AC is
to the part taken away
DF, then shall the resi-
due CB be to the residue FE as the whole AB is to
the whole DE.

Because a AB. DE :: AC. DF, b therefore by a hyp.
permutation AB. AC :: DE. DF. c and thence b 10. 5.
by division AC. CB :: DF. FE. b wherefore c 17. 5.
again by permutation AC. DF :: CB. FE. d that d hyp. &
is, AB. DE :: CB. FE. Which was to be dem. 11. 5.

Coroll.


Hence, If like proportionals be subtracted
from like proportionals, the residues shall be
proportional.

2. Hence is converse ratio demonstrated.

Let AB. CB :: DE. FE. I say that AB. AC ::
DE. DF. For by a permutation AB. DE :: CB. a 16. 5.
FE. b therof. AB. DE :: AC. DF. whence again b 17. 5.
by permutation AB. AC :: DE. DF. Which was
to be dem.

PROP.

The fifth Book of



A B C D E F the first A be equal to the third C, then the fourth D is so to the sixth F; and if A be less than C, so D is less than F.

a hyp.

b cor. 4. 5.

c hyp. and

8. 5.

d schol. 13.

5.

e 10. 5.

f 7. 5.

g 11. 5. and

9. 5.

1. Hyp. Let $A = C$. Because $a E.F :: B.C$, by b in-
version shall be $F.E :: C.B$. e But $\frac{C}{B} \supset \frac{A}{B}$, therefore

$\frac{F}{E} \supset \frac{A}{B}$ or $\frac{D}{E}$, e therefore $D \sqsubset F$. W. W. to be dem.

2. Hyp. By the same way of argument, if $A \sqsupset C$, it will appear that $D \supset F$.

3. Hyp. If $A = C$. Because $F.E :: C.B :: f A.B :: D.E$. g therefore is $D = F$. Which was to be dem.

PROP. XXI.



A B C D E F the first A be greater than the third C; then is the fourth D greater than the sixth F: but if the first be equal to the third, then is the fourth equal to the sixth; if less, so is the other likewise.

a hyp.

1. Hyp. If $A \sqsubset C$; then because $a D.E :: B.C$.

b 8. 5.

therefore inversely $E.D :: C.B$. but $\frac{C}{B} \supset \frac{A}{B}$ c there-

c schol. 13.

fore $\frac{E}{D} \supset \frac{A}{B}$, that is $\frac{E}{F}$, d therefore $D \sqsubset F$.

5.

d 10. 5.

2. Hyp.

2. *Hyp.* By the like arguments, if $A \propto C$, then is $D \propto F$.

3. *Hyp.* If $A = C$; then because $E. D :: e C. e$ 7. 5. $B :: e A. B :: f E. F$, g therefore is $D = F$. *f hyp.*
Which was to be dem. g 9. 5.

PROP. XXII.



If there be any number of magnitudes A, B, C , and others equal to them in number D, E, F , which taken two and two are in the same ratio ($A. B :: D. E$. and $B. C :: E. F$) they shall be in the same ratio also by equality, ($A. C :: D. F$).


Take $G. H$ equimultiples of A, D ; and I, K of B, E ; and also L, M of C, F .

Because $a A. B :: a hyp. D. E$, b therefore $G. b$ 4. 5. $I :: H. K$. and in like manner $I. L :: K. M$. therefore, if $G \propto H$, $=$,

$\propto, L$, c then is $H \propto, =, \propto M$, d therefore $A. c$ 20. 5. $C :: D. F$. By the same way of demonstration if d 6. *def.* $C :: D. F$. further $C. N :: F. O$, then by equality $A. N :: D. O$. Which was to be dem.

PROP.

The fifth Book of
P R O P. XXIII.



If there be three magnitudes *A, B, C*, and others *D, E, F*, equal to them in number, which taken two and two are in the same ratio, and their proportionality inordinate (*A. B :: E. F.* and *B. C :: D. E.*) they shall be in the same ratio also by equality.

Take *G, H, I* equimultiples of *A, B, D*; and also *K, L, M* equimultiples of *C, E, F*. Then *G. H :: a A. B :: b E. F* *a :: L. M.* Moreover because *b B. C :: D. E*, thence is *c H. K :: I. L*; therefore *G, H, K*, and *I, L, M* are according to 21.5. Wherefore if *G* be $\sqsubset$,

a 15. 5.
b hyp.
c 4. 5.

$\sqsupset$, $\sqsupset$ *K*, then is likewise *I* $\sqsupset$, $\sqsupset$ *M*. and so *d* consequently *A. C :: D. F.* Which was to be
d 6. def. 5. demonstrated.

If there be more magnitudes than three, this way of demonstration holds good in them also.

Coroll.

*22. & 23. From hence * it follows that ratio's compounded of the same ratio's are among themselves the same; as also the same parts of the same ratio's, are among themselves the same.

P R O P. XXIV.


A ——— B ——— G If the first magnitude *AB*
C ——— have the same ratio to the second
D ——— E ——— H cond C, which the third *DE*
F ——— has to the fourth *F*; and if
the fifth *BG* have the same
ratio to the second *C*, which the sixth *EH* has to the
fourth *F*, then shall the first compounded with the
fifth (*AG*) have the same ratio to the second *C*,
which the third compounded with the sixth (*DH*) has
to the fourth *F*.

For

For because a $AB.C :: DE.F$, and by the Hyp. a hyp.
and inversion $C.BG :: F.EH$; therefore by b e- b 22. 5.
equality $AB.BG :: DE.EH$. whence by com-
pounding $AG.BG :: DH.EH$. Also c $BG.C :: c$ hyp.
 $EH.F$. Therefore again by b equality $AG.C ::$
 $DH.F$. Which was to be dem.

P R O P. XXV.


If four magnitudes be proportional ($AB.$
 $CD :: E.F$) the greatest AB and the least
 F shall be greater than the rest CD , and E .



Make $AG = E$, and $CH = F$. Because
 $AB.CD :: a$ $E.F :: b$ $AG.CH$. c thence a hyp.
is $AB.CD :: GB.HD$. d but $AB < CD$. b 7. 5.
 e therefore $GB < HD$. But $AG + F = c$ 19. 5.
 $E + CH$, therefore $AG + F + GB < d$ hyp.
 $E + CH + HD$, that is, $AB + F < e$ schol. 14.
 $+ CD$. Which was to be dem. 5.

These Propositions which follow are not Euclide's,
but taken out of other Authors, and here subjoined
because of their frequent use.

P R O P. XXVI.




If the first have a
greater proportion to
the second, than the
third to the fourth,
then contrarywise, by

conversion, the second shall have a less proportion to
the first, than the fourth to the third.

Let $\frac{A}{B} < \frac{C}{D}$. I say that $\frac{B}{A} > \frac{D}{C}$. For conceive

$\frac{C}{D} = \frac{E}{B}$; a therefore $\frac{A}{B} < \frac{E}{B}$ b whence $A < E$. c there- a 13. 5.
fore $\frac{B}{A} > \frac{B}{E}$, d or $\frac{D}{C}$ Which was to be dem. b 10. 5.
 c 8. 5.
 d cor. 4. 5.

P R O P. XXVII.




If the first have a greater pro-
portion to the second, than the
third to the fourth; then alter-
nately the first shall have a greater
proportion to the third, than the second to the fourth.

Let

Let $\frac{A}{B} \sqsubset \frac{C}{D}$. then I say $\frac{A}{C} \sqsubset \frac{B}{D}$. For conceiv

a 10. 5. $\frac{E}{B} = \frac{C}{D}$. a therefore $A \sqsubset E$, b and therefore $\frac{A}{C} \sqsubset \frac{E}{D}$
 b 1. 5. $\frac{B}{D}$
 c 16. 5. c or $\frac{B}{D}$. Which was to be dem.


PROP. XXVIII.



If the first have a greater proportion to the second than the third to the fourth, then the first compounded with the second shall have a greater proportion to the second, than the third compounded with the fourth to the fourth.

Let $\frac{AB}{BC} \sqsubset \frac{DE}{EF}$. I say that $\frac{AC}{BC} \sqsubset \frac{DE}{EF}$. For conceive $\frac{GB}{BC} = \frac{DE}{EF}$. a therefore is $AB \sqsubset GB$. add BC to each part. then b will $AC \sqsubset GC$. c therefore $\frac{AC}{BC} \sqsubset \frac{GC}{BC}$. d that is $\frac{DF}{EF}$. Which was to be dem.

PROP. XXIX.



If the first compounded with the second have a greater proportion to the second, than the third compounded with the fourth has to the fourth; then by division the first shall have a greater proportion to the second, than the third to the fourth.

Let $\frac{AC}{BC} \sqsubset \frac{DF}{EF}$. then I say $\frac{AB}{BC} \sqsubset \frac{DE}{EF}$. For conceive $\frac{GC}{BC} = \frac{DE}{EF}$. a theref. $AC \sqsubset GC$. Take away BC, that is common; then b remains $AB \sqsubset GB$. c Therefore $\frac{AB}{BC} \sqsubset \frac{GB}{BC}$ d or $\frac{DE}{EF}$. Which was to be dem.

PROP.

PROP. XXX.

If the first compounded with the second have a greater proportion to the second, than the third compounded with the fourth has to the fourth, then by converse ratio shall the first compounded with the second have a lesser ratio to the first, than the third compounded with the fourth shall have to the third.

Let $\frac{AC}{BC} < \frac{DF}{EF}$. Then I say that $\frac{AC}{AB} > \frac{DF}{DE}$.

For because that $\frac{AC}{BC} < \frac{DF}{EF}$, *a* therefore by division $\frac{AB}{BC} < \frac{DE}{EF}$ by conversion *c* therefore $\frac{BC}{AB} > \frac{EF}{DE}$ and *d* therefore by composition $\frac{AC}{AB} > \frac{DF}{DE}$. Which was to be dem.

a hyp. b 29. 5. c 26. 5. d 28. 5.

PROP. XXXI.

If there be three magnitudes A, B, C, and others also D, E, F equal to them in number; and if there be a greater proportion of the first of the former to the second, than there is of the first of the last to their second ($\frac{A}{B} < \frac{D}{E}$;) and there be also a greater proportion of the second of the first magnitudes to the third, than there is of the second of the last magnitude to their third ($\frac{B}{C} < \frac{E}{F}$.) Then by equality also shall the ratio of the first of the former magnitudes to the third be greater than the ratio of the first of the latter magnitudes to the third ($\frac{A}{C} < \frac{D}{F}$.)

Con-

- a 10. 5.
b 8. 5.
c 13. 5.
d 10. 5.
e 8. 5.
f 12. 5.

Conceive $\frac{G}{C} = \frac{E}{F}$. *a* therefore is $E \sqsubset G$, and *b*
therefore $\frac{A}{G} \sqsubset \frac{A}{B}$. Again conceive $\frac{H}{G} = \frac{D}{F}$.
c therefore $\frac{H}{G} \sqsupset \frac{A}{B}$, therefore much more $\frac{H}{G} \sqsupset \frac{A}{C}$.
d wherefore $A \sqsubset H$, *e* and consequently $\frac{A}{C} \sqsubset \frac{H}{C}$
f or $\frac{D}{F}$.

P R Q P. XXXII.

A ——— D ——— If there be three mag-
 B ——— E ——— nitudes A, B, C , and o-
 C ——— F ——— thers D, E, F , equal to
 G ——— them in number; and
 H ——— there be a greater pro-
portion of the first of the
former magnitudes to the second, than there is of the
second of the latter to the third $\left(\frac{A}{B} \sqsubset \frac{E}{F}\right)$ and also
the ratio of the second of the former to the third be
greater than the ratio of the first of the latter to the
second $\left(\frac{B}{C} \sqsubset \frac{D}{E}\right)$. then by equality also shall the
proportion of the first of the former to the third, be
greater than that of the first of the latter to the third
 $\left(\frac{A}{C} \sqsubset \frac{D}{E}\right)$.

The demonstration of this Proposition is alto-
gether like that of the precedent.

P R O P. XXXIII.

A — E — B If the proportion of the whole AB
 C — I — D to the whole CD be greater than
 F the proportion of the part taken a-
way AE to the part taken away
 CF ; then shall also the ratio of the
remaindes EB to the remainder FD be greater than
that of the whole AB to the whole CD .

Be-

Because that $\frac{AB}{CD} = \frac{AE}{EF}$, *b* therefore by permutation $\frac{AB}{AE} = \frac{CD}{CF}$, *c* therefore by converse ratio $\frac{AB}{EB} = \frac{CD}{FD}$, and by permutation again $\frac{AB}{CD} = \frac{EB}{FD}$.

PROPOSITION XXXIV.

A ———	D. ———	If there be any number of magni- tudes, and others al- so equal to them in number; and the
B ———	E ———	
C ———	F ———	
G ———	H ———	

proportion of the first of the former to the first of the latter, be greater than that of the second to the second, and that greater than the proportion of the third to the third, and so forward: all the former magnitudes together shall have a greater ratio to all the latter together, than all the former, leaving out the first, shall have to all the latter, leaving out the first; but less than that of the first of the former to the first of the latter; and lastly greater than that of the last of the former to the last of the latter.

You may please to consult Interpreters for the demonstration hereof, we having for brevity sake omitted it, and because 'tis of no use in these Elements.


The End of the fifth Book.

THE

THE


THE SIXTH BOOK OF EUCLIDE'S ELEMENTS

Definitions.



I. Like right-lined figures (ABC, DCE) are such whose several angles are equal one to the other, and also their sides about the equal angles, proportional.


The angle $B = DCE$, and $AB. BC :: DC. CE$. Also the angle $A = D$, and $BA. AC :: CD. DE$. Lastly the angle $ACB = E$, and $BC. CA :: CE. ED$.



II. Reciprocal figures are (BD, BF) when in either figure are the terms antecedents and consequents of ratio's, that is, $AB. BG :: EB. BC$.

III. A right line AB is said to be cut according to mean and extreme proportion, when as the whole AB is to the greater segment AC , so is the greater segment AC to the less CB ($AB. AC :: AC. CB$).

IV. The




IV. The altitude of any figure ABC, is a perpendicular line AD drawn from the top A to the base BC.

V. A ratio is said to be compounded of two ratio's, when the quantities of the ratio's being multiplied the one into the other, do produce any ratio. As the ratio of A to C is compounded of the ratio's

of A to B and B to C. For $\frac{A}{B} \times \frac{B}{C} = \frac{A}{C}$ a 10. def. 3. b 15. 4.


PROP. I.



Triangles ABC, ACD, and parallelograms BC-AE, CDFA which have the same height, are in proportion one to the other, as their bases BC, CD are.

a Take as many as you please, BG, GH, equal to BC, and also DI = CD. and join AG, AH, a 3. 1. AI.

b The triangles ACB, ABG, AGH are equal, b 38. 1. and b also the triangle ACD = ADI. Therefore the triangle ACH is as multiple of the triangle ACB, as the base HC is of the base BC; and the triangle ACI as multiple of the triangle ACD, as the base CI is of CD. But if $HC = n \times BC$, $CI = n \times CD$, then is likewise the triangle AHC = $n \times$ triangle ACI; and d therefore d 6. def. 5. BC. CD :: the triangle ABC. ACD :: e Pgr. CE. e 41. 1. 6. CF. Which was to be dem. 15. 5.



Hence, Triangles ABC , DEF , and Pgrs. $AGBC$, $DEFH$, whose bases BC , EF are equal, are in such proportion as their altitudes, AI , DK are.

- a 3. 1. a Take $IL = CB$, and $KM = EF$; and join
 b 7. 5. LA , LG , MD , MH . then is it evident that the
 c 1. 6. triangles ABC , $DEF :: b$ AIL , $DKM :: c$ AIL , DK .
 d 41. 1. & d pgrs. $AGBC$, $DEFH$. Which was to be dem.
 15. 5.

P R O P. II.

If to one side BC of a triangle ABC be drawn a parallel right line DE , the same shall cut the sides of the triangle proportionally (AD , $BD :: AE$, EC). And if the sides of the triangle be proportionally cut (AD , $BD :: AE$, EC) then a right line DE joined at the sections D , E shall be parallel to the remaining side of the triangle BC . Draw CD and BE .


- a 17. 1. 1. Hyp. Because the triangle DEB a = DEC ,
 b 7. 5. b therefore shall be the triangle ADE . $DBE ::$
 c 2. 6. ADE , ECD . But the triangle ADE . $DBE :: c$
 d 11. 5. AD , DB , and the triangle ADE . $DEC :: AE$,
 e 1. 6. EC , d therefore AD . $DB :: AE$. EC .
 f 19. 5. 2. Hyp. Because AD . $DB :: AE$. EC , e that is
 g 39. 1. the triangle ADE $DBE :: ADE$. ECD ; f there-
 fore is the triangle $DBE = ECD$; and g there-
 fore DE , BC are parallels. Which was to be dem.

Scholium.

If there be drawn many parallels to one side of any triangle, then all the segments of the sides shall

shall be proportional; as is easily deducible from the precedent.

PROP. III.




If an angle BAC of a triangle BAC be bisected, and the right line AD , that bisects the angle, cut the base also; then shall the segments of the base have the same ratio that the other sides of the triangle have ($BD. DC :: AB. AC$.) And if the segments of the base have the same ratio, that the other sides of the triangle have ($BD. DC :: AB. AC$.) then a right line AD drawn from the top A to the section D , shall bisect that angle BAC of the triangle.

Produce BA , and make $AE = AC$, and join CE .

1. Hyp. Because $AE = AC$, therefore is the angle $ACE = E$ $b = \frac{1}{2} BAC$ $c = DAC$; d a 5. 1. therefore DA, CE are parallels. e Wherefore b 32. 1. $BA. AE (AC) :: BD. DC$. c hyp.

2. Hyp. Because $BA. AC (AE) :: BD. DC$, f d 27. 1. therefore are DA, CE parallels; and g therefore e 2. 6. is the angle $BAD = E$; and the angle DAC g f 2. 6. $= ACE$ $h = E$. k therefore the angle $BAD = g$ 29. 1. DAC . Wherefore the angle BAC is bisected. n 5. 1. Which was to be dem. k 1. ax,

PROP. IV.



Of equiangular triangles ABC, DCE , the sides are proportional which are about the equal angles, B, DCE , ($AB. BC :: DC. CE$, &c.) And the sides AB, DC , &c. which are subtended under the equal angles ACB, E , &c. are homologous, or of like ratio.

Set the side BC in a direct line to the side CE , and produce BA and ED till they meet. a 32. 1.

H 3

Be-

- b hyp. Because the angle B $b = ECD$, c therefore
 c 28. 1. BF, CD are parallel; Also because the angle
 BCA $b = CED$, e therefore are CA, EF parallel.
 d 34. 1. Therefore the figure CAFD is a Pgr. d therefore
 AF = CD, and AC = d ED. Whence it is evi-
 e 2. 6. dent that AB. AF (CD) :: e BC. CE. f by permu-
 f 16. 5. tation therefore AB. BC :: CD. CE. also BC.
 CE :: FD (AC.) DE. f and thence by per-
 g 22. 5. mutation BC. AC :: CE. DE. g Wherefore also
 by equality AB. AC :: CD. DE. Therefore, &c.


Coroll.

Hence AB. DC :: BC. CE :: AC. DE.

Scol.

Hence, If in a triangle FBE there be drawn AC a parallel to one side FE, the triangle ABC shall be like to the whole FBE.

PROP. V.




If two triangles ABC, DEF have their sides proportional (AB. BC :: DE. EF, and AC. BC :: DF. EF, and also AB. AC :: DE. DF) those triangles are equiangular, and those angles e-

qual under which are subtended the homologous sides.

- a 23. 1. At the side EF a make the angle FEG = B,
 b 32. 1. and the angle EFG = C; b whence the angle
 c 4. 6. G = A. Therefore GE. EF c :: AB. BC :: DE.
 d hyp. EF. e and therefore GE = DE. Likewise GF.
 e 11. 5. and EF c :: AC. CB :: d DF. FE. e therefore GF =
 9. 5. DF. Therefore the triangles DEF, GEF are
 f 8. 1. mutually equilateral. f Therefore the angle D =
 g 32. 1. G = A, and the angle FED f = FEG = B, and
 g consequently the angle DEF = C. There-
 fore, &c.

PROP.

PROP. VI.




If two triangles ABC , DEF have one angle B equal to one angle DEF , and the sides about the equal angles B, DEF proportional ($AB.BC :: DE.EF$) then those triangles ABC, DEF are equian-

gular, and have those angles equal, under which are subtended the homologous sides.

At the side EF make the angle $FEG = B$, and the angle $EFG = C$; then will the angle $G = A$. a 32. 1. Therefore $GE.EF :: b AB.BC :: c DE.EF$, d and b 4. 6. therefore $DE = GE$. But the angle $DEF = b f = c$ hyp. GEF ; therefore the angle $D = g = G = A$, h and d 9. 5. consequently the angle $EFD = C$. *W.W. to be dem.* e hyp.

PROP. VII.




If two triangles ABC, DEF have one angle A equal to one angle D , and the sides about the other angles ABC, E , proportional ($AB.BC :: DE.EF$) and if they have both of the re-

maining angles C, F either less or not less than a right angle; then shall the triangles ABC, DEF be equiangular, and have those angles equal about which the proportional sides are.

For, if it can be, let the angle $ABC = E$, and make the angle $ABG = E$. Therefore, whereas the angle $A = D$, b thence is the angle AGB a hyp. $= F$. Therefore $AB.BG :: c DE.EF :: d AB.b$ 32. 1. $BC.$ e therefore $BG = BC$. f therefore the angle c 4. 6. $BGC = BCG$. g Therefore BGC or C is less d hyp. than a right angle, and b consequently AGB or e 9. 5. F is greater than a right: Therefore the angles f 5. 1. C and F are not of the same species or kind, g cor. 17. 1, which is against the Hypothesis. h cor. 12. 1.

PROP. VIII.



If in a right-lined triangle ABC , from the right angle BAC there be drawn AD a perpendicular to the base BC ; then the triangles about the perpendicular (ADB , ADC) are like both to the whole triangle ABC , and also one to the other.

a hyp.

b 12. ax.

c 32. & 4.6

d 21. 6.

e 1. def.8.


For because BAC , ADB are a right angles, b and so equal, and B common; the triangles BAC , ADB c are like. By the same argument BAC , ADC are like d whence also ADB , ADC will be like. Which was to be dem.

Coroll.

Hence, 1. $BD. DA :: DA. DC$.

2. $BC. AC :: AC. DC$. and $CB. BA :: BA. BD$.

PROP. IX.



a 31. 1.

b 31. 1.

c 2. 6.

d 28. 3.


From a right line given AB to cut off any part required, as $\frac{1}{3}$ (AG .)

From the point A draw an infinite line AC any-wise, in which, a take any three equal parts AD, DE, EF . join FB . to which from D b draw the parallel DG . And the thing is done.

For $GB. AG :: c FD. AD$; whence by d composition $AB. AG :: AF. AD$. therefore, whereas $AD = \frac{1}{3}$ of AF , therefore is $AG = \frac{1}{3}$ of AB . Which was to be done.

PROP.

PROP. X.




To divide a right line given
AB not divided (in *F* and *G*)
 as another line given *AC* was
 cut (in *D* and *E*.)

Let a right line *BC* join
 the extremities of the line
 divided, and of the line not
 divided; and to that line from the points *E*, *D*
 draw the parallels *EG*, *DF* meeting with the ^a 31. 1.
 right line that is to be cut, in *G* and *F*; then
 the thing is done.

For let *DH* be a drawn parallel to *AB*. Then
AB. *DE* :: *b* *AF*. *FG*. and *DE*. *EC* *b* :: *DI*. ^b 2. 6.
IH :: *c* *FG*. *GB*. Which was to be done. ^c 34. 1. &
 7. 5.

Schol.



Hence is learnt to cut a right line given *AB* into
 as many equal parts as you please (suppose 5;)
 which will be more easily performed thus.


Draw an infinite line *AD*, and another *BH* pa-
 rallel to it, and infinite also. Of those take equal
 parts, *AR*, *RS*, *SV*, *VN*; and *BZ*, *ZX*, *XT*, *TL*;
 in

in each line less parts by one, than are required in AB; then let the right lines LR, TS, XV, ZN be drawn; these lines so drawn shall cut the right line given AB into five parts.

a 33. 1.
b constr.
c 2. 6.

For RL, ST, VX, NZ are *a* parallels; therefore, whereas AR, RS, SV, VN are *b* equal; *c* thence AM, MO, OP, PQ are equal also. Likewise, because that $BZ = ZX$, therefore is $BQ = PQ$. and therefore AB is cut into five parts. Which was to be dem.

PROP. XI.



Two right lines being given AB , AD , to find out a third in proportion to them (DE .)

Join BD , and from AB being produced take $BC = AD$. Thro' C draw CE parallel to DB ; with which let AD produced meet in E . then is DE the proportional required.

a 2. 6.

For $AB, BC (AD) a :: AD, DE$. Which was to be dem.




b cor. 9.

Or thus: Make the angle ABC right, and also the angle ACD right, then $b AB, BC :: BC, BD$.

PROP.


PROP. XII.



Three right lines being given DE, EF, DG to find out a fourth proportional GH .

Join EG , and thro' F draw FH parallel to EG ; with which let DG produced to H meet. Then it is evident that $DE. EF :: DG. GH$. a 2. 6. Which was to be done.

PROP. XIII.




Two right lines being given AE, EB , to find out a mean proportional EF .

Upon the whole line AB as a diameter describe a semicircle AFB , and from E erect a perpendicular EF meeting with the periphery in F . then $AE. EF :: EF. FB$. For let AF and FB be drawn; a then from the right angle of the a 31. 1. right-angled triangle AFB is drawn a right line FE perpendicular to the base. b Therefore $AE. b$ cor. 8. 6. $FE :: FE. EB$. Which was to be done.

Coroll.

Hence, A right line drawn in a circle from any point of the diameter perpendicularly, and extended to the circumference, is a mean proportional betwixt the two segments of the diameter.

PROP. XIV.




Equal Parallelograms having one angle ABC equal to one EBG , have the sides BD , BF which are about the equal angles reciprocal ($AB \cdot BG :: EB \cdot BC$;) And those Parallelograms BD , BF which have

one angle ABC equal to one EBG , and the sides which are about the equal angles reciprocal, are equal.

- For let the sides AB , BG about the equal angles make one right line; *a* wherefore EB , BC shall do the same. Let FG , DC be produced till they meet.
- a* sch. 15. 1. 1. Hyp. $AB \cdot BG :: b :: f$ $BD \cdot BH :: c$ $BF \cdot BH ::$
 - b* 1. 6. d $BE \cdot BC$. *e* therefore, &c.
 - c* 7. 5. 2. Hyp. $BD \cdot BH :: f$ $AB \cdot BG :: g$ $BE \cdot BC :: h$
 - d* 1. 6. $BF \cdot BH$. *k* Therefore the Pgr. $BD = BF$. Which
 - e* 11. 5. was to be dem.
 - f* 1. 6.
 - g* hyp.
 - h* 1. 6.
 - k* 11. and
 - 9. 5.

PROP. XV.




Equal triangles having one angle ABC equal to one DBE , their sides which are about the equal angles are reciprocal ($AB. BE :: DB. BC$.) And those triangles that have one angle ABC equal to one DBE , and have also the sides that are about the equal angles reciprocal ($AB. BE :: DB. BC$) are equal.

Let the sides CB, BC , which are about the equal angles be set in a strait line; a therefore a *sch. 15.1.* ABE is a right line. Let CE be drawn.

1. Hyp. $AB. BE :: b$ the triangle $ABC. CBE c :: b$ 1. 6.
the triangle $DBE. CBE :: d$ $DB. BC. e$ therefore, &c. c 7. 5.

1. Hyp. The triangle $ABC. CBE :: f$ $AB. BE :: g$ $DB. BC h ::$ the triangle $DBC. CBE. k$ e 11. 5.
Therefore the triangle $ABC = DBE$. Which f 1. 6.
was to be dem. g hyp. h 1. 6. k 11, and $9. 5.$

PROP. XVI.



If four right lines be proportional ($AB. FG :: EF. CB$) the rectangle AC comprehended under the extremes AD, CB , is equal to the rectangle EG comprehended under the means FG, EF . And if the rectangle AC comprehended under the extremes AB, CB be equal to the rectangle EG comprehended under the means FG, EF , then are the four right lines proportional, ($AB. FG :: EF. CB$)


1. Hyp.

- a 12. ax. 1. Hyp. The angles B and F are right, and consequently equal, and by Hyp. $AB.FG::EF.CB$,
 b 14. 6. therefore the rectangle $AC = EG$.
 c hyp. 2. Hyp. The rectangle $AC = EG$, and the
 d 14. 6. angle $B = F$; therefore $AB.FG::EF.CB$.
Which was to be dem.

Coroll.

- c 12. 6. Hence, it is easy to apply a rectangle given EG to a right line given AB; (viz.) e by making $AB.EF::FG.BC$.

PROP. XVII.



If three right lines be proportional ($AB.EF::EF.CB$) the rectangle AC made under the extremes AB, CB is equal to the square EG made of the middle EF . And if the rectangle AC comprehended under the extremes AB, CB be equal to the square EG made of the middle EF , then the three lines are proportional, ($AB.EF::EF.CB$).

Take $FG = EF$.


- a hyp. 1. Hyp. $AB.EF::a EF (FG.) CB$. therefore
 b 16. 6. the rectangle $AC b = EG c = EFq$.
 c 29. def. 1. 2. Hyp. The rectangle $AC d$ to the square
 d hyp. $EG = EFq$. e therefore $AB.EF::FG (EF.) BC$.
 e 16. 6. *Which was to be dem.*

Coroll.

Let $A \times B = Cq$. therefore $A.C::C.B$.

PROP.

PROP. XVIII.



From a right line given AB to describe a right-lined figure $AGHB$ like and alike situate to a right-lined figure given $CEFD$.

Resolve the right-lined figure given into triangles; a Make the angle $ABH = D$, a and a 23. 1. the angle $BAH = DCF$, a and the angle $AHG = CFE$, b and the angle $HAG = FCE$. then $AGHB$ shall be the right-lined figure sought.

For the angle B $b = D$, and the angle BAH b *constr.* $b = DCF$. c wherefore the angle $AHB = CFD$. c 32. 1. b also the angle $HAG = FCE$, and the angle AHG $b = CFE$. c wherefore the angle $G = E$, and the whole angle GAB $d = ECD$, and the d 2. *ax.* whole angle GHB $d = EFD$. The Polygons therefore are mutually equiangular. Moreover because the triangles are equiangular, therefore $AB.BH$ $e :: CD.DF$; and $AG.GH$ $e :: CE.EF$. e 4. 6. Likewise $AG.AH :: e$ $CE.CF$. and $AH.AB :: f$ 22. 5. $CF.CD$. f From whence by equality $AG.AB :: g$ 6. *def.* 5. $CF.CD$. After the same manner $GH.HB :: EF.FD$. Therefore the Polygons $ABHG$, $CDEF$ are like and alike situate. Which was to be done.

PROP. XIX.



Like triangles ABC , DEF are in duplicate ratio of their homologous sides, BC , EF .


a Let there be a 11. 6. made BC , $EF :: EF.BG$. and let AG be drawn. Because that $AB.DE$

b cor. 4. 6. $DE :: BC$. $EF :: EF$. BG , and the angle $B = E$.
 c constr. d therefore is the triangle $ABG = DEF$. But the
 d 15. 6. triangle ABC . $ABG :: e BC$. BG , and $f \frac{BC}{BG} = \frac{BC}{EF}$
 e 1. 6.
 f 10. def. 5.
 g 11. 5. twice; therefore $\frac{ABC}{ABG}$ that is $\frac{ABC}{DEF} g = \frac{BC}{EF}$ twice.
Which was to be dem.

Coroll.

Hence, If three right lines (BC , EF , BG) be proportional, then as the first is to the third, so is a triangle made upon the first BC , to a triangle like and alike described upon the second EF : or so is a triangle described upon the second EF to a triangle like and alike described upon the third.

PROP. XX.



Like Polygons $ABCDE$, $FGHIK$ are divided into equal triangles ABC , FGH , and ACD , FHI , and ADE , FIK ; both equal in number, and homologous to the wholes (ABC . $FGH :: ABCDE$. $FGHIK :: ACD$. $FHI :: ADE$. FIK .) And the Polygons $ABCDE$, $FGHIK$ have a double ratio one to the other what one homologous side BC has to the other homologous side GH .

a hyp.
 b 6. 6.

1. For the angle $B = G$, and AB . $BC :: a :: FG$. GH . b therefore the triangles ABC , FGH
 are

are equiangular. After the same manner are the triangles AED, FKI like. Whereas therefore the angle BCA $b =$ GHF, and the angle ADE $b =$ FIK, and the whole angles BCD, GHI, and the whole angles CDE, HIK are c equal, there remains the angle ACD $d =$ FHI, and the angle ADC $=$ FIH. e from whence also the angle CAD $=$ HFI. therefore the triangles ACD, FHI are like. Therefore, &c.

2. Because that the triangles BCA, GHF are like, therefore is $\frac{BCA}{GHF} = \frac{BC}{GH}$ twice. For the same reason is $\frac{CAD}{HFI} = \frac{CD}{HI}$ twice; lastly $\frac{DEA}{IKF} = \frac{DE}{IK}$ twice. now whereas that BC. GH $g ::$ CD. HI g *hyp. and* $g ::$ DE. IK. b therefore is the triangle BCA. GHF $::$ CAD. HFI $::$ DEA. IKF $k ::$ the polygon ABCDE. FGHK $:: \frac{BC}{GH}$ twice.

Coroll.


1. Hence, If there be three right lines proportional, then as the first is to the third, so is a polygone made upon the first to a polygone made on the second, like and alike described; or so is a polygone described upon the second to a polygone made on the third like and alike described.

By which is found out a method of enlarging or diminishing any right-lined figure in a ratio given : As if you would make a pentagone quintuple of that pentagone whereof CD is the side, then be-
twixt AB and 5 AB find out a mean proportional * upon this raise a pentagone like to that given, and it shall be quintuple of the pentagone given.

2. Hence also, If the homologous sides of like figures be known, then will the proportion of the figures be evident, viz. by finding out a third proportional.

I


PROP.



Right lined figures ABC , DIE which are like to the same right-lined figure HFG , are also like one to the other.

a 1. def. 6. For the angle $A = H = D$; and the angle $C = G = E$; and the angle $B = F = I$. Also $AB.AC :: HF.HG :: DI.DE$. & $a AC.CB :: HG.GF :: DE.EI$. And $AB.BC :: HF.FG :: DI.IE$. Therefore ABC , DIE are alike. Which was to be dem.

PROP. XXII.



If four right lines be proportional ($AB.CD :: EF.GH$) the right-lined figures also described upon them being like and in like sort situate, shall be proportional ($ABI.CDK :: EM.GO$.) And if the right-lined figures described upon the lines, like and alike situate, be proportional ($ABI.CDK :: EM.GO$.) then the right lines also shall be proportional ($AB.CD :: EF.GH$.)

a 19. 6. 1. Hyp. $\frac{ABI}{CDK} a = \frac{AB}{CD} \text{ twice } a = \frac{EF}{GH} \text{ twice } a = \frac{EM}{GO}$

b hyp. b therefore $ABI.CDK :: EM.GO$.

c 20. 6. 2. Hyp. $\frac{AB}{CD} \text{ twice } a = \frac{ABI}{CDK} b = \frac{EM}{GO} = c \frac{EF}{GH}$


d cor. 23. 5. twice. Therefore $AB.CD :: EF.GH$. Which was to be dem.

Schol.

Schol.

Hence is deduced the manner and reason of multiplying surd quantities. ex. gr. Let $\sqrt{5}$ to be multiplied into $\sqrt{3}$. I say that the product will be $\sqrt{15}$. For by the definition of multiplication it ought to be, as $1. \sqrt{3} :: \sqrt{5}.$ to the product. Therefore by this $q. 1. q. \sqrt{3} :: q. \sqrt{5}.$ $q.$ of the product. therefore the square of the product is 15. Wherefore $\sqrt{15}$ is the product of $\sqrt{3}$ into $\sqrt{5}$. Which was to be dem.

THEOREM.



If a right line AB be cut any-wise in D , the rect-*Pet. Herig.* angle comprehended under the parts AD, DB is a mean proportional betwixt their squares. Likewise the rectangle comprehended under the whole AB and one part AD , or DB , is a mean proportional betwixt the square of the whole AB and the square of the said part AD , or DB .

Upon the diameter AB describe a semicircle; from D erect a perpendicular DE meeting with the periphery in E . join AE, BE .

It's evident that $AD. DE :: DE. DB.$ *b* there-*a* cor. 8. 6. fore $ADq. DEq. :: DEq. DBq.$ *c* that is, $ADq. b$ 22. 6. $ADB :: ADB. DBq.$ Which was to be dem. *c* 17. 6.


Moreover $BA. AE :: d AE. AD.$ *e* therefore $BAq. d$ cor. 8. 6. $AEq. :: AEq. ADq.$ *f* that is, $BAq. BAD :: BAD. e$ 22. 6. $ADq.$ After the same manner $ABq. ABD :: ABD. f$ 17. 6. $BDq.$ Which was to be dem.

Or thus: suppose $Z = A + E$. It is manifest that $Aq. AE :: a A. E :: a AE. Eq.$ alio $Zq. ZA :: a a$ 1. 6. $Z. A :: ZA. Aq.$ and $Zq. ZE :: a Z. E :: ZE. Eq.$

I 2

PROP.

PROP. XXIII.



Equiangular parallelograms AC, CF, have the ratio one to the other; which is compounded of their sides.


$$\left(\frac{AC}{CF} = \frac{BC}{CG} + \frac{DC}{CF} \right)$$

Let the sides about the equal angles C be a set in a direct line, and let the Pgr. CH be completed. Then is the ratio of $\frac{AC}{CF} = \frac{AC}{CH} + \frac{CH}{CF} = \frac{BC}{CG} + \frac{DC}{CE}$. Which was to be dem.

Coroll.

Andr. Hence, and from 34. 1. it appears, 1. That triangles which have one angle equal (as C) have a ratio compounded of the ratios of the right lines (AC to CB, and LC to CF) containing the equal angle.

* 35. 1.



2. That all rectangles, and * consequently all parallelograms, have their ratio one to the other compounded of the ratios of base to base, and altitude to altitude. After the like manner you may argue in triangles.


3. From hence is apparent how to give the proportion of triangles and parallelograms. Let there be two Pgrs. X and Z, whose bases are AC, CB, and altitudes CL, CF. Make CL, CF :: CB, O. * then will it be X.Z :: AC, O.

* 14. 6.

and 1. 6.

PROP.


PROP. XXIV.



In every parallelogram $ABCD$, the parallelograms EG , HF which are about the diameter AC are like to the whole, and also one to the other.

For the Pgrs. EG , HF have each of them one angle common with the whole, a therefore they are equiangular to the whole, a 29. 1. and also one to the other. Also both the triangles ABC , AEI , IHC a and the triangles ADC , AGI , IFC are equiangular mutually; b therefore b 4. 6. $AE.EI :: AB.BC$, and b $AE.AI :: AB.AC$, and b $AI.AG :: AC.AD$. c Therefore by equality, c 22. 5. $AE.AG :: AB.AD$. d Therefore the Pgrs. EG , d 1. def. 8. BD are like, After the same manner are HF , 6 . BD like also. Therefore, &c.

PROP. XXV.




Unto a right-lined figure given $ABEDC$ to describe another figure P like and alike situate, which also shall be equal to another right-lined figure given F .

a Make the rectangle $AL = ABEDC$; b also a 45. 1. upon BL make the rectangle $BM = F$; betwixt b 44. 1. AB and BH c find out a mean proportional NO ; c 13. 6. Upon NO b make the polygone B like to the d 18. 6. right-lined figure given $ABEDC$. I say the polygone

e cor. 20. 6. lygone P so made shall be equal to F that was
f r. 6. given.

g 14. 5. For ABEDC (AL) P :: *e* AB. BH :: *f* AL.
a constr. BM. Therefore P *g* = BM. *b* = F. Which was
 to be done.

PROP. XXVI.



If from a Parallelogram ABCD, be taken away another Parallelogram AGFE, like unto the whole, and in like sort set, having also an angle common with it EAG; then is that Parallelogram about the same diagonal AC with the whole.

If you deny AC to be the common diagonal, then let AHC be it, cutting EF in H, and let HI be drawn parallel to AE. Then are Pgrs. EI, DB *a* like. *b* therefore AE.EH :: AD.DC :: *c* AE.EF. and *d* consequently EH = EF. *f* Which is absurd.

a 24. 6.


b r. def. 6.

c hyp.

d 9. 5.

f 9. ax.

PROP. XXVII.



Of all parallelograms, AD, AG applied to the same right line AB, and wanting in figure by the parallelograms CE, KI like and alike set to the Pgr. AD which is described upon the half line, the greatest is that AD which is applied to the half being like to the defect KI.

For because that GE *a* = GC, and KI added in common. *b* thence is KE = CI *c* = AM. add CG in common. *d* then is AG = to the Gnomon MBL. But the Gnomon MBL *e* = CE (AD.) Therefore AG = AD. Which was to be demonstrated.

a 43. 1.

b 2. ax.


c 36. 1.

d 2. ax.

e 9. ax.

PROP.

PROP. XXVIII.




Upon a right line given AB , to apply a parallelogram AP equal to a right-lined figure given C , deficient by a parallelogram ZR which is like to another parallelogram given D . * Now it is requisite that the * 27. 6.
right-lined figure given C , whereunto the Pgr. to be applied AP must be equal, be not greater than the Pgr. AF which is applied upon the half line, the defects being like, namely the defect of the Pgr. AF , which is applied to the half line, and the defect of the Pgr. D to be applied whose defect is to be like to the Pgr. given.

Bisect AB in E ; upon EB make the Pgr. a 18. 6.
 EG like to the Pgr. D ; and b let $EG = C + I$. b sch. 45. 1.
 c Make the Pgr. $NT = I$, and like to the Pgr. c 25. 6.
given D , or EG ; Draw the diameter FB ; Make
 $FO = KN$, and $FQ = KT$; thro' O and Q
draw the parallels SR, QZ . Then is the Pgr.
 AP that which was sought.

For the Pgrs. D, EG, OQ, NT, ZR are all d constr. e
like one to the other, and the Pgr. $EG = e$ NT 24. 6.
 $+ C = e$ $OQ + C$. f wherefore $C =$ to the e constr.
Gnomon OBQ $g = AO + PG = h$ $AO + EP$ f 3. ax.
 $= AP$, Which was to be done. g 2. ax.
 h 43. 1.

PROP. XXIX.




Upon a right line given AB . to apply a Pgr. AN equal to a right-lined figure given C , exceeding by a Pgr. OP , which shall be like to another Pgr. given D .

a 18. 6. Bisect AB in E . Upon EB a make a Pgr. EG
b 25. 6. like to D , which was given. And *b* let the Pgr. $HK = EG + C$, and like to D given, or EG . Make $FEL = c$ IH , and c $FGM = IK$. Thro' L, M draw the parallels MN and RN , and AR parallel to NM . produce ABP , GBQ . Draw the diameter FBN . Then is AN the parallelogram required.

d constr. For the pgrs. D, HK, LM, EG are *d* like, *e* therefore the pgr. OP is like to the pgr. LM , or D .
e 24. 1. Also $LM f = HK f = EG + C$. *g* Therefore $C =$
f constr. to the Gnomon ENG . But $AL h = LB k = BM$.
g 3. ax. *l* therefore $C = AN$. Which was to be done.

k 43. 1.
l 2. and 1.
 ax.

PROP. XXX.



To cut a finite right line given AB according to extreme and mean ratio ($AB. AG :: AG. GB$.)


a 11. 2.

b 17. 6.

a Cut AB in G , in such wise that $AB \times BG = AG^2$. *b*
 Then $BA. AG :: AG. GB$. Which was to be done.

PROP.

PROP. XXXI.



In right-angled triangles BAC , any figure BF described upon the side BC subtending the right angle BAC , is equal to the figures BG, AL , described upon the sides BA, AC , containing the right angle, like and alike situate to the former BF .

From the right angle BAC let down a perpendicular AD . Because that $DC. CA :: a CA. a cor. 8.6.$
 CB, b therefore $AL. BF :: DC. CB. Also, be b cor. 20.6.$
because $DB. BA :: a BA. BC, b$ therefore $BG. BF$
 $:: DB. BC. c$ therefore $AL + BG. BF :: DC + c 24. 5.$
 $DB (BC.) BC. d$ Therefore $AL + BG = BF. d schol. 14.$
Which was to be dem. 5.


Or thus: $BG. BF :: e BAq. BCq. And e AL. e 22. 6.$
 $BF :: ACq. BCq. f$ therefore $BG + AL. BF :: f 24. 5.$
 $BAq + ACq. BCq. g$ Therefore whereas $BAq + g sch. 14. 5.$
 $ACq = b BCq. b$ thence is $BG + AL = BF. Which h 47. 1,$
was to be dem.

Coroll.

From this Proposition you may learn how to add or subtract any like figures, by the same method that is used in adding and subtracting of squares, in *Schol. 47. 1.*

PROP.

PROP. XXXII.



If two triangles ABC , DCE having two sides proportional to two (AB , $AC :: DC$, DE .) be so compounded or set together at one angle ACD that their homologous sides

be also parallel AB to DC , and AC to DE .) then the remaining sides of those triangles (BC , CE) shall be found placed in one strait line.

a 29. 2.

b hyp.

c 6. 6.

d 2. ax.



e 32. 1.

f 1. ax.

g 14. 1.

For the angle A $a = ACD$ $a = D$, and AB , AC $b :: DC$, DE . c therefore the angle $B = DCE$. Therefore the angle $B + A$ $d = ACE$. But the angle $B + A + ACB$ $e = 2$ right. f therefore the angle $ACE + ACB = 2$ right. g therefore BCE is a right line. Which was to be dem.

PROP. XXXIII.



In equal circles $DBCA$, $HFGP$, the angles BDC , FHG have the same ratio with their peripheries BC , FG on which they insist; whether the angles be set at the centers (as BDC , FHG) or at the circumferences, A , E : And in like sort are the sectors BDC , FHG , because described upon the centers.

Draw

Draw the right lines BC, FG. Make $CI=CB$,
and $GL=FG=LP$, and join DI, HL, HP.

The arch $BC = CI$. *a* also the arch FG, GL , *a* 28. 3.
 LP are equal. *b* therefore the angle $BDC=CDI$, *b* 27. 3.

and the angle $FGH=GHL=LHP$. Therefore
the arch BI is as multiple of the arch BC , as the

angle BDI is of the angle BDC . And in like man-
ner is the arch EP as multiple of the arch FG ,

as the angle FHP is of the angle FHG . But if
the arch $BI \square, =, \sqsupset FP$. *c* then likewise is the

angle $BDI \square, =, \sqsupset FHP$. Therefore is the arch *c* 27. 3.
 $BC.FG :: d$ the angle $BDC.FHG$ *d* 6. def. 5.

e :: $\frac{BDC}{2}, \frac{FHG}{2}$ *e* 15. 5.
f 20. 3.

f :: A. E. Which was to be dem-

Moreover, the angle $BMC g = CNI$; *b* and *g* 27. 3.
therefore the segment $BCM = CIN$. *k* Also the *h* 24. 3.

triangle $BDC = CDI$, *l* wherefore the sector *k* 4. 1.
 $BDCM = CDIN$. After the same manner are *l* 2. ax.

the sectors FHG, GHL, LHP equal one to the
other. Therefore since accordingly as the arch

$BI \square, =, \sqsupset FGP$, so is likewise the sector
 $BDI \square, =, \sqsupset FHP$; *m* thence shall be the sector *m* 6. def. 5

$BDC.FHG ::$ the arch $BC.FG$. Which was to
be demonstrated.

Coroll.

I. Hence, As sector is to sector, so is angle to *II. 5.*
angle.

2. The angle BDC in the center is to four right
angles, as the arch BC , on which it insists, to the
whole circumference.


For as the angle BDC is to a right angle, so
is the arch BC to a quadrant. Therefore BDC is
so to four right angles as the arch BC is to four
quadrants, that is, the whole circumference. Also
the angle $A. 2$ right :: the arch BC periphery.

3. Hence, The arches IL, BC of unequal circles
which subtend equal angles, whether at the centers,
as IAL and BAC , or at the periphery are like.

For

The sixth Book of

For IL . periph. $::$ angle IAL (BAC) $\frac{1}{4}$ right.
 Also Arch BC , periph $::$ angle BAC . $\frac{1}{4}$ right.
 Therefore IL . periph $::$ BC . periph. And consequently the arches IL and BC are like. Whence



4. Two semidiameters AB , AC cut off like arches IL , BC from concentrical peripheries.

The End of the sixth Book.

THE

THE SEVENTH BOOK
OF
EUCLIDE'S ELEMENTS.

Definitions.

I. **U**Nity is that, by which every thing that is, is called One.

II. Number is a multitude composed of units.

III. One number is a part of another, the lesser of the greater, when the lesser measures the greater.

Every part is denominated from that number by which it measures the number whereof it is a part; as 4 is called the third part of 12, because it measures 12 by 3.

IV. But the lesser number is termed Parts, when it measures not the greater.

All parts whatsoever are denominated from those two numbers, by which the greatest common measure of the two numbers measures each of them; as 10 is said to be $\frac{2}{3}$ of the number 15; because the greatest common measure, which is 5, measures 10 by 2, and 15 by 3.

V. A number is Multiple (or Manifold) a greater in comparison of a lesser, when the lesser measures the greater.

VI. An even number is that which may be divided into two equal parts.

VII. But an odd number is that which cannot be divided into two equal parts; or, that which differs from an even number by an unit.

VIII. A number evenly even is that which an even number measures by an even number.

IX. But a number evenly odd is that which an even number measures by an odd number.

X. A

X. A number oddly odd is that which an odd number measures by an odd number.

XI. A prime (or first) number is that, which is measured only by an unit.

XII. Numbers prime the one to the other, are such as only an unit doth measure, being their common measure.

XIII. A composed number is that which some certain number measures.

XIV. Numbers composed the one to the other, are they, which some number, being a common measure to them both, does measure.

In this, and the preceding definition, unity is not a number.

XV. One number is said to multiply another, when the number multiplied is so often added to it self, as there are units in the number multiplying, and another number is produced.

Hence in every multiplication a unit is to the multiplier, as the multiplicand is to the product.

Obs. That many times, when any number are to be multiplied as (*A* into *B*) the conjunction of the letters denotes the product: So $AB = A \times B$, and $CDE = C \times D \times E$.

XVI. When two numbers multiplying themselves produce another, the number produced is called a plain number; and the numbers which multiplied one another, are called the sides of that number: So $2 (C) \times 3 (D) = 6 = CD$ is a plain number.

XVII. But when three numbers multiplying one another produce any number, the number produced is termed a solid number; and the numbers multiplying one another, are the sides thereof: So $2 (C) \times 3 (D) \times 5 (E) = 30 = CDE$ is a solid number.

XVIII. A square number is that which is equally equal; or, which is contained under two equal numbers. Let *A* be the side of a square; the square is thus noted, AA , or Aq . or A^2 .

XIX. A

XIX. A Cube is that number which is equally equal equally, or, which is contained under three equal numbers. Let *A* be the side of a Cube; the Cube is thus noted, *AAA*, or *Ac.* or *A³*.

In this definition, and the three foregoing, unity is a number.

XX. Numbers are proportional, when the first is as multiple of the second, as the third is of the fourth; or, the same part; or, when a part of the first number measures the second, and the same part of the third measures the fourth, equally: and on the contrary. So $A. B :: C. D$, that is, $3. 9 :: 5. 15$.

XXI. Like plane, and solid numbers are they, which have their sides proportional: *Namely, not all the sides, but some.*

XXII. A perfect number is that which is equal to all its aliquot parts.

As 6 and 28. But a number that is less than its aliquot parts is called an Abounding number; and a greater a Diminutive: so 12 is an Abounding, 15 a Diminutive number.

XXIII. One number is said to measure another, by that number, which, when it multiplies, or is multiplied by it, it produces.

In Division, a unit is to the quotient as the divisor is to the dividend. Note, that a number placed under another with a line betwixt them, signifies Division: So $\frac{A}{B}$ = A divided by B, and $\frac{CA}{B}$ = C x A divided by B.

Two numbers are called Terms or Roots of Proportion, lesser than which cannot be found in the same proportion.

Postulates or Petitions.

1. **T**hat numbers equal or manifold to any number may be taken at pleasure.
2. That a greater number may be taken than any number whatsoever.
3. That

3. That Addition, Subtraction, Multiplication, Division, and the Extractions of roots or sides of square and cube numbers, be also granted as possible.

Axiomes.

1. **W**Hatsoever agrees with one or many equal numbers, agrees likewise with the rest.

2. Those parts that are the same to the same part, or parts, are the same amongst themselves.

3. Numbers that are the same parts of equal numbers, or of the same number, are equal amongst themselves.

4. Those numbers, of whom the same number, or equal numbers, are the same parts, are equal amongst themselves.

5. Unity measures every number by the units that are in it; that is, by the same number.

6. Every number measures it self by a unity.

7. If one number, multiplying another, produce a third, the multiplier shall measure the product by the multiplicand; and the multiplicand shall measure the same by the multiplier.

Hence, *No prime number is either a plane, solid, square, or cube number.*

8. If one number measures another, that number by which it measures shall measure the same by the units that are in the number measuring, that is by the number itself that measures.

9. If a number measuring another, multiply that by which it measures, or be multiplied by it, it produces that number which it measures.

10. How many numbers soever any number measures, it likewise measures the number composed of them.

11. If a number measure any number, it also measures every number which the said number measures.

12. A number that measures the whole and a part taken away doth also measure the residue.

PROP.

PROP. I.

A....E..G..B 8 5 3 If two unequal num-
 C...F..D $\frac{5}{3}$ $\frac{3}{2}$ $\frac{2}{1}$ bers AB, CD, being given,
 H--- the lesser CD be conti-
 nually taken from the
 greater AB (and the residue EB from CD, &c.) by
 an alternate subtraction, and the number remaining
 do never measure the precedent, till the unity GB be
 taken; then are the numbers which were given AB,
 CD, prime the one to the other.

If you deny it, let AB, CD have a common
 measure, namely, the number H. Therefore H
 measuring CD, does *a* also measure AE; and *a* II. ax. 7.
 consequently the remainder EB; *a* therefore it
 likewise measures CF, and *b* so the remainder *b* 12. ax. 7.
 FD; *a* wherefore it also measures EG. But it
 measured the whole EB, and *b* therefore it must
 measure that which remains GB, a unity, itself
 being a number. *c* Which is absurd. *c* 9. ax. 7.

PROP. II.

9 6 Two numbers AB,
 A.....E.....B 15 9 6 CD being given, not
 6 3 prime the one to the
 C.....F...D $\frac{9}{2}$ $\frac{6}{1}$ $\frac{3}{2}$ other to find out their
 G--- greatest common mea-
 sure FD.

Take the lesser number CD from the greater
 AB as often as you can. If nothing remains, *a* 6. ax. 7.
 it is manifest that CD is the greatest common
 measure. But if there remains something (as EB)
 then take it out of CD, and the residue FD out
 of EB, and so forward till some number (FD)
 measure the said EB (*b* for this will be before *b* 1. 7.
 you come to a unity.) FD shall be the greatest
 common measure.

For FD *c* measures EB, and *d* therefore also CF; *c* const.
 and *e* consequently the whole CD; *d* therefore like- *d* 11. ax. 7.
 wise AE; and so measures the whole AB. Wherefore *e* 12. ax. 7.
 K it

it is evident that FD is a common measure. If you deny it to be the greatest, let there be a greater (G;) then whereas G measures CD, it must likewise measure AE, *e* and the residue EB, *g* *suppos.* *d* as also CF, *e* and by consequence the residue h *g.* *ax.* 1. FD, *g* the greater the less. *b* Which is absurd.

Coroll.

Hence, A number that measures two numbers, does also measure their greatest common measure.

PROP. III.

A 12

B 8

D 4

C 6

E .. 2

F ---

Three numbers being given, A, B, C, not prime to one another, to find out their greatest common measure E.

Find out D the greatest common measure of the two numbers A, B. If D measures C the third,

it is clear that D is the greatest common measure of all the three numbers. If D does not measure C, at least D and C will be composed the one to the other, by the Coroll of the Prop. preceding. Therefore let E be the greatest common measure of the said numbers D and C, and it appears to be the number required.

a *constr.* For E *a* measures C and D, and D measures A and B; therefore *b* E measures each of the numbers A, B, C; neither shall any greater (F) measure them; for if you affirm that, *c* then F measuring A and B, does likewise measure D their greatest common measure; and in like manner, F measuring D and C, does also measure E *c* their greatest common measure, *d* the *e* *g.* *ax.* 1. greater the less. *e* Which is absurd.

Coroll.

Hence, A number that measures three numbers, does also measure their greatest common measure.

PROP.

PROP. IV.

A.....6 Every less number *A* is of every
 B.....7 greater *B* either a part or parts.
 B.....18 If *A* and *B* be prime to one
 B.....9 another, *a* *A* shall be as many a 4. def.7.
 parts of the number *B*, as there
 are unities in *A* (as $6 = \frac{6}{7} 7$.) But if *A* mea-
 sures *B*, it is plain that *A* is a part of *B* (as 6 b 3. def.7.
 $= \frac{1}{3} 18$.) Lastly, if *A* and *B* be otherwise com-
 posed to one another, *c* the greatest common c 4. def.7.
 measure shall determine how many parts *A* does
 contain of *B*; as $6 = \frac{2}{3} 9$.

PROP. V.

A.....6 D....4
 6 6 4 4
 B.....G.....C 12. E....H....F 8.

If a number *A* be a part of a number *BC*, and
 another number *D* the same part of another number
EF; then both the numbers together (*A* + *D*) shall
 be the same part of both the numbers together (*BC*
 + *EF*), which one number *A* is of one number *BC*.

For if *BC* be resolved into its parts *BG*, *GC*,
 equal to *A*; and *EF* also into its parts *EH*, *HF*
 equal to *D*; *a* the number of parts in *BC* shall a hyp.
 be equal to the number of parts in *EF*. There-
 fore since $A + D = BG + EH = GC + HF$, thence b const. and
 $A + D$ shall be as often in $BC + EF$, as *A* is in 2. ax. 1.
BC. Which was to be dem.

Or thus. Let $a = \frac{x}{2}$, and $b = \frac{y}{2}$, then 2. $a = x$, c 2. ax. 1.
 and 2. $b = y$. wherefore 2. $a + b = x + y$. there-
 fore $a + b = \frac{x + y}{2}$. Which was to be dem.

PROP. VI.

$\begin{array}{ccc} 3 & 3 & 4 & 4 \\ A \dots G \dots B & 6 & D \dots H \dots E & 8 \\ C \dots \dots 9 & & F \dots \dots \dots 12 & \end{array}$
 If a number AB be parts of a number C , and another number DE the same parts of another number F ; then both numbers together $AB + DE$ shall be of both numbers together $C + F$ the same parts, that one number AB is of one number C .

- Divide AB into its parts AG, GB ; and DE into its parts DH, HE . The multitude of parts in both AB, DE is equal by supposition; wherefore since AG *a* is the same part of the number C , that DH is of the number F ; therefore $AG + DH$ *b* shall be the same part of the compounded number $C + F$, that one number AG is of one number C . *b* In like manner $GB + HE$ is the same part of the said $C + F$, that one number GB is of one number C . *c* Therefore $AB + DE$ is the same parts of $C + F$, that AB is of C . Which was to be dem.
- a* hyp.
- b* 5. 7.
- c* 2. ax. 7.

Or thus. Let $a = \frac{2}{3} x$, and $b = \frac{2}{3} y$, and $x + y = g$, then, because $3 a = 2 x$, and $3 b = 2 y$, is $3 a + 3 b = 2 x + 2 y = 2 g$. therefore $a + b = \frac{2}{3} g = \frac{2}{3} x + y$.

PROP. VII.

$\begin{array}{ccc} 5 & 3 & 6 & 6 \\ A \dots E \dots B & 8 & G \dots C \dots \dots F \dots D & 16 \\ 6 & 10 & & \end{array}$
 If a number AB be the same part of a number CD , that a part taken away AE is of a part taken away CF ; then shall the residue EB be the same part of the residue FD , that the whole AB is of the whole CD .

- a* 1. post. 7. Let EB be the same part of the number GC that
- b* 5. 7. AB is of CD , or AE of CF . *b* Therefore $AE + EB$ is the same part of $CF + GC$ that AE is of CF , or
- c* 6. ax. 1. AB of CD . *c* therefore $GF = CD$. Take away
- d* 3. ax. 1. CF common to both, and *d* there remains $GC =$
- e* 2. ax. 7. FD . *e* Wherefore EB is the same part of the residue FD (GC) that the whole AB is of the whole CB . Which was to be dem.

Or

Or thus. Let $a + b = x$; and $c + d = y$; and $x = 3y$, in like manner as $a = 3c$; I say $b = 3d$. For $3c + 3d = 3y = x = a + b$. take away from f 1. 2. both $3c = a$, and b there remains $3d = b$. *g hyp. Which was to be dem.*

PROP. VIII.

6 2 4 2 2 If a number AB
A.....H..G....E..L..B 16 be the same parts of
18 6 a number CD, that a
C.....F.....D 24 part taken away AE
is of a part taken
away CF; the residue also EB shall be of the re-
sidue FD the same parts, that the whole AB is of
the whole CD.

Divide AB into AG, GB, parts of the number CD; also AE into AH, HE, parts of the number CF; and take $GL = AH = HE$. *a* wherefore *a* 3. ax. 1, $HG = EL$. And because b $AG = GB$, *c* therefore *b* constr. $HG = LB$. Now whereas the whole AG is the *c* 3. ax. 1. same part of the whole CD that the part taken away AH is of the part taken away CF, *d* the *d* 7. 7. residue HG or EL shall be the same part also of the residue FD that AG is of CD. In like manner, because GB is the same part of the whole CD, that HE or GL are of CF, *d* therefore the residue LB shall be the same part of the residue FD that GB is of the whole CD. Therefore $EL + LB$ (EB) is the same parts of the residue FD, that the whole AB is of the whole CD. *Which was to be dem.*

Or thus more easily. Let $a \div b = x$, and $c + d = y$. Also $y = \frac{2}{3}x$ as well as $c = \frac{2}{3}a$; or, *e* which is *e* 9. ax. 7. the same, $3y = 2x$; and $3c = 2a$. I say $d = \frac{2}{3}b$. For $3c + 3d = 3y = 2x = 2a + 2b$ 1. 2. *b. g* Therefore $3c + 3d = 2a + 2b$. take away *g* 1. ax. 1. from each $3c = 2a$. and *b* there remains $3d = 2b$ *h hyp.* *a b. l* therefore $d = \frac{2}{3}b$. *Which was to be dem.* *k* 3. ax. 1. *l* 8. ax.

K 3

PROP.

PROP. IX.

A....4
 4 4
 B....G....C8
 5 D....5
 5 5
 E....H....F10

If a number A be a part of a number BC , and another number D the same part of another number EF ; then alternately what part or parts the first A is of the third D , the same part or parts shall the second BC be of the fourth EF .

A is supposed $\neg D$. therefore let BG, GC , and EH, HF , parts of the numbers BC, EF be equal; BG and GC to A ; and EH, HF to D . The multitude of parts is put equal in both. But it is clear that
 a 1. & x. 7. GB is a the same part or parts of EH , that GC is of
 and 4. 7. HF ; wherefore BC ($BG + GC$) is the same part
 b 5. & 6. 7. or parts of EF ($EH + HF$) that BG alone (A) is
 of EH alone (D). Which was to be dem.

Or thus. Let $a = \frac{b}{3}$ and $c = \frac{d}{3}$ or $3a = b$ and

* 15. 5. $\frac{3c}{a} = d$. then is $\frac{c^*}{a} = \frac{3c}{3a} = \frac{d}{b}$.

PROP. X.

A..G..B4
 C.....6
 5 5
 D....H....E10
 F.....15

If a number AB be parts of a number C , and another number DE the same parts of another number F ; then alternately, what part or parts the first AB is of the third DE , the same parts or part shall the second C be of the fourth F .

AB is taken $\neg DE$, and $C \neg F$. Let AG, GB , and DH, HE be parts of the numbers C and F , viz. as many in AB , as in DE . It is manifest that
 a 9. 7. AG is the same part of C , that DH is of F . a
 whence alternately AG is of DH , and likewise GB
 b 5. & 9. 7. of HE , and b so conjointly AB of DE the same
 part, or parts, that C is of F . Which was to be dem.

Or thus. Let $a = \frac{2b}{3}$, & $c = \frac{2d}{3}$. or $3a = 2b$, &
 $3c = 2d$. Then is $\frac{c}{a} = \frac{3c}{3a} = \frac{2d}{2b} = \frac{d}{b}$.

PROP.

PROP. XI.

If a part taken away AE
be to a part taken away CF ,
as the whole AB is to the
whole CD , the residue also
 EB shall be to the residue FD ,
as the whole AB is to the
whole CD .

4 3
A...E...B7
8 6

C.....F.....D14

First, let AB be $\supset CD$; a then AB is either
a part or parts of the number CD ; and likewise
 AE is b the same part or parts of CF ; c theref.
the residue EB is the same part or parts of the
residue FD , that the whole AB is of the whole
 CD . b and so $AB. CD :: EB. FD$. But if AB be
 $\subset CD$, then according to what is already shewn,
will $CD. AB :: FD. EB$. therefore by inversion
 $AB. CD :: EB. FD$.

a 4. 7.
 b 20. def. 7.
 c 7. or 8. 7.

PROP. XII.

A, 4. C, 2. E, 3. If there be divers numbers,
B, 8. D, 4. F, 6. how many soever, proportional
($A. B :: C. D :: E. F$;) then
as one of the antecedents A is to one of the conse-
quents B , so shall all the antecedents ($A+C+E$) be
to all the consequents ($B+D+F$.)

First, let A, C, E , be $\supset B, D, F$; then (be-
cause of the same proportions) a shall A be the
same part or parts of B that C is of D ; b and
likewise conjointly $A+C$ shall be the same part
or parts of $B+D$ that A alone is of B alone. In
like manner $A+C+E$ is the same part or parts of
 $B+D+F$ that A is of B . c Therefore $A+C+E.$
 $B+D+F :: A. B$. But if A, C, E , be put greater
than B, D, F , the same may be shewn by in-
version.

a 20. def. 7.
 b 5. & 6. 7

c 20. def. 7.

PROP. XIII.

A, 3. C, 4. If there be four numbers proportional
B, 5. D, 12. ($A.B::C.D.$) then alternately they
shall also be proportional, ($A.C::B.D.$)

First let A and C be $\sqsupset$ B and D, and A $\sqsupset$
a 20. def. 7. C. By reason of the same proportion a shall A
b 9. & 10. be the same part or parts of B, that C is of D.
7. Therefore alternately A is the same part or
parts of C that B is of D. and so $A.C::B.D.$
But if A be $\sqsubset$ C, and A and C supposed $\sqsupset$ B
and D, the matter will be the same by invert-
ing the proportions.

PROP. XIV.

A, 9. D, 6. If there be numbers, how many soever,
B, 6. E, 4. A, B, C, and as many more equal to
C, 3. F, 2. them in multitude, which may be com-
pared two and two in the same propor-
tion ($A.B::D.E.$ and $B.C::E.F.$;) they shall also, of
equality, be in the same proportion ($A.C::D.F.$)

a 13. 7. For because $A.B::D.E.$ a therefore alternately
is $A.D::B.E::a.C.F.$ a and so again by chan-
ging $A.C::D.F.$ Which was to be dem.

PROP. XV.

1. D.. If a unit measure any number B,
B... 3. E..... 6. and another number D do equally
measure some other number E; al-
ternately also shall a unit measure the third number
D, as often as the second B does the fourth E.

a 9. 7. For seeing 1 is the same part of B, that D is
of E; a therefore alternately shall 1 be the
same part of D, that B is of E. Which was to be
demonstrated.

PROP.

PROP. XVI.

B, 4. A, 3. If two numbers, *A*, *B*, multiplying themselves the one into
 A, 3. B, 4. the other, produce any numbers
 AB, 12. BA, 12. *AB*, *BA*; the numbers produced
AB and *BA* shall be equal the one to the other.

For because $AB = A \times B$, *a* therefore shall *r* ^{a 15.def.7.}
 be as often in *A*, as *B* in *AB*, *b* and by conse- ^{b 15.7.}
 quence alternately *r* shall be as often in *B* as *A*
 in *AB*. But for that $BA = B \times A$, *a* therefore
 shall *r* be as often in *B*, as *A* in *BA*. therefore
 as often as *r* is in *AB*, so often is *r* in *BA*. and
c so $AB = BA$. Which was to be dem. c 4.ax.7.

PROP. XVII.

A, 3. If a number *A* multiplying
 B, 2. C, 4. two numbers *B*, *C*, produce other
 AB, 6. AC, 12. numbers, *AB*, *AC*; the numbers
 produc'd of them shall be in the
 same proportion that the numbers multiplied are.
 (*AB*. *AC* :: *B*. *C*.)

For being $AB = A \times B$, *a* therefore shall *r* be as *a* 15.def.7.
 often in *A*, as *B* in *AB*. *a* Likewise because AC
 $= A \times C$, therefore shall *r* be as often in *A*, as *C*
 in *AC*. and so also *B* as often in *AB* as *C* in *AC*.
b wherefore *B*. *AB* :: *C*. *AC*. *c* and therefore also *b* 20.def.7.
 alternately *B*. *C* :: *AB*. *AC*. Which was to be dem. c 13.7.

PROP. XVIII.

C, 5. C, 5. If two numbers *A*, *B*, mul-
 A, 3. B, 9. tipling any number *C*, produce
 AC, 15. BC, 45. other numbers *AC*, *BC*; the
 numbers produced of them shall
 be in the same proportion that the numbers multiply-
 ing are. (*A*. *B* :: *AC*. *BC*.)

For a $AC = CA$, and $BC = CB$; so the same *a* 16. 7.
C multiplying *A* and *B* produces *AC* and *BC*.
b therefore *A*. *B* :: *AC*. *BC*. Which was to be dem. b 17.7.

Schol.

Schol.

Hence is deduced the vulgar manner of reducing fractions ($\frac{3}{4}$, $\frac{2}{5}$) to the same denomination. For multiply 9 both by 3 and 5, and they produce $\frac{27}{4} = \frac{3}{4}$; because by this, 3. 5 :: 27. 45. Likewise multiply 5 by 7 and 9; there arises $\frac{45}{9} = \frac{2}{9}$; because 7. 9 :: 35. 45.

P R O P. XIX.

A, 4. B, 6. C, 8. D, 12. If there be four numbers in proportion (A. B :: C. D) the number produced of the first and fourth (AD) is equal to the number which is produced of the second and third (BC.) And if the number which is produced of the first and fourth (AD) be equal to that produced of the second and third (BC) those four numbers shall be in proportion (A. B :: C. D.)

a 17. 7.

b hyp.

c 18. 7.

d 9. 5.

e hyp.

f 7. 5.

g 17. 7.

h 18. 7.

k 11. 5.

1. Hyp. For AC, AD a :: C. D b :: A. B c :: AC. BC. d therefore AD = BC. Which was to be dem.

2. Hyp. Because e AD = BC, therefore AC. AD f :: AC. BC. But AC. AD g :: C. D. and AC. BC h :: A. B. k therefore C. D :: A. B. Which was to be demonstrated.

P R O P. XX.

A. B. C. If there be three numbers in proportion (A. B :: B. C) the number contained under the extremes (AC) is equal to the square made of the middle (BB.)

And if the number contained under the extremes be equal to that (Bq.) produced of the middle, those three numbers shall be in proportion ($\frac{A}{B} :: \frac{B}{C}$)

a 1. ax. 7.

b 19. 7.

1. Hyp. For take D = B. a therefore A. B :: D (B.) C. b wherefore AC = BD, a or BB. Which was to be demonstrated.

2. Hyp.

2. Hyp. Because $AC : c = BD$, d therefore $A. c$ hyp.
 $B :: D (B) C$. Which was to be dem. d 19. 7.

PROP. XXI.

$A \dots G \dots B 5$. $E \dots \dots \dots 10$ Numbers AB , CD ,
 $C \dots H \dots D 3$. $F \dots \dots \dots 6$ being the least of all
that have the same
proportion with them (E , F) do equally measure the
numbers, E , F , having the same proportion with
them; the greater AB the greater E , and the lesser
 CD the lesser F .

For $AB. CD a :: E. F. b$ therefore alternately a hyp.
 $AB.E :: CD. F. c$ therefore AB is the same part b 13. 7.
or parts of E that CD is of F . Not parts; for if c 20. def. 7.
so, then let AG , GB be parts of the number E ;
and CH , HD , parts of the number F . c therefore
 $AG. E :: CH. F.$ and by inversion $AG. CH d :: d$ 13. 7.
 $E. F e :: AB. CD.$ therefore AB, CD are not the e hyp.
least in their proportion; which is contrary to
the Hypothesis. Therefore, &c.

PROP. XXII.

$A, 4$. $D, 12$. If there be three numbers A ,
 $B, 3$. $E, 8$. B, C ; and other numbers equal
 $C, 2$. $F, 6$. to them in multitude, D, E, F ;
which may be compared two and
two in the same proportion: And if also the pro-
portion of them be perturbed ($A. B :: E. F.$ and $B.$
 $C :: D. E$.) then of equality they shall be in the same
proportion ($A. C :: D. F$.)

For because $A, B a :: E, F$, therefore shall $AF a$ hyp.
 $= BE$; and because $B, C :: a D, E$, b therefore b 19. 7.
 $BE = CD. c$ and consequently $AF = CD. d$ Where- c 1. ax. 1.
fore $A, C :: D, F$. Which was to be dem. d 19. 7.

PROP.

P R O P. XXIII.

A, 9. B, 4. Numbers prime the one to the
 C----D--- other, A, B, are the least of all
 E-- numbers that have the same proportion with them.

If it be possible, let C and D be less than A and B, and in the same proportion; *a* therefore C measures A equally as D measures B, namely, by the same number F; and so C shall be *b* as often in A as *r* is in E; *c* likewise alternately, E as often in A as *r* in C. By the like inference as many times as *r* is in D, so many times shall E be in E. Therefore E measures both A and B; which consequently are not prime the one to the other, *contrary to the Hypothesis*.

P R O P. XXIV.

A, 9. B, 4. Numbers A, B, being the least
 C--- of all that have the same proportion
 D--- E--- with them, are prime the one to the others.

If it be possible, let A and B have a common measure C; and let the same measure A by D, and B by E; *a* therefore $CD=A$, *b* and $CE=B$.
a 9. *ax*. 7. *b* 17. 7. Wherefore $A:B::D:E$. But D and E are lesser than A and B, as being but parts of them. Therefore A and B are not the least in their proportion, *against the Hypothesis*.

P R O P. XXV.

A, 9. B, 4. If two numbers A, B, be prime
 C, 3. D-- the one to the other, the number C measuring one of them A, shall be prime to the other number B.

For if you affirm any other D to measure the numbers B and C, *a* then D measuring C does also measure A; and consequently A and B are not prime the one to the other. *Which is against the Hypothesis*.

P R O P.

PROP. XXVI.

A, 5.	C, 8.	<i>If two numbers; A, B, be prime to any number C, the number also produced of them A, B shall be prime to the same C.</i>
B, 3.		
<u>AB, 15.</u>	E - - - -	
	F - - - -	

If it be possible, let the number E be a common measure to AB, and C; and let $\frac{AB}{E}$ be = F; *aa 9. ax. 7.*

thence $AB = EF$; *b* wherefore also $E : A :: B : F$. *b* 19. 7.
But because A is prime to C , which E measures, *c* 25. 7.
c therefore E and A are prime to one another, *d* 23. 7.
and so least in their own proportion, *e* and consequently they must measure B and F ; namely *e* 21. 7.
 F shall measure B , and A shall measure F .
Therefore seeing E measures both B and C , they
shall not be prime to one another: *Contrary*
to the Hypothesis.

PROP. XXVII.

A, 4. B, 5.
Aq, 16.
D, 4.

If two numbers A, B, be prime to one another, that also which is produc'd of one of them (Aq) shall be prime to the other B.

Take $D = A$; therefore each of D , and A a 1. ar. 7.
are prime to B . b wherefore AD or Aq is prime b 26. 7.
to B . Which was to be dem.

PROP. XXVIII.

*A, 5. C, 4.
B, 3. D, 2.
AB, 15. CD, 8.
If two numbers A, B, be prime
to two numbers C, D, each to ei-
ther of both, the numbers also
produced of them AB, CD shall
be prime to one another.*

For being A and B are prime to C, therefore a shall AB also be prime to the same. And by the same

- b 26. 7. same reason shall AB be prime to D. b Therefore AB is prime to CD. Which was to be dem.

P R O P. XXIX.

- A, 3. B, 2. If two numbers A, B, be prime
 Aq, 9. Bq, 4. to one another, and each multiply-
 Ac, 27. Bc, 8. ing himself produce another number
 (Aq, and Bq;) then the numbers
 produced of them (Aq, Bq) shall be prime to one ano-
 ther. And if the numbers given at first, A, B, multiply-
 ing the said produced numbers (Aq, Bq) produce others
 (Ac, Bc) those numbers also shall be prime to one ano-
 ther: And this shall ever happen about the extremes.
- a 27. 7. For because A is prime to B, a therefore Aq
 shall be prime to B. and Aq being prime to B,
 a therefore Aq shall also be prime to Bq. Again,
 because A is as well prime to B and Bq. as Aq
 is to the said B and Bq, b therefore shall A x
 Aq, that is, Ac, be prime to B x Bq, that is,
 to Bc: And so forth of the rest.
- b 28. 7.

P R O P. XXX.

- 8 5 If two numbers AB,
 A.....B.....C 1; D---- BC, be prime the one to
 the other, then both ad-
 ded together (AC) shall be prime to either of them
 AB, BC. And if both added together AC be prime to
 any one of them AB, the numbers also given in the
 beginning AB, BC shall be prime to one another.
1. Hyp. For if you would have AC, AB to be
 a 12. ax. 7. composed, let D be the common measure: a
 this shall measure the residue BC: and there-
 fore AB, BC, are not prime to one another;
 which is against the Hypothesis.
2. Hyp. AC, AB being taken for prime to one
 another, let D be the common measure of AB,
 BC. b But seeing that measures the whole AC,
 b 10. ax. 7. therefore AC, AB, are not prime to one ano-
 ther; contrary to the Hypothesis.

Cgroll

Hence
of two,
to the

A, 5.

For
both A
ber; cA, 4.
B, 6.

AB, 2

then fl
or B,
Sup

ber A

D. A
fore l
e and
meafA, 1
B, 2.L
let t
For
num
quer
of t

Coroll.

Hence, A number which being compounded of two, is prime to one of them, is also prime to the other.

PROP. XXXI.

A, 5. B, 8. *Every prime number A is prime to every number B, which it measures not.*

For if any common measure does measure both A, B, a then A will not be a prime number; *contrary to the Hypothesis.*

a 11. def. 7.

PROP. XXXII.

A, 4. D, 3. *If two numbers A, B, multiplying one another produce another AB, and some prime number D measure the number produced of them AB; then shall it also measure one of those numbers, A or B, which were given at the beginning.*

Suppose the number D not to measure the num-

ber A, and let $\frac{AB}{D} = E$. a then $AB = DE$, b whence a 9. ax. 7.

D. A :: B. E. c But D is prime to A; d there- b 19. 7.
fore D and A are the least in their proportion; c hyp. and
e and consequently D measures B as often as A 31. 7.
measures E. Which was to be dem. d 23. 7.
e 21. 7.

PROP. XXXIII.

A, 12. *Every composed number A is mea-*
B, 2. *sured by some prime number B.*

Let one or more numbers a measure A, of which let the least be B; that shall be a prime number: a 13. def. 7.
For if it be said to be composed, then some a lesser number shall measure it, b which shall also conse- b 11. ax. 7.
quently measure A. Wherefore B is not the least of them which measure A, *contrary to the Hyp.*

PROP.

P R O P. XXXIV.

A, 9. Every number A , is either a prime, or measured by some prime number.

For A is either necessarily a prime or a composed number. If it be a prime, 'tis that we affirm. If composed, a then some prime number measures it. Which was to be dem.

P R O P. XXXV.

A, 6. B, 4. C, 8. H - I - K ----
D, 2. L ---
E, 3. F, 2. G, 4.

How many numbers soever A, B, C , being given, to find the least numbers E, F, G , that have the same proportion with them.

a 23. 7. If A, B, C , be prime to one another, a they shall be the least in their proportion. If they be composed, b let their greatest common measure be D , which let measure them by E, F, G . These are then least in the proportion A, B, C .
b 3. 7. For $D \times E, F, G$, c produces ABC , d therefore they are all in the same proportion. But allow other numbers H, I, K to be the least in the same proportion; e which shall therefore equally
e 21. 7. measure ABC , namely by the number L . f there-
f 9. ax. 7. fore $L \times H, I, K$, shall produce A, B, C , g and con-
g 1. ax. 1. sequently $ED = A = HL$. h from whence $E. H ::$
h 19. 7. k suppos. $L. D$. But $E \nmid K$; l therefore $L \nmid D$, and so
l 20. def. 7. D is not the greatest common measure of A, B, C . Which is against the Hypothesis.

Coroll.

Hence, The greatest common measure of how many numbers soever, does measure them by the numbers which are least of all that have the same proportion with them. Whereby appears the vulgar method of reducing fractions to the least terms.

P R O P.

PROP. XXXVI.

Two numbers being given *A*, *B*, to find out the least number which they measure.

A, 5. *B*, 4. 1. *Case*. If *A* and *B* be prime the
AB, 20. one to the other, *AB* is the num-
D----- ber required. For it is manifest
E--- *F*--- that *A* and *B* measure *AB*. If it
be possible, let *A* and *B* measure
some other number *D* $\neg$ *AB*, if you please by
E, and *F*. *a* therefore *AE* $\neg$ *D* $\neg$ *BF*, *b* and so *A*. *a* 9. *ax*. 7.
B :: *F*. *E*. But because *A* and *B* *c* are prime the *c* 1. *ax*. 1.
one to the other, *d* and so least in their propor- *b* 19. 7.
tion, *A* shall *e* equally measure *F* as *B* does *E*. *c* *hyp*.
But *B*. *E* *f* :: *AB*. *AE* (*D*) *g* Therefore *AB* shall *d* 23. 7.
also measure *D*, which is less than it self. Which *e* 21. 7.
is absurd. *f* 17. 7.
g 20. *def*. 7.

A, 6. *B*, 4. *F*--- 2. *Case*. But if *A* and
C, 3. *D*, 2. *G*--- *H*--- *B* be composed one to
AD, 12. another, *h* let there be *h* 35. 7.
found *C* and *D* the
least in the same proportion. *k* therefore *AD* $\neg$ *BC*; *k* 19. 7.
and *AD* or *BC* shall be the number sought for.

For it is *l* plain that *B* and *D* do measure *AD* *l* 7. *ax*. 7.
or *BC*. Conceive *A* and *B* to measure *F* $\neg$ *AD*,
namely *A* by *G*, and *B* by *H*. *m* therefore *AG* = *m* 9. *ax*. 7.
F = *BH*. *n* whence *A*. *B* :: *H*. *G* *o* :: *C*. *D*. *p* and *n* 19. 7.
consequently *C* equally measures *H* as *D* does *G*. *o* *constr*.
But *D*. *G* *q* :: *AD*. *AG* (*F*) therefore *AD* *r* mea- *p* 21. 7.
sures *F*, the greater the less. Which is absurd. *q* 17. 7.
r 20. *def*. 7.

Coroll.

Hence, If two numbers multiply the least
that are in the same proportion, the greater the
less, and the less the greater, the least number
which they measure shall be produced.

L

PROP.

PROP. XXXVII.

A, 2. B, 3. *If two numbers A, B, measure*
 E 6 *any number C, D, the least num*
 C ---- F ---- D *ber which they measure E shall al*
so measure the same CD.

If you deny it, take E from CD as often as
you can, and leave FD $\sqsupset$ E. therefore seeing
A and B a measure E, b and E measures CF,
likewise A and B will measure CF. But a they
measure the whole CD; d therefore also they
measure the residue FD; and consequently E is
not the least which A and B measure; contrary
to the Hypothesis.

a hyp.
 b constr.
 c 11. ax. 7.
 d 12. ax. 7.

PROP. XXXVIII.

A, 3. B, 4. C, 6. *Three numbers being given*
 D, 12. *A, B, C, to find out the least*
which they measure.

a 36. 7. *a Find D to be the least that two of them A*
and B do measure; which if the third C do
also measure, it is manifest that D is the num
ber sought for. But if C do not measure D, let
E be the least that C and D do measure. E
shall be the number required.

A, 2. B, 3. C, 4. *For it appears by the 11.*
 D, 6. E, 12. *ax. 7. that A, B, C measure*
 F --- *E; and it is easily shewn*
that they measure no other

b 37. 7. *less than F. For if you affirm they do, b then D*
measures F, b and consequently E measures the
same F, the greater the less. Which is absurd.

Coroll.

Hence it appears, that if three numbers mea-
 sure any number, the least also, which they
 measure, shall measure the same.

PROP.

PROP. XXXIX.

A, 12. If any number *B* measure a number
B, 4. C, 3. *A*, the number measured *A*, shall
have a part *C* denominated of the
number measuring *B*.

For because $\frac{A}{B} a = C$, *b* shall $A = BC$. *c* there- *a* hyp.
b 9. ax. 7.
fore $\frac{A}{C} = B$. Which was to be dem. *c* 7. ax. 7.

PROP. XL.

A, 15. If a number *A* have any part
B, 3. C, 5. whatsoever *B*, the number *C*, from
which the part *B* is denominated,
shall measure the same.

For being $BC a = A$, *b* thence $\frac{A}{C} = B$. Which *a* hyp. $\mathcal{E}$
b 9. ax. 7.
was to be dem. *b* 7. ax. 7.

PROP. XLI.

$\frac{1}{2}$ G, 12. To find out a number *G*, which being
 $\frac{1}{3}$ H --- the least, contains the parts given
 $\frac{1}{4}$ $\frac{1}{2}, \frac{1}{3}, \frac{1}{4}$.

a Let *G* be found the least which the deno- *a* 38. 7.
minators, 2, 3, 4, measure; *b* it is evident that *b* 39. 7.
G has the parts $\frac{1}{2}, \frac{1}{3}, \frac{1}{4}$. If it be possible let *H*
□ *G* have the same parts; *c* therefore 2, 3, 4, *c* 40. 7:
measure *H*; and so *G* is not the least which
2, 3, 4 measure: against the constr.

The End of the seventh Book,

THE EIGHTH BOOK OF EUCLIDE'S ELEMENTS

PROP. I.

A, 8. B, 12. C, 18. D, 27.
E-F--G---H----

I*F there be divers numbers how many soever in continual proportion, A, B, C, D, and their extremes A, D, prime to one another; then the numbers A, B, C, D, are the least of all numbers that have the same proportion with them.*

For, if it be possible, let there be as many others E, F, G, H, less than A, B, C, D, and in the same proportion with them. *a* Therefore of equality $A.D :: E.H$. and consequently A and D are prime numbers, *b* and so the least in their proportion, *c* equally measuring E and H, which are less than themselves. *Which is absurd.*

a 14. 7.

b 23. 7.

c 21. 7.

PROP. II.

I.

A, 2. B, 3.

Aq, 4. AB, 6. Bq, 9.

Ac, 8. AqB, 12. ABq, 18. Bc, 27.

To find out the least numbers continually proportional, as many as shall be required, in the proportion given of A to B.

Let A and B be the least in the proportion given; then Aq, AB, Bq, shall be three last in the same continual proportion that A is to B.

For

For AA. AB $a :: A. B a :: AB. BB$. Likewise a 17. 7.
because A and B are b prime to one another, cb 24. 7.
shall Aq, Bq, be also prime to one another, dc 29. 7.
and so Aq, AB, Bq, are $::$ the least in the pro- d 1. 8.
portion of A to B.

Moreover, I say Ac, AqB, ABq, Bc, are the
four least in the proportion of A to B. For AqA.
AqB $e :: A. B. e :: ABA$ (AqB.) $ABB. e$ and $A. e$ 17. 7.
 $B :: ABq. BBq$ (Bc.) Therefore since Ac, and Bc, f 29. 7.
are f prime to one another, likewise g shall Ac, g 1. 8.
AqB, ABq, Bc be the four least $::$ in the pro-
portion of A to B. In the same manner may you
find out as many proportional numbers as you
please. Which was to be done.

Coroll.

1. Hence, If three numbers, being the least,
are proportional, their extremes shall be squares;
if four, cubes.

2. How many extremes proportional soever there
be, being by this prop. found to be the least in the
given proportion, they are prime to one another.

3. Two numbers, being the least in the given
proportion, do measure all the mean numbers
whatsoever of the least in the same proportion;
because they arise from the multiplication of
them into certain other numbers.

4. Hence also it appears by the construction,
that the series of numbers 1, A, Aq, Ac; 1, B, Bq.
Bc; Ac, AqB, ABq, Bc consists of an equal multi-
tude of numbers; and consequently, the extreme
numbers of how many soever the least continually
proportionals are the last of as many other conti-
nually proportionals from a unite, as the extreme
Ac, Bc, of the continually proportionals Ac, AqB.
ABq, Bc, are the least of as many proportionals
from a unite, 1, A, Aq, Ac; and 1, B, Bq, Bc.

5. 1, A, Aq, Ac; and B, BA, BAq; and Bq,
ABq, are $::$ in the proportion of 1 to A. Also
B, Bq, Bc; and A, AB, ABq; and Aq, AqB are
 $::$ in the proportion of 1 to B.

L 3

PROP.

PROP. III.

A, 8. B, 12. C, 18. D, 28. *If there be numbers continually proportional, how many soever, A, B, C, D, being also the least of all that have the same proportion with them; their extremes A, D, are prime to one another.*

a 2. 8.

For if there be a found as many numbers the least in the proportion of A to B, they shall be no other than A, B, C, D; therefore, by the second Coroll. of the precedent Prop. the extremes A and D are prime to one another. Which was to be dem.

PROP. IV.

A, 6. B, 5. C, 4. D, 3. *Proportions how many soever being given, in the least numbers (A, B, C, D) to find out the least numbers continually proportional in the proportions given.*

I -- K --- L ---

- a 36. 7. a Find out E the least number which B and C
 b 3. post. 7. do measure; and let B measure E b as often as A
 does another F, viz. by the same number H. b
 Also let C measure the said E as often as D mea-
 sures another G. then F, E, G, shall be the least
 c 9. ax. 7. in the proportions given. For $AH : c = F$, and
 d 18. 7. $BC : c = E$; therefore $A : B :: AH : BH :: F : E$.
 e 7. 5. In like manner $C : D :: E : G$; therefore F, E, G,
 are continually proportional in the proportions
 given. And they are moreover the least in the
 said proportions: For conceive other numbers,
 f 21. 7. I, K, L, to be the least; f then A and B must e-
 qually measure I and K, f and C and D likewise
 K and L; and so B and C measure the same K.
 g 37. 7. Wherefore also E measures the same number
 K, which is less than it self. Which is absurd.

A, 6.

A, 6. B, 5. C, 4. D, 3. E, 5. F, 7.

H, 24. G, 20. I, 15. K, 21.

But three proportions being given, A to B, C to D, and E to F; find out as before three numbers H, G, I, the least continually in the proportions of A to B, and C to D. Then if E measures H take another number K which may be equal to H 3. post. 7. measur'd by F; and those four numbers H, G, I, K, shall be continually the least in the given proportions; which we need go no other way to prove than we did in the first part.

A, 6. B, 5. C, 4. D, 3. E, 2. F, 7.

H, 24. G, 29. I, 25.

M, 48. L, 40. K, 30. N, 105.

If E do not measure I, let K be the least which E and I do measure; and as often as I measures K, let G as often measure L, and H also M. so likewise let F measure N as often as E measures K. The four numbers, M, L, K, N shall be least, continually in the given proportions; which we may demonstrate as before.

PROPOSITION V.

C, 4. E, 3. *Plane numbers CD,*
D, 6. F, 16. ED, 18. *EF, are in that proportion to one another which is composed of their sides.*
 $CD, 24. EF, 48.$

For because $CD. ED :: C.E; a$ and $ED. EF :: a$ 17. 7.

D. F. & $\frac{CD}{EF} b = \frac{CD}{ED} + \frac{D}{E} c$ then shall be the proportion b 20. def. 5.

portion $\frac{CD}{EF} = \frac{C}{E} + \frac{D}{F}.$ Which was to be dem.

PROPOSITION VI.

A, 16. B, 24. C, 36. D, 54. E, 81. *If there be numbers continually proportional how many soever, A, B, C, D, E, and the first A do not measure the second B, neither shall any of the other measure any one of the rest.*
F, 4. G, 6. H, 9.

- a 20. def. 7. Because A does not measure B, *a* neither shall any one measure that which next follows; being A.B :: B.C :: C.D. &c. *b* Take three numbers, F, G, H, the least in the proportion of A to B, therefore since A does not measure B, *a* neither shall F measure G. *c* therefore F is not a unit. But F and H are prime to one another; and so, *d* being of equality A. C :: F. H. and F does not measure H, *a* neither shall A measure C; and consequently neither shall B measure D, nor C measure E, &c. because A. C *e* :: B. D *e* :: C. E, &c. In like manner four or five numbers being taken the least in the proportion of A to B, it will appear that A does not measure D and E; nor does B measure E and F, &c. Wherefore none of them shall measure any other. Which was to be dem.

P R O P. VII.

A, 3. B, 6. C, 12. D, 24. E, 48.

If there be numbers continually proportional how many soever A, B, C, D, E, and the first A measure the last E, it shall also measure the second B.

- a 6. 7. If you deny that A measures B, *a* then neither shall it measure E; Which is contrary to the Hyp.

P R O P. VIII.

A, 24. C, 36. D, 54. B, 81. If between two G, 8. H, 12. I, 18. K, 27. numbers A, B, there E, 32. L, 48. M, 72. F, 108. fall mean proportional numbers in continual proportion C, D; as many mean continually proportional numbers as fall between them, so many also mean continually proportional numbers shall fall between two other numbers E, F, which have the same proportion with them. (L, M.)

- a 35. 7. *a* Take G, H, I, K, the least :: in the proportion of A to C; *b* of equality shall G.K :: A.B *c* :: E.F. But G, and K *d* are prime to one another. *e*
 b 14. 7. Wherefore G measures E as often as K does F.
 c hyp. Let H measure L, and I likewise M by the same
 d 3. 8. number. *f* therefore E, L, M, F, are in such proportion as G, H, I, K, that is as A, B, C, D. Which
 e 21. 7. was to be dem.
 f constr.

PROP.

P R O P. IX.

I.
 E,2. F,3.
 G,4. H,6. I,9.
 A,8. C,12. D,18. B,17.
 If two numbers A, B
 be prime to one another,
 and mean numbers in
 continual proportion C, D
 fall between them; as ma-
 ny mean numbers in continual proportion as fall be-
 tween them, so many means also (E, G ; and F, I)
 shall fall in continual proportion between either of
 them and a unity.

It is evident that $1, E, G, A$, and $1, F, I, B$,
 are $\therefore$, and as many as A, C, D, B , namely by
 the 4. Coroll. 2. 8. Which was to be dem.

P R O P. X.

A,8. I,12. K,18. B,27.
 E,4. DF,6. G,9.
 D,2. F,3.
 I.
 If between two numbers
 A, B , and a unit, numbers
 continually proportional
 $(E, D, \text{and } F, G,)$ do fall,
 how many mean numbers
 in continual proportion fall between either of them
 and a unit, so many means also shall fall in conti-
 nual proportion between them, I, K .

For E, DF, G , and $A, DqF (I) DG, (K,) B$,
 are $\therefore$ by 2. 8. therefore, &c.

P R O P. XI.

A,2. B,3.
 Aq,4. AB,6. Bq,9.
 Between two square numbers,
 Aq, Bq , there is one mean pro-
 portional number AB : and
 the square Aq to the square Bq is in double propor-
 tion of that of the side A to the side B .

a It is manifest that Aq, AB, Bq , are $\therefore$; b and a 17. 7.
 consequently also $\frac{Aq}{Bq} = \frac{A}{B}$ doubly. W.W. to be dem. b 10. def. 7.

PROP.

P R O P. XII.

Ac, 27. AqB, 36. ABq, 48. Bc, 64.

A, 3. B, 4.

Aq, 9. AB, 12. Bq, 16.

Between two
cube numbers,
Ac, Bc, there
are two mean

proportional numbers AqB, ABq: and the cube Ac
is to the cube Bc in treble proportion of that in
which the side A is to the side B.

a 2. 8. For Ac, AqB, ABq, Bc, are $\therefore$ in the propor-
b 10. def. 5. tion of A to B; b and therefore $\frac{Ac}{Bc} = \frac{A}{B}$ trebly.

Which was to be dem.

P R O P. XIII.

A, 2. B, 4. C, 8.

Aq, 4. AB, 8. Bq, 16. BC, 32. Cq, 64.

Ac, 8. AqB, 16. ABq, 32. Bc, 64. BqC, 128. BCq, 256.
(Cc, 512.

If there be numbers in continual proportion how
many soever A, B, C; and every of them multiplying
it self produce certain numbers; the numbers produ-
ced of them Aq, Bq, Cq, shall be proportional: And
if the number first given A, B, C, multiplying their
products Aq, Bq, Cq, produce other numbers, Ac, Bc,
Cc, they also shall be proportional; and this shall
ever happen to the extremes.

a 2. 8. For Aq, AB, Bq, BC, Cq a are $\therefore$ b therefore
b 14. 7. of equality Aq.Bq::Bq.Cq. Which was to be dem.

a Also Ac, AqB, ABq, Bc, BqC, BCq, Cc, are
 $\therefore$; b therefore likewise of equality Ac. Bc::
Bc. Cc. Which was to be dem.

P R O P. XIV.

Aq, 4. AB, 12. Bq, 36.

A, 2.

B, 6.

If a square number Aq
measure a square number
Bq, the side also of the one
(A) shall measure the side of the other (B:) and if the
side of one square A measure the side of another B, the
square Aq shall likewise measure the square Bq.

1. Hyp.

1. *Hyp.* For $Aq \cdot AB :: AB \cdot Bq$, therefore see-22. $\text{\textcircled{E}}$ 11.8
ing by the Hypothesis Aq measures Bq , b it b 7.8.
shall measure also AB . But $Aq \cdot AB :: A \cdot B$. c 20. *def.* 7.
therefore A measures B . Which was to be dem.

2. *Hyp.* A measures B . c therefore Aq shall as
well measure AB , c as AB measures Bq ; d and con-d 11. *ax.* 7.
sequently Aq measures Bq . Which was to be dem.

PROP. XV.

$A, 2.$ $B, 6.$ If a cube num-
 $Ac, 8.$ $AqB, 24.$ $ABq, 72.$ $Bc, 216.$ ber Ac measures a
cube number Bc ,
then the side of the one (A) shall measure the side
of the other (B .) And if the side A of one cube Ac
measure the side B of the other Bc , also the cube
 Ac shall measure the cube Bc .

1. *Hyp.* For Ac, AqB, ABq, Bc a are $∴$, there- 22. $\text{\textcircled{E}}$ 12.8
fore Ac , b measuring the extreme Bc , shall also b *hyp.*
 c measure the second AqB . But $Ac \cdot Aq \cdot B :: A \cdot c$ 7.8.
 $B \cdot d$ therefore A shall also measure B . d 20. *def.* 7.

2. *Hyp.* A measures B ; d therefore Ac mea-
sures AqB , which also measures ABq , and that c 11. *ax.* 7.
 Bc ; e therefore Ac shall measure Bc . Which
was to be dem.

PROP. XVI.

$Aq, 4.$ $B, 9.$ If a square number Aq do not mea-
 $Aq, 16.$ $Bq, 81.$ sure a square number Bq , neither shall
the side of the one A measure the side
of the other B : And if A the side of the one square
 Aq do not measure B the side of the other Bq , nei-
ther shall the square Aq measure the square Bq .

1. *Hyp.* For if you affirm that A measures B , a 14. 8.
then Aq also shall measure Bq . against the *Hyp.*

2. *Hyp.* If you maintain Aq to measure Bq ;
 a then likewise A shall measure B . contrary to
the Hypothesis.

PROP.

PROP. XVII.

- A, 2. B, 3. *If a cube number Ac do not mea-*
 Ac, 8. Bc, 27. *sure a cube number Bc, neither*
shall the side of one A, measure the
side of the other B: And if A the side of one cube
Ac do not measure B the side of the other Bc, neither
shall the cube Ac measure the cube Bc.
- a 15. 8. 1. Hyp. Let A measure B; a then Ac shall
 measure Bc. against the Hyp.
 2. Hyp. Let Ac measure Bc; then A shall mea-
 sure B; which is also against the Hypothesis.

PROP. XVIII.

- C, 6. D, 2. *Between two like plane num-*
 CD, 12. *bers CD and EF there is one*
 E, 9. F, 3. DE, 18. *mean proportional number DE:*
 EF, 27. *And the plane CD is to the*
plane EF in double proportion
of that which the side C has to the homologous side
(or of like proportion) E.
- * 21. def. 7. Being * by the Hypoth. C. D :: E. F. there-
 fore by inversion C. E :: D. F. But C. E a :: CD.
 a 17. 7. DE; a and D. F :: DE. EF. b therefore CD. DE ::
 b 11. 5. DE. EF. c Wherefore the proportion of CD to
 c 10. def. 5. EF is double to that of CD to DE, that is, to
 the proportion of C to E, or D to F.

Coroll.

Hence it is apparent, that between two like
 plane numbers there falls one mean proportional
 in the proportion of the homologous sides.

PROP.

P R O P. XIX.

CDE, 30. DEF, 60. FGE, 120. FGH, 240.

CD, 6. DF, 12. FG, 24.

C, 2. D, 3. E, 5. F, 3. G, 6. H, 10.

Between two like solid numbers CDE, FGH, there are two mean proportional numbers DFE, FGE. And the solid CDE is to the solid FGH, in treble proportion of that which the homologous side C has to the homologous side F.

Whereas by the * hyp. C.D :: F.G, & D.E :: * 21.def.7. G.H. therefore *a* by inversion shall C.F :: D. *a* 13. 7. G :: *a* E.H. But CD.DF *b* :: C. F, and DF.FG *b* 17. 7. *b* :: D. G; *c* wherefore CD.DF :: DF.FG :: *c* 11. 5. E.H. *d* and accordingly CDE. DFE :: DFE. *d* 17. 7. FGE :: E. H :: FGE. FGH. Therefore between CDE. FGH, fall two mean proportionals DFE, FGE. *e* And so it is plain that the proportion *e* 10.def.5. of CDE to FGH is treble to that of CDE to FDE, or C to F. *Which was to be dem.*

Coroll.

Hereby it is manifest, that between two like solid numbers there fall two mean proportionals in the proportion of the homologous sides.

P R O P. XX.

A, 12. C, 18. B, 27. *If between two numbers D, 2. E, 3. F, 6. G, 9. A, B, there fall one mean proportional number C; these numbers A, B, are like plane numbers.*

a Take D and E the least in the proportion of A *a* 35. 7. to C, or C to B. then D measures A equally as E does C, viz. by the same number F; *b* also D equally *b* 21. 7. measures C, as E does B, viz. by the same number G. *c* Therefore DF = A, and EG = B. *d* and con- *c* 9. 4x. 7. sequently A and B are plane numbers. But because *d* 16.def.7. EF = Cc = DG, *e* shall D. E :: F. G. and alter- *e* 19. 7. nately

f 21. def. 7. nately $D.F :: E.G.$ f Therefore the plane numbers A and B are also like. Which was to be dem.

PROP. XXI.

A, 16. C, 24. D, 36. B, 54. If between two numbers A, B there fall two mean proportional numbers C, D; those numbers A, B are like solid numbers.

a 2. 8. a Take E, F, G, the least $::$ in the proportion
b 10. 8. of A to C. b then E and G are like plane numbers: let the sides of this be H and P, and of that
c 21. def. 7 K and L. c therefore $H.K :: P.L :: d$ E. F. But E,
d cor. 18. 8. F, G, do e equally measure A, C, D, viz. by the
e 21. 7. same number M. and likewise the said numbers E, F, G, do equally measure the numbers C, B, D, viz. by the same number N. f Therefore $A = EM$
f 9. ax. 7. $= HPM$, f and $B = GN = KLN$; g and so A and
g 17. def. 7. B are solid numbers. But for that C $f = FM$, and
h 17. 7. D $f = FN$, therefore shall M. N $h :: FM. FN ::$
k 7. 5. $:: C.D :: E.F :: H.K :: P.L.$ m wherefore A and
l constr. B are like solid numbers. Which was to be dem.
m 21. def. 7. Lemma.

AE, BF, CG, DH, If proportional numbers A, B, C, D, measure proportional numbers AE, BF, CG, DH, by the numbers E, F, G, H, these numbers (E, F, G, H,) shall be proportional.

For being AEDH $a = BFCG$, a and $AD = BC$,
a 19. 7. $\frac{AEDH}{AD} = \frac{BFCG}{BC}$ c that is, $EH = FG$.
b 1. ax. 7. b thence will
c 9. ax. 7. a Therefore $E.F :: G.H.$ Which was to be dem.

Coroll.

d 15. def. 7. Hence $\frac{Bq}{Aq} = \frac{B}{A} \times \frac{B}{A}$ d For 1. B :: B. Bq d and 1. A ::

e lem. prec. A. Aq. e theref. $\frac{1}{A} B :: \frac{B}{A} \cdot \frac{Bq}{Aq}$ d theref. $\frac{Bq}{Aq} = \frac{B}{A} \times \frac{B}{A}$

In like manner $\frac{B}{Ac} \times \frac{B}{Ac} = \frac{Bc}{Acc}$, and so of the rest.

PROP.

PROP. XXII.

Aq, B, C. If three numbers Aq, B, C be
4. 8. 16. continually proportional, and the
first Aq a square, the third C
shall also be a square.

For because Aq C a = Bq, b thence is $C = \frac{Bq}{Aq}$ a 20. 7.
b ax. 7.
 $c = Q \cdot \frac{B}{A}$; But it is plain that $\frac{B}{A}$ is a number, d be- c cor. of the
cause $\frac{Bq}{Aq}$ or C is a number. Theref. if three, &c. d hyp. and
14. 8.

PROP. XXIII.

Ac, B, C, D. If four numbers Ac, B, C, D,
8, 12, 18, 27. be continually proportional, and
the first of them Ac a cube, the
fourth also D shall be a cube.

For because Ac D a = BC, b therefore $D = \frac{BC}{Ac}$ a 19. 7.
b 7. ax. 7.
 $c = \frac{B}{Ac} \times C$; that is (because Ac C = d Bq, and c cor. of the
prec. lem.
b thence $C = \frac{Bq}{Ac}$) $D = \frac{B}{Ac} \times \frac{Bq}{Ac} = \frac{Bc}{Acc}$ c = $C \times \frac{B}{Ac}$ d 20. 7.
But it is evident c that $\frac{B}{Ac}$ is a number, because $\frac{Bc}{Acc}$ e 15. 8.
or D is supposed a number. Therefore if four
numbers, &c.

PROP. XXIV.

A, 16. 24. B, 36. If two numbers A, B, be in the
C, 4. 6. D, 9. same proportion one to another,
that a square number C is to a
square number D, and the first A be a square number,
the second also B shall be a square number.

Between C and D being square numbers, * and * 8. 8.
so between A and B having the same proportion, a 11. 8.
a falls one mean proportional. Therefore b being b hyp.
A

c 22. 8. A is a square number, c B also shall be a square number. Which was to be dem.

Coroll.

1. Hence, If there be two like numbers AB, CD ($A.B :: C.D$) and the first AB be a square, the second also CD shall be a square.

*12. & 18
8.

* For $AB. CD :: Aq. Cq.$

2. From hence it appears, that the proportion of any square number to any other not square, cannot possibly be declared in two square numbers. Whence it cannot be $Q. Q :: 1. 2.$ nor $1. 5 :: Q. Q$ &c.

PROP. XXV.

C, 64. 96. 144. D, 216. If two numbers A, B, be in the same proportion one to another, that a cube number C is to a cube number D, the first of them A being a cube number; the second B shall likewise be a cube number.

a 12. 8.
b 8. 8.
c hyp.
d 23. 8.

a Between the cube numbers C and D, b and so between A and B having the same proportion, fall two mean proportionals; therefore c because A is a cube, d shall B be a cube also. Which was to be dem.

Coroll.

1. Hence, If there be two numbers ABC, DEF ($A. B :: D. E$, and $B. C :: E. F$;) and the first ABC be a cube, the second DEF shall be a cube also.

*12. & 19,
8.

* For $ABC. DEF :: Ac. Dc.$

1. It is perspicuous from hence, that the proportion of any cube number to any other number not a cube cannot be found in two cube numbers.

PROP.

P R O P. XXVI.

A, 20. C, 30. B, 45. *Like plane numbers A, B,*
D, 4. E, 6. F, 9. *are in the same proportion one*
to another, that a square num-
ber is in to a square number.

Between A and B a falls one mean propor- a 18. 8.
tional number C; b take three numbers D, E, F, b 2. 8.
the least :: in the proportion of A to C. the
extremes D, F, b shall be square numbers. But
of equality A. B c :: D. F. therefore A. B :: Q. Q. c 14. 7.
Which was to be dem.

P R O P. XXVII.

A, 16. C, 24. D, 36. B, 54. *Like solid numbers*
E, 8. F, 12. G, 18. H, 27. *A, B, are in the same*
proportion one to ano-
ther, that a cube number is in to a cube number.

a Between A and B fall two mean proportio- a 19. 8.
nal numbers, namely C and D; b take four b 2. 8.
numbers E, F, G, H the least :: in the propor-
tion of A to C; b the extremes E, H, are cube
numbers. But A. B c :: E. H :: C. C. *Which* c 14. 7.
was to be dem.

Schol.

1. From hence is inferred, that no numbers in See *Clas-*
proportion superparticular, superbipartient, or *vius*.
double, or any other manifold proportion not
denominated from a square number, are like
plane numbers.

2. Likewise, that neither any two prime
numbers, nor any two numbers prime one to
another, not being squares, can be like plane
numbers.

The End of the eighth Book.

M

THE

THE NINTH BOOK OF EUCLIDE'S ELEMENTS.

PROP. I.

A, 6. B, 54.

Aq, 36. 108. AB, 324.

I If two like plane numbers *A*, *B* multiplying one another, produce a number *AB*, the number produced *AB* shall be a square number.

a 17. 7.

b 18. 8.

c 8. 8.

d 22. 8.

For $A : B :: Aq : AB$; wherefore since one mean proportional *b* falls between *A* and *B*, *c* likewise one mean proportional number shall fall between *Aq* and *AB*: therefore being the first *Aq* is a square number, *d* the third *AB* shall be a square number too. Which was to be demonstrated.

Or thus. Let *ab*, *cd* be like plane numbers; namely $a : b :: c : d$. & therefore $ad = bc$. and so likewise $ab : cd$, or $ad : bc = ad : ad = Q : ad$.

PROP. II.

A, 6. B, 54.

Aq, 36. AB, 324.

If two numbers *A*, *B*, multiplying one another, produce a square number *AB*, those numbers *A*, *B* are like plane numbers.

a 17. 7.

b 11. 8.

c 8. 8.

d 20. 8.

For $A : B :: Aq : AB$; wherefore being between *Aq*, *AB*, *b* there falls one mean proportional number, *c* likewise one mean shall fall between *A* and *B*. *d* therefore *A* and *B* are like planes. Which was to be dem.

PROP.

PROP. III.

A, 2. Ac, 8. Acc, 64. If a cube number *Ac* multiplying it self produce a number *Acc*. the number produced *Acc* shall be a cube number.

For 1. $A a :: A. Aq b :: Aq. Ac$. therefore between *A* and *Ac* fall two mean proportionals. But 1. $Ac a :: Ac. Acc. c$ therefore between *Ac* and *Acc*, fall also two mean proportionals: and so by consequence seeing *Ac* is a cube, *d Acc* shall be a cube also. Which was to be dem.

Or thus; $aaa(Ac)$ multiplied into it self makes $aaaaaa(Acc)$; this is a cube, whose side is *aa*.

PROP. IV.

Ac, 8. Bc, 27. If a cube number *Ac* multiplying a cube number *Bc*, produce a number *AcBc*, the produced number *AcBc* shall be a cube.

For $Ac. Bc a :: Acc. AcBc$. But between *Ac* and *Bc* two mean proportional numbers fall; c therefore there fall as many between *Acc* and *AcBc*. So that whereas *Acc* is a cube number, *d AcBc* shall be such also. Which was to be dem.

Or thus. $AcBc = aaabbb(ababab) = C: ab$.

PROP. V.

Ac, 8. B, 27. If a cube number *Ac* multiplying a number *B* produce a cube number *AcB*, the number multiplied *B* shall also be a cube.

For $Acc. AcB. a :: Ac. B$. But between *Acc* and *AcB* two mean proportionals; c therefore also as many shall fall between *Ac* and *B*. whence *Ac* being a cube number, *d B* shall be a cube number too. Which was to be dem.

M 2

PROP.

PROP. VI.

A, 8. Aq, 64. Ac, 512. If a number A multiplying it self produce a cube Aq , that number A it self is a cube.

a hyp. For because Aq is a cube, and AqA (Ac) b 19. def. 7. also a cube; therefore c shall A be a cube. Which c 5. 9. was to be dem.

PROP. VII.

A, 6. B, 11. AB, 66. If a composed number A multiplying any number B , produce a number AB , the number produced AB shall be a solid number.

Being A is a composed number, a some other a 13. def. 7. number D measures it, conceive by E . b 9. def. 7. fore $A = DE : c$ whence $DEB = AB$ is a solid c 17. def. 7. number. Which was to be dem.

PROP. VIII.

1. a, 3. a^2 , 9. a^3 , 27. a^4 , 81. a^5 , 243. a^6 , 729.

If from a unit there be numbers continually proportional how many soever (1. a, a^2 , a^3 , a^4 , &c.) the third number from a unit a^2 is a square number; and so are all forward, leaving one between (a^4 , a^6 , a^8 , &c.) But the fourth a^3 is a cube number; and so are all forward, leaving two between (a^6 , a^9 , &c.) The seventh also a^6 , is both a cube number and a square; and so are all forward, leaving five between (a^{12} , a^{18} , &c.)

For 1. $a^2 = Q. a$. and $a^4 = aaaa = Q. aa$. and $a^6 = aaaaaa = Q. aaa$, &c.

2. $a^3 = aaa = C. a$. and $a^6 = aaaaaa = C. aa$ and $aaaaaaaaa = C. aaa$, &c.

3. $a^6 = aaaaaa = C. aa = Q. aaa$. therefore, &c.

a hyp. Or according to Euclide; Because 1. $a :: a. a^2$. b 20. 7. b shall $a^2 = Q. a$. therefore seeing a^2 , a^3 , a^4 , are $::$, c 12. 8. c the third a^4 shall be a square number; and so likewise a^6 , a^8 , &c. Also because 1. $a :: a^3. a^4$. d 23. 8. therefore shall $a^3 b = a^2 \times a = C. a$. d therefore the fourth from a^3 , namely a^6 , shall likewise be a cube, &c. and consequently a^6 is both a cube and a square number, &c.

PROP.

PROP. IX.

1. a. 4. a^2 , 16. a^3 , 64. a^4 , 256, &c. If from a unit there be numbers how many soever continually proportional (1. a, a^2 , a^3 , &c.) and the number following the unit (a) be a square; then all the rest, a^2 , a^3 , a^4 , &c. shall be squares too. But if the number next the unit (a) be a cube, then all the following numbers a^2 , a^3 , a^4 , &c. shall be cube numbers.

1. Hyp. For a^2 , a^4 , a^6 , &c. are square numbers by the prec. Prop. also being a is taken to be a square, a therefore the third a^3 shall be a square, a 22. 8. and likewise a^5 , a^7 , &c. and so all.

2. Hyp. a is taken to be a cube, b therefore a^4 , b 23. 8. a^7 , a^{10} are cubes: but by the prec. a^3 , a^6 , a^9 , c 20. 7. &c. are cubes: lastly, because 1. a :: a. aa. c there d 3. 9. fore shall $a^2 = Q$: a. but a cube multiplied into c 23. 8. it self d produces a cube; therefore a^2 is a cube, e and consequently the fourth from it a^5 , and in like manner a^8 , a^{11} , &c. are cubes. therefore all. Which was to be dem.

Peradventure more clearly thus. let b be the side of the square number a, and so the series a, a^2 , a^3 , a^4 , &c. will be otherwise expressed, thus, bb, b^4 , b^6 , b^8 , &c. It is evident that all these numbers are squares, and may be thus expressed, Q: b, Q: bb: Q. bbb, Q: bbbb, &c.

In like manner, if b be the side of the cube a, the series may be expressed thus, b^3 , b^6 , b^9 , b^{12} . &c. or C: b, C: b^3 , C: b^6 , C: b^9 , &c.

PROP. X.

1. a, a^2 , a^3 , a^4 , a^5 , a^6 , If from a unit there be numbers how many soever continually proportional (1. a, a^2 , a^3 , &c.) and the number next the unit (a) be not a square number; then is none of the rest following a square number, excepting a^2 the third from the unit, and so all forward, leaving one be-

tween ($a^4, a^5, a^6, &c.$) But if that (a) which is next after the unit, be not a cube number, neither is any other of the following numbers a cube, saving a ; the fourth from the unit, and so of all forward, leaving two between, $a^6, a^9, a^{12}, &c.$

a hyp. 1. *Hyp.* For if it be possible, let a^5 be a square number; therefore because $a. a^2 :: a^4, a^5$, and by inversion $a^5. a^4 :: a^2. a$; and also a^5 and a^4 are square numbers, and the first a^2 a square, therefore a shall be likewise a square; contrary to the Hypothesis.

d 24. 7. 2. *Hyp.* If it may be, let a^4 be a cube; being d of equality $a^4. a^6 :: a. a^5$ and inversely $a^6. a^4 :: a^3. a$; and also being a^6 and a^4 are cubes, and the first a^3 a cube; therefore a shall be a cube also; against the Hyp.

PROP. XI.

1. $a, a^2, a^3, a^4, a^5, a^6.$ If there be numbers 1, 3, 9, 27, 81, 243, 729. how many soever in continual proportion from a unit (1, $a, a^2, a^3, &c.$) the less measures the greater by some one of them that are amongst the proportional numbers.

a 5. ax. 7. Because 1. $a :: a. aa.$ therefore $\frac{aa}{a} = a = \frac{aaa}{aa}$

& 20. def. 7. *b 14. 7.* Also because 1. $aa b :: a. aaa.$ therefore $\frac{aaa}{a} = aa =$

$\frac{a^4}{aa} = \frac{a^5}{a^3}, &c.$ Lastly because 1. $a^3 b :: a. a^4.$ therefore

$\frac{a^4}{a} = a^3 = \frac{a^6}{a^2}, &c.$

Coroll.

Hence, If a number that measures any one of proportional numbers, be not one of the said numbers, neither shall the number by which it measures the said proportional numbers, be one of them.

PROP.

PROP. XII.

If there be numbers how many soever in continual proportion from a unit (1, 2, a2, a3, a4.) whatsoever prime numbers B measure the last a4, the same (B) shall also measure the number (a) which follows next after the unit.

If you say B does not measure a, a then B is a prime to a; b and also B is prime to a2; c and b so consequently to a4, which is supposed to measure. Which is absurd.

Coroll.

1. Therefore every prime number that measures the last, does also measure all those other numbers that precede the last.

2. If any number not measuring that next to the unit, does yet measure the last, it is a composed number.

3. If the number next to the unit be a prime, no other prime number shall measure the last.

PROP. XIII.

If from a unit be numbers in continual proportion how many soever (a, a2, a3, &c.) and that after the unit (a) a prime; then shall no other measure the greatest number, but those which are amongst the said proportional numbers.

If it be possible, let some other E measure a4, viz. by F. a then F shall be some other beside a, a2, a3. But because E measuring a4, does not measure a, b therefore E shall be a composed number, c and some prime number measure it, d which does consequently measure a4. e and so is no other than a, therefore a measures E. So also may F be shewn to be a composed number, measuring a4, 9.

M 4

and

measuring $a4$, and so that a measures F . Therefore
 f 9. ax. 7. fore seeing $EF \div f = a4 = a \times a3$, g shall a . $E :: F$.
 g 19. 7. $a3$. Consequently, whereas a measures E , b like-
 h 20. def. 7. wise F shall equally measure $a3$, viz. by the
 k cor. 11. 9. same number G . k Nor shall G be a . or $a2$. there-
fore, as before, G is a composed number, and a
measures it. Wherefore being that $FG \div f = a3$
 $= a2 \times a$ g shall a . $F :: G$ $a2$. and so because A
measures F , b G shall equally measure $a2$. viz.
by the same number H , k which is not a . There-
fore being $GH \div a2 = a2$, l thence H . $a :: a$.
 i 23. def. 7. G . and because a measures G (as before) m H
 n 20. def. 7. also shall measure a , which is a prime num-
ber. Which is impossible.

P R O P. XIV.

$A, 30.$ If certain prime numbers $B, C,$
 $B, 2. C, 3. D, 5.$ D , do measure the least number
 $E - - F - -$ A , no other prime number E shall
measure the same, besides those
that measured it at first.

a 9. ax. 9. If it is possible, let $\frac{A}{E}$ be $= F$. a then $A = EF$.

b 32. 7. b therefore every of the prime numbers $B, C, D,$
measures one of those E, F , not E , which is
taken to be a prime; therefore F which is less
than it self A ; contrary to the Hyp.

P R O P. XV.

$A, 9. B, 12. C, 16.$ If three numbers continually
 $D, 3. E, 4.$ proportional A, B, C , be the
least of all that have the same
proportion with them; any two of them added to-
gether shall be prime to the third.

a 35. 7. a Take D and E the least in the proportion of
 b 2. 8. A to B ; b then $A = Dq$, & b $C = Eq$, b and $B = DE$.
But

But because Dc is prime to E , d therefore shall c 2. 47.
 $D + E$ be prime to both D and E . * therefore d 30. 7.
 $D \times D + E e = Dq + DE$ ($f A + B$) is prime to * 26. 7.
 E , and so to C or Eq . Which was to be dem. e 3. 2.

g In like manner $DE + Eq$ ($B + C$) is prime f before
to D , and consequently to $A = Dq$. Which was g 27. 7.
to be dem. h 26. 7.

Lastly, because Bb is prime to $D + E$, it shall k 4. 2.
also be prime to the square of it $k Dq + 2 DE$ l 30. 7.
 $+ Eq$ ($A + 2 B + C$); l wherefore the said B
shall be prime to $A + B + C$, l and so likewise
to $A + C$. Which was to be dem.

PROP. XVI.

$A, 3. B, 5. C---$ If two numbers A, B , be
prime to one another, the se-
cond B shall not be to any other C , as the first A is
to the second B .

If you affirm $A. B :: B. C$. then whereas A
and Ba are the least in their proportion, Aba 23. 7.
shall measure B as many times as B does C ; b 21. 7.
but A measures itself also; therefore A and Bc 6. ax. 7.
are not prime to one another. against the Hyp.

PROP. XVII.

$A, 8. B, 12. C, 18. D, 27. E---$ If there be
numbers how
many soever in
continual proportion A, B, C, D , and the extremes of
them A, D be prime one to another, the last D shall
not be to any other number E , as the first A is to the
second B .

Suppose $A. B :: D. E$. then alternately A .
 $D :: B. E$. therefore seeing A and B are the
least in their proportion, Aba shall measure a 23. 7.
 B, c and B likewise C , and C the follow- b 21. 7.
ing number D, d and so A shall measure c 20. def. 7.
the said number D . Wherefore A and Dd 11. ax. 7.
are

are not prime to one another, contrary to the Hypothesis.

P R O P. XVIII.

A, 4. B, 6. C, 9.
Bq, 36.

No numbers being given A, B, to consider if there may be a third number found proportional to them C.

a 9. ax. 7. If A measure Bq by any number C, a then
b 20. 7. $AC = Bq$. from whence it is manifest that A.
B :: B. C. Which was to be dem.

A, 6. B, 4. Bq, 16. But if A do not measure Bq, there will not be any third proportional. For suppose A. B :: B. C. a then

c 7. ax. 7, $AC = Bq$. c and consequently $\frac{Bq}{A} = C$. namely A measures Bq. Which is against the Hypoth.

P R O P. XIX.

A, 8. B, 12. C, 18. D, 27.
BC, 216.

Three numbers being given A, B, C, to consider if a fourth proportional to them D may be found.

a 9. ax. 7. If A measures BC by any number D, a then
b ax. 19. 7, $AD = BC$; b therefore it appears that A. B :: C. D. which was required.

But if A do not measure BC, then there can no fourth proportional be found; which may be shewn as in the preceding Prop.

P R O P. XX.

A, 2. B, 3. C, 5.
D, 30. G ----

More prime numbers may be given than any multitude whatsoever of prime numbers A, B, C, propounded.

a 38. 7. a Let D be the least which A, B, C, measure; If
b 33. 7. $D + 1$ be a prime, the case is plain; if composed, b then some prime number, conceive G, measures

ures $D + 1$; which is none of the three A, B, C ; For if it be, seeing it c measures the whole c *suppos.*
 $D - 1$, d and the part taken away D , e it shall d *constr.*
 also measure the remaining unit. Which is absurd. e 12. ax. 7.
 Therefore the propounded number of prime num-
 bers is increased by $D + 1$; or at least by G .

PROP. XXI.

5 5 3 3 2 2
 $A \dots E \dots B \dots F \dots C \dots G \dots D$ 20.

If even numbers, how many soever, AB, BC, CD ,
 be added together, the whole AD shall be even.

a Take $EB = \frac{1}{2} AB$, and $FC = \frac{1}{2} BC$, and
 $GD = \frac{1}{2} CD$. b it is plain that $EB + FC + GD = \frac{1}{2} AD$. c therefore AD is an even num-
 ber. Which was to be dem. a 6. def. 7. b 12. 7. c 6. def. 7.

PROP. XXII.

$A \dots F. B \dots G. C \dots H. D \dots L. E$ 22
 9 7 5 3

If odd numbers, how many soever, AB, BC, CD ,
 DE , be added together, and their multitudes even,
 the whole also AE shall be even,

A unit being taken from each odd number,
 there will a remain AF, BG, CH, DL , even
 numbers, b and thence the number compounded a 7. def. 7.
 of them will be even, add to them the c even b 21. 9.
 number made of the remaining units, and the c hyp.
 whole AE will thereby be even. Which was d 21. 9.
 to be dem.

PROP.

PROP. XXIII.

$\overset{7}{A} \dots \overset{5}{B} \dots \overset{1}{C} \dots \overset{1}{E} \dots \overset{3}{D}$ 15. If odd numbers, how many soever, AB , BC , CD , be added together, and the multitude of them be odd, the whole AD shall be odd.

- For CD one of the odd numbers being taken away, the number compounded of the others AC
 a 22. 9. a is even. Whereto add $CD - 1$, b the whole
 b 21. 2. AE is also even; wherefore the unit being re-
 c 7. def. 7. stored, the whole AD c will be odd. Which was to be dem.

PROP. XXIV.

$\overset{4}{A} \dots \overset{5}{B} \dots \overset{1}{D} \dots \overset{6}{C}$ 10. If an even number AB be taken away from an even number AC , that which remains BC shall be even.

- a 7. def. 7. For if BD ($BC - 1$) be odd, a BC ($BD + 1$) will be even. Which was to be dem. But if you
 b hyp. say BD is even, because AB b is even, c thence
 c 21. 9. AD will be so; a and consequently AC ($AD + 1$) will be odd, contrary to the Hypoth. therefore BC is even. Which was to be dem.

PROP. XXV.

$\overset{6}{A} \dots \overset{1}{D} \dots \overset{3}{C} \dots \overset{7}{B}$ 10. If from an even number AB , an odd number AC be taken away, the remaining number CB shall be odd.

- a 7. def. 7. For $AC - 1$ (AD) a is even. b therefore DB
 b 24. 9. is even; c and consequently CB ($DB - 1$) is
 c 7. def. 7. odd. Which was to be dem.

PROP. XXVI.

$\overset{4}{A} \dots \overset{6}{C} \dots \overset{1}{D} \dots \overset{11}{B}$ 11. If from an odd number AB be taken away an odd number CB , that which remains AC shall be even.

For

For $AB - 1$ (AD) and $CB - 1$ (CD) a are a 7. def. 7.
even; b therefore AD - CD (AC) is even. b 24. 9.
Which was to be done.

P R O P. XXVII.

$A, 4$ 6 If from an odd number
 $A, D, \dots C, \dots B$ II. AB be taken away an even
 5 number CB , the residue AC
shall be odd.

For $AB - 1$ (DB) a is even, and CB is sup- a 7. def. 7.
posed to be even; b therefore the residue CD is b 24. 9.
even: c therefore $CD - 1$ (CA) is odd. *Which* c 7. def. 7
was to be dem.

P R O P. XXVIII.

$A, 3$ If an odd number A multiplying an
 $B, 4$ even number B produce a number AB , the
 $AB, 12$ number produced AB shall be even.

For AB a is compounded of the odd a hyp. and
number A taken as many times as a unit is 15. def. 7.
contained in B an even number. b Therefore AB b 21. 9.
is an even number.

Schol.

In like manner, if A be an even number, AB
shall be an even number also.

P R O P. XXIX.

$A, 3$ If an odd number A multiplying an
 $B, 5$ odd number B , produce a number AB , the
 $AB, 15$ number produced AB shall be odd.

For AB a is compounded of the odd a 15. def. 7.
number B taken as often as a unit is included in
 A likewise an odd number. b Therefore AB is b 23. 9.
an odd number. *Which was to be dem.*

Schol.

Schol.

$\frac{B, 12}{A, 3}$ (C, 4. 1. An odd number A measuring an even number B , measures the same by an even number C .

a 9. ax. 7. For if C be affirmed to be odd, then because
b 29. 9. $B = AC$. b therefore B shall be odd, against the Hypothesis.

$\frac{B, 15}{A, 3}$ (C, 5. 2. An odd number A measuring an odd number B , measures the same by an odd number C .

a 28. 9. For if C be said to be even, a then AC , or B will be even, contrary to the Hypothesis.

$\frac{B, 16}{A, 3}$ (C, 5. 3. Every number (A and C) that measures an odd number B , is itself an odd number.

For if either A or C be affirmed to be even, B
a 28. 9. a shall be an even number, against the Hypoth.

P R O P. XXX.

$\frac{B, 24}{A, 3}$ (C, 8. $\frac{D, 12}{A, 3}$ (E, 4.

If an odd number A measure an even number B , it shall also measure the half of it D .

a hyp. a Let $\frac{B}{A}$ be $= C$. b then C is an even number.

b 1. Schol.

29. 9. Therefore let E be $= \frac{1}{2} C$, then $Bc = CA d = 2$

c 9. ax. 7. $EA e = 2 D$. f therefore $EA = D$; g and conse-

d 1. 2. quently $\frac{D}{A} = E$. Which was to be dem.

e hyp.

f 7. ax. 1.

P R O P. XXXI.

g 7. ax. 7. $A, 5. B, 8. C, 16. D---$ If an odd number A be prime to any number B , it shall also be prime to the double thereof C .

a 3. schol. If it be possible, let some number D measure A
29. 9. and C , a then D measuring the odd number A shall be odd it self, b and so shall measure B the half of the even number C . theref: A and B are not prime one to another. Which is against the Hyp.
b 30. 9. Coroll.

Coroll.

It follows from hence that an odd number which is prime to any number of double progression, is also prime to all the numbers of that progression.

P R O P. XXXII.

1. A, 2. B, 4. C, 8. D, 16. *All numbers A, B, C, D, &c. in double progression from the binarie are evenly even only.*

It is evident that all these numbers 1, A, B, C, D, a are even, and $b \div 2$, namely in a double proportion, c and so every less measures the greater by some one of them. d Wherefore all are evenly even. But for that A is a prime number, e no number besides these shall measure any of them. Therefore they are evenly even only. *Which was to be dem.*

P R O P. XXXIII.

A, 30. B, 15. *If of a number A, the half B be odd, the same A is evenly odd only.*

Being an odd number B a measures A by two an even number, b therefore B is evenly odd. If you affirm it to be evenly even, c then some even number D measures it by an even number E. whence $2 B d = A, d = DE$. e wherefore $2 E :: D. B$. and therefore as 2 f measures the even number E, g so D an even number measures B an odd. *Which is impossible.*

P R O P. XXXIV.

A, 24. *If an even number A be neither doubled from two, nor have its half part odd, it is both evenly even and evenly odd.*

It is undoubtable, that A is evenly even, because the half of it is not odd. But because, if A be divided into two equal parts, and its half again into

7. def. 7. into two equal parts and so on, we shall at length light upon some *a* odd number (not upon the number two, because *A* is not supposed to be doubled upward from two) which shall measure *b* 1 sch. 29. *A* by an even number, so *b* otherwise *A* it self should be odd, against the Hypoth. Therefore *A* also evenly odd. Which was to be dem.

P R O P. XXXV.

A.....8.
 4 8
 B....F.....G 12.
 C.....18.
 9 6 4 8
 D.....H.....L....K.....N 27.

If there be numbers in continual proportion how many soever *A*, *B*, *G*, *C*, *D*, *N*, and the number *FG* be taken from the second, and *KN* from the last, equal to the first *A*; as the excess of the second *BF* is to the first *A*, so shall the excess of the last *DK* be to all the numbers that precede it, *A*, *B*, *G*, *C*.

- a* hyp. From *DN* take *NL* = *BG*, and *NH* = *C*.
b 17. 5. Because *DN*. *C* (*HN*) *a* :: *HN*. *BG* (*LN*) *a* ::
c 12. 5. *LN* (*BG*.) *A* (*KN*.) *b* therefore by dividing
d 3. ax. 1. each, shall *DH*. *HN* :: *HL*. *LN* :: *LK*. *KN*. *c*
 wherefore *DK* *C* + *BG* + *A* :: *LK* (*d* *BF*.) *KN*
 (*A*.) Which was to be dem.

Coroll.

- e* 18. 5. Hence *e* by compounding. *DN* + *BG* + *C*.
A + *BG* + *C* :: *BG*. *A*.

P R O P. XXXVI.

1. *A*, 2. *B*, 4. *C*, 8. *D*, 16.
E, 31. *G*, 62. *H*, 124. *L*, 248. *F*, 496.
 M, 31. *N*, 465.
 P ---- *Q* ----

If from a unit be taken how many numbers soever *1*, *A*, *B*, *C*, *D*, in double proportion continually, until the whole added together *E* be a prime number; and if this whole

whole E multiplying the last produce a number F , that which is produced F shall be a perfect number.

Take as many numbers E, G, H, L , likewise in double proportion continually; then a of e a 14. 7.
equality $A. D :: E. L.$ b therefore $AL = DE$ $c = b$ 19. 7.

$F. d$ whence $L = \frac{F}{2}$. Wherefore E, G, H, L, F , c hyp. d 7. ax. 7.

are $::$ in double proportion. Let $G - E$ be $= M$, and $F - E = N$; e then $M. E :: N. E + G$ $+ e$ 35. 9.

$H + L.$ f But $M = E.$ g therefore $N = E + f$ 3. ax.

$G + H + L.$ b therefore $F = 1 + B + C + D$ g 14. 5.

$+ E + G + H + L = E + N.$ Moreover be h 2. ax. 1.

cause D k measures DE (F) l therefore every k 7. ax. 7.

one, $1, A, B, C, m$ measuring D , as m also $E,$ l 11. ax. 7.

G, H, L , does measure F . And further, no other m 11. 9.

number measures the said F . For if there do, let n 9. ax. 7.

it be P , which measures F by $Q.$ n therefore PQ o 19. 7.

$= F = D. F.$ o therefore $E. Q :: P. D.$ therefore p 13. 9.

seeing A a prime number measures D , p and so q 20. def. 7.

no other P measures the same, q consequently

E does not measure $Q.$ Wherefore E being sup- r 31. 7.

posed a prime number, r it shall be prime to $Q.$ s 23. 7.

f wherefore E and Q are the least in their pro- t 21. 7.

portion; t and so E measures P as many times u 13. 7.

as Q does D ; u therefore Q is one of them $A,$

$B, C.$ Let it be B , seeing then of equality $B.$

$D :: E. H :: x$ and so $BH = DE = F = PQ.$ x 19. 7.

and so also $Q. B :: H. P.$ y therefore $H = P.$ y 14. 5.

therefore P is also one of them $A, B, C,$ &c.

against the Hypothesis. Wherefore no other be- z 22. def. 7.

sides the foresaid numbers measures F , and z con-

sequently F is a perfect number. Which was to

be demonstrated.

The End of the ninth Book.

N

THE

THE TENTH BOOK OF EUCLIDE'S ELEMENTS

Definitions.

- I. **C**ommenfurable magnitudes are those, which are measured by one and the same measure.

The note of commensurability is $\sqcap$, as


 $A \sqcap B$; that is, the line A of 8 foot is
commensurable to the line B of 13 foot;
because D a line of one foot measures both
 A and B . Also $\sqrt{18} \sqcap \sqrt{50}$; because
 $\sqrt{2}$ measures both $\sqrt{18}$ and $\sqrt{50}$. For $\sqrt{\frac{18}{2}} = \sqrt{9} = 3$. and $\sqrt{\frac{50}{2}} = \sqrt{25} = 5$.
wherefore $\sqrt{18} \sqcap \sqrt{50} :: 3. 5$.

II. Incommensurable magnitudes are such, of which no common measure can be found.

Incommensurability is denoted by this mark $\sqn$; as $\sqrt{6} \sqn \sqrt{15}$ (5;) that is, $\sqrt{6}$ is incommensurable to the number 5, or to a magnitude designed by that number; because there is no common measure of them, as shall appear hereafter.

III. Right lines are commensurable in power, when the same space does measure their squares.

The



The mark of this commensurability is Γ ; as $AB \Gamma CD$. i. e. the line AB of 6 foot is in power commensurable to the line CD , which is expressed by $\sqrt{20}$. because E the space of one foot square does as well measure ABq (36) as the rectangle XY (20) to which the square of the line CD ($\sqrt{20}$) is equal. The same note Γ sometimes signifies commensurable in power only.

IV. Lines incommensurable in power are such; to whose squares no space can be found to be a common measure.

This incommensurability is denoted thus; $\varsigma \Gamma$; $\sqrt{8}$ i. e. the numbers or lines 5, and $\sqrt{8}$ are incommensurable in power, because their squares 25 and 8 are incommensurable.

V. From which it is manifest, that to any right line given right lines infinite in multitude are both commensurable and incommensurable; some in length and power, others in power only. The right line given is called a Rational line.

The note of which is ρ .

VI. And lines commensurable to this line, whether in length and power, or in power only, are also called Rational, ρ .

VII. But such as are incommensurable to it, are called Irrational.

And denoted thus ρ .


VIII. Also the square which is made of the said given right line is called Rational, ρv .

IX. And likewise such figures as are commensurable to it, are Rational, $\rho \alpha$.

X. But such as are incommensurable, Irrational, $\rho \alpha$.

XI. And those right lines also, which contain them in power, are Irrational ρ .

Schol.



acor. 15. 4.
b 47. 1.

That the last seven definitions may be rendered more clear by an example, let there be a circle $ADBP$, whose semidiameter is CB , inscribe therein the sides of the ordinate figures, as of a Hexagone BP , of a triangle AP , of a square BD , of a Pentagone FD . Therefore, if according to the 5. def. the semidiameter CB be the Rational line given, expressed by the number 2. to which the other lines BP , AP , BD , FD , are to be compared, then $BP a = BC = 2$, wherefore BP is $\dot{p} \sqcup BC$, according to the 6. def. Also $AP b = \sqrt{12}$ (for $ABq (16) - BPq (4) = 12$) therefore $AB \dot{p} \sqcup BC$. likewise according to the 6. def. and $APq (12)$ is $\dot{p} v$ by the 9. def. Moreover $BD b = \sqrt{DCq + BCq} = \sqrt{8}$; whence BD is $\dot{p} \sqcup BC$; and $BDq \dot{p} v$. Lastly, $FDq = 10 - \sqrt{10}$. (as shall appear by the praxis to be delivered at the 10. 13.) shall be $\dot{p} v$, according to the 10. def. and consequently $FD = \sqrt{10} - \sqrt{20}$ is $\dot{p}$, according to the 11. def.

A Postulate.

That any magnitude may be so often multiplied, till it exceed any magnitude whatsoever of the same kind.

Axiomes.

1. A magnitude measuring how many magnitudes soever, does also measure that which is composed of them.

2. A

2. A magnitude measuring any magnitude whatsoever, does likewise measure every magnitude which that measures.

3. A magnitude measuring a whole magnitude and a part of it taken away, does also measure the residue.

PROP. I.

Two unequal magnitudes AB, C, being given, if from the greater AB there be taken away more than half (AH) and from the residue (HB) be again taken away more than half (HI) and this be done continually, there shall at-length be left a certain magnitude IB, less than the less of the magnitudes first given C.

a Take C so often, till its multiple a *post.* 10. **ACD** DE do somewhat exceed AB, and there be $DE = FG = GE = C$. Take from AB more than half HA, and from the remainder HB more than half HI, and so continually, till the parts AH, HI, IB, be equal in multitude to the parts DF, FG, GE. Now it is plain, that FE, which is not less than $\frac{1}{2}$ DE, is greater than HB, which is less than $\frac{1}{2}$ AB $\supset$ DE. And in like manner GE, which is not less than $\frac{1}{2}$ FE, is greater than IB $\supset$ $\frac{1}{2}$ HB. therefore C, or GE $\supset$ IB. Which was to be dem.

The same may also be demonstrated, if from AB the half AH be taken away, and again from the residue HB the half HI, and so forward.

P R O P. II.

Two unequal magnitudes being given (AB, CD) if the less AB, be continually taken from the greater CD, by an interchangeable subtraction, and the residue do not measure the magnitude going before; then are the magnitudes given incommensurable.

If it be possible, let some magnitude E be the common measure. Then because AB taken from CD, as often as it can be, leaves a magnitude FD less than it self, AC E and FD taken from AB leaves GB, and so forward; *a* therefore at length some magnitude GB—E shall be left, therefore E *b* measuring AB, *c* and so CF, *b* and the whole CD, *d* shall also measure the residue FD. *c* consequently also AG; *d* 3. ax. 10. *d* wherefore it shall likewise measure the remainder GB, less than it self. Which is absurd.

P R O P. III.

Two commensurable magnitudes being given AB, CD, to find out their greatest common measure FB.

Take AB from CD, and the residue ED from AB, and FB from ED, till FB measure ED (which will come to pass at length, *a* because by the Hyp. AB $\sqsubset$ CD) FB shall be the magnitude required.

For FB *b* measures ED, *c* and so also AF; but it measures it self too, *d* therefore likewise AB, *c* and consequently CE, *d* and so the whole CD. Wherefore FB is the common measure of AB, CD. If you affirm G to be a common measure greater than that, *e* 2. ax. 10. then G measuring AB and CD, *e* measures also *f* 3. ax. 10. CE and *f* the remainder ED, *e* and so AF; and *f* consequently the remainder FB, the greater the less. Which is absurd.

Coroll.

Hence
nitudes,
measure

A—
B—
C—
Three
C, to f
a Fi
of any
measu
tude f
a F
measu
magn
c ther
great
great


He
meas
sure

A—
C—
B—
to a
a
of A
ofte
the
D.
qu

Coroll.

Hence, A magnitude that measures two magnitudes, does also measure their greatest common measure.

PROP. IV.



Three commensurable magnitudes being given A, B, C, to find out their greatest common measure.


a Find out D the greatest common measure *a* 3. 10. of any two A, B; *a* also E the greatest common measure of D and C. therefore E is the magnitude sought for.

a For it is clear, E measuring D and C, *b* does *b* constr. & measure the three A, B, C. Conceive another 2. ax. 10. magnitude F greater than that to measure them; *c* then F measures D, *c* and consequently E the *c* cor. 3. 10. greatest common measure of D, and C, the greater the less. Which is absurd.

Coroll.

Hence also it appears, that if a magnitude measure three magnitudes, it shall likewise measure their greatest common measure.

PROP. V.



a C being found the greatest common measure *a* 3. 10. of A, B; as often as C is contained in A and B, so often is 1 contained in the numbers D and E; *b* b 17. def. 7. therefore C. A :: 1. D; wherefore inversely A. C :: D. 1. *b* but likewise C. B :: 1. E. *c* therefore of equality A. B :: D. E :: N. N. W. W. to be dem.

PROP. VI.

E ————— $F, 1.$ If two magnitudes
 A ————— $C, 4.$ $A, B,$ have such pro-
 B ————— $D, 3.$ portion one to another
 as number C has to

number D , those magnitudes A, B shall be commensurable.
 a sch. 10. 6. What part 1 is of the number C , a that let E
 b constr. be of A . Therefore because $E.A :: 1.C$. and $A.B$
 c hyp. $c :: C.D$. d therefore of equality shall $E.B :: 1.D$.
 d 21. 5. Wherefore seeing 1 e measures the number D , f
 e 5. ax. 7. likewise E measures B ; but it g also measures A .
 f 20. def. 7. b therefore $A \sqsubset B$. Which was to be dem.
 g constr.
 h 1. def. 10.

PROP. VII.

A ————— Incommensurable magni-
 B ————— tudes $A, B,$ have not that
 proportion one to another,
 which number has to number.

a 6. 10. If you affirm $A.B :: N.N.$ a then $A \sqsubset B$
 against the Hypothesis.

PROP. VIII.

A ————— If two magnitudes $A, B,$ have
 B ————— not that proportion one to ano-
 ther, which number has to num-
 ber, those magnitudes are incommensurable.

a 5. 10. Conceive $A \sqsubset B$ a then $A.B :: N.N.$ con-
 trary to the Hypothesis.

PROP. IX.

A ————— The squares described of right
 B ————— lines commensurable in length,
 $E, 4.$ have that proportion one to ano-
 $F, 3.$ ther, that a square number has
 to a square number. And squares, which have that pro-
 portion

portion on
 a square
 surable i
 right line
 proportion
 has to a
 not such
 ber has t
 commensf

1. Hy
 For a
 Aq A
 Bq B
 fore A
 fore A

2. Hy
 A
 B
 twice
 fore A
 Which

3. H
 For s
 is shev

4. H
 B, Fo
 as abo

Li
 trary
 7

portion one to another, that a square number has to a square number, shall also have their sides commensurable in length. But such squares as are made of right lines incommensurable in length, have not that proportion one to another, which a square number has to a square number. And squares which have not such proportion one to another as a square number has to a square number, have not their sides commensurable in length.

1. Hyp. A. $\square$ B. I say Aq. Bq :: Q. Q.

For a let A.B :: number E. number F. therefore

$\frac{Aq}{Bq} (b \frac{A}{B} \text{ twice}) c = \frac{E}{F} \text{ twice. } d = \frac{Eq}{Fq} \text{ e there-}$ a 5. 10.
fore Aq. Bq :: Eq. Fq :: Q. Q. Which was to be dem. b 20. 6.
c sch. 23. 5.

2. Hyp. Aq. Bq :: Eq. Fq :: Q. Q. I say A $\square$ B. For

$\frac{A}{B} \text{ twice } (f \frac{Aq}{Bq}) g = \frac{Eq}{Fq} h = \frac{E}{F} \text{ twice. i there-}$ d 11. 8.
fore A. B :: E. F :: N. N. k wherefore A $\square$ B. e 11. 5.
f 20. 6.
g hyp.
h 11. 8.
i sch. 23. 5.
k 6. 10.

3. Hyp. A $\square$ C. I deny that Aq. Bq :: Q. Q. For suppose Aq. Bq :: Q. Q. then A $\square$ B, as is shewn before, against the Hyp.

4. Hyp. Not Aq. Bq :: Q. Q. I say that A $\square$ B, For conceive A $\square$ B. then Aq. Bq :: Q. Q. as above, against the Hyp.

Coroll.

Lines $\square$ are also $\square$. but not on the contrary. And lines $\square$ are not therefore $\square$. but $\square$ are also $\square$.

PROP.

P R O P. X.

If four magnitudes be proportional ($C. A :: B. D$) and the first C be commensurable to the second A , the third B shall be commensurable to the fourth D . And if the first C be incommensurable to the second A , also the third B shall be incommensurable to the fourth D .

a 5. 10.
b 6. 10.
c 7. 10.
d 8. 10.

If $C \sqsubset A$, a then $C. A :: N. N$ b :: $B. D. b$ therefore $B \sqsubset D$. But if $\sqsubset A$, c then shall not $C. A :: N. N$:: $B. D. d$ wherefore $B \sqsubset D$. Which was to be dem.

Lemma 1.

To find out two plane numbers, not having the proportion which a square number hath to a square.

Any two plane numbers not like will satisfy this Lemma, as those numbers which have superparticular, superbipartient, or double proportion; or any two prime numbers. See sch. 27. 8.

Lemma 2.

B, 5. K — 1 — 1 — 1 — M
C, 3. H — 1 — 1 — 1 — R

To find out a line HR , to which a right line given KM hath the proportion of two numbers given B, C .
a sch. 10. 6. a Divide KM into as many equal parts as there are units in the number B . and let as many of these, as there are units in the number C , b make the right-line HR , it is manifest that $KM. HR :: B. C$.

Lemma 3.

To find out a line D , to the square of which, the square of a right line given KM hath the proportion of two numbers given B, C .

Allow $B. C a :: KM. HR$. and between KM and HR , b find a mean proportional D . Therefore $KMq. Dq c :: KM. HR d :: B. C$.

P R O P.

PROP. XI.

A ——— B, 20. To find two right lines
E ——— C, 16. incommensurable to a right
D ——— line given A, one D in
length only, the other E in
power also.

1. Take the numbers B, C, a so that there a 1.lem.10.
be not B.C :: Q. Q. b and let B.C :: Aq. Dq. c 10.
it is plain that A $\square$ D. But Aq d $\square$ Dq. b 3.lem.10.
Which was to be done. 10.

2. d Make A.E :: E.D. I say Aq $\square$ Eq. For A. c 9. 10.
D :: Aq. Eq. therefore since A $\square$ D, as before : f d 6. 10.
therefore Aq $\square$ Eq. Which was to be done. d 13. 6.
e 20. 6.
f 10. 10.

PROP. XII.

Magnitudes (A, B) commensurable to the
same magnitude C, are also commensurable
one to the other.

Because A $\square$ C, and C $\square$ B, a let A. a 5. 10.
D, 18. E, 8. C :: N. N :: D. E, and C.
F, 2. G, 3. B :: N. N :: F. G. b take b 4. 8.
H, 5. I, 4. K, 6. three numbers H, I, K, the
A B Cleaft :: in the proportions of D to E,
and F to G. Now because A.C c :: D.E c :: H.I, c constr.
and C. B c :: F.G :: I.K. d therefore of equality d 22. 5.
A.B :: H.K :: N. N. e therefore A $\square$ B. Which e 6. 10.
was to be dem.

Schol.

Hence, Every right line commensurable to a
rational line is also it self rational. And all 12. 10. and
right lines rational are commensurable to one def. 6.
another, at least in power. Also, every space
commensurable to a rational space is rational
too : and all rational spaces are commensurable def. 9.
one to another. But magnitudes whereof one
is rational, the other irrational, are incommen- def. 7. &
surable amongst themselves. 10.

PROP.

PROP. XIII.

A —————
 C —————
 B —————

If there be two magnitudes A , B , and one of them A commensurable to a third C , but the other B incommensurable, those magnitudes A , B are incommensurable.

a hyp. Conceive $B \sqsubset A$. then being C a $\sqsupset A$, b
b 12. 10. therefore $C \sqsupset B$, against the Hyp.

PROP. XIV.

A —————
 B —————
 C —————
 D —————

If there be two magnitudes commensurable A , B ; and one of them A incommensurable to any other magnitude C , the other also B shall be incommensurable to the same C .

Imagine $B \sqsubset C$. then for that A a $\sqsupset B$, b therefore $A \sqsupset C$, against the Hyp.

a hyp.
b 12. 10.

$A B C$

PROP. XV.

A —————
 B —————
 C —————
 D —————

If four right lines be proportional ($A. B :: C. D$.) and the first A be in power more than the second B by the square of a right line commensurable to it self in length, then also the third C shall be more in power than the fourth D by the square of a right line commensurable to it self in length. But if the first A be more in power than the second B by the square of a right line incommensurable to it self in length, then shall the third C be more in power than the fourth D by the square of a right line incommensurable to it self in length.

a hyp. For because $A. B a :: C. D$. b therefore $Aq. Bq ::$
b 22. 6. $Cq. Dq.$ c therefore by division $Aq - Bq. Bq :: Cq -$
c 17. 5. $Dq. Dq.$ d wherefore $\sqrt{Aq - Bq. Bq} :: \sqrt{Cq -$
d 22. 6. $Dq. Dq}$ and so inverſely $B. \sqrt{Aq - Bq} :: D. \sqrt{Cq -$
e cor. 4. 5. $Dq. f$ therefore of equality $A. \sqrt{Aq - Bq} :: C.$
f 22. 5. $\sqrt{Cq - Dq}$. consequently if $A \sqsupset$, or $\sqsupset \sqrt{Aq - Bq}$,

-Bq. g t
Dq. Wh

A —

D

commens
the whole
of the pa
at first

1. Hy

AB, BC

AC

2. H

AC, A

(BC) a

to be de

Hen

posed

them,

other

D

surab

the n

of th

shall

1.


mea

- A

agai

-Bq. g then likewise $C \sqsupseteq$, or $\sqsupseteq \sqrt{} : Cq - g$ 10. 10.
Dq. Which was to be dem.

PROP. XVI.



If two magnitudes commensurable AB, BC , be composed, the whole magnitude AC shall be

commensurable to each of the parts AB, BC . And if the whole magnitude AC be commensurable to either of the parts AB , or BC , those two magnitudes given at first AB, BC , shall be commensurable.


1. Hyp. a Let D be the common measure of a 3. 10.
 AB, BC ; b so D measures AC . and therefore b 1. ax. 10.
 $AC \sqsupseteq AB$, and BC . Which was to be dem. c 1. def. 10.

2. Hyp. a Let D be the common measure of d 3. ax. 10.
 AC, AB . d therefore D measures $AC - AB$
(BC) and consequently $AB \sqsupseteq BC$. Which was
to be dem.

Coroll.

Hence it follows, if a whole magnitude composed of two, be commensurable to any one of them, the same shall be commensurable to the other also.

PROP. XVII.



If two incommensurable magnitudes AB, BC , be composed, the whole magnitude also AC shall be incommen-

surable to either of the two parts AB, BC . And if the whole magnitude AC be incommensurable to one of them AB , the magnitudes first given AB, BC , shall be incommensurable.

1. Hyp. If it can be, let D be the common measure of AC, AB . a therefore D measures AC a 3. ax. 10.
 $- AB$ (BC) b and therefore a No $AB \sqsupseteq BC$, b 1. def. 10.
against the Hypoth.


2. Hyp.

c 16. 10. 2. Hyp. Conceive $AB \perp BC$. c therefore $AC \perp AB$, against the Hyp.

Coroll.

Hence also, If one magnitude, composed of two, be incommensurable to any one of them, the same also shall be incommensurable to the other.

PROP. XVIII.




If there be two unequal right lines AB , GK and upon the greater AB a parallelogram ADB equal to the fourth part of a square made of the less line GK , and wanting in

figure by a square, be applied, and divide the said AB into parts commensurable in length AD , DB ; then shall the greater line AB be more in power than the less GK by the square of a right line FD commensurable in length to the greater. And if the greater AB be in power more than the less GK by the square of the right line FD commensurable unto it self in length, and a parallelogram ADB equal to the fourth part of the square made of the less line GK , and wanting in figure by a square, be applied to the greater AB , then shall it divide the same into parts

- a 10. 1. AD , DB commensurable in length.
 b 28. 6. a Divide GK equally in H , and b make the
 c 8. 2. rectangle $ADB - GHq$. Cut off $AF = DB$. then is
 d constr. $\&$ $ABq c = 4 ADB d (4 GHq \text{ or } GKq) + FD$. Now
 4. 2. in the first place, if $AD \perp DB$. then shall AB e
 e 16. 10. $\perp BD$ c $\perp 2 DB$ f ($AF = DB$, or $AB - FD$) b
 g cor. 16. therefore $AB \perp FD$. Which was to be dem. But
 10. secondly, if $AB \perp FD$, b then shall $AB \perp AB$
 k 12. 10. $- FD (2 DB)$ k therefore $AB \perp DB$. l where-
 l 16. 10. fore $AD \perp DB$. Which was to be dem.

PROP.

PROP. XIX.



If there be two right lines unequal AB , GK , and to the greater AB be applied a parallelogram ADB equal to the fourth part of a square made upon the less GK , and


wanting in figure by a square, and also thus applied divide the said AB into parts AD , DB incommensurable in length; the greater line AB shall be in power more than the less GK by the square of the right line FD incommensurable to the greater in length. And if the greater line AB be more in power than the less GK by the square of a right line FD incommensurable unto it self in length, and if also upon the greater AB be applied a parallelogram ADB equal to the fourth part of the square of the less GK and wanting in figure by a square, then shall it divide the said greater line AB into parts incommensurable in length AD , DB .

Suppose all the same that was done and said in the prec. Prop. Therefore first, If $AD \perp DB$, *a* then shall $AB \perp DB$. *b* Wherefore $AB \perp 2 DB$ ($AB - FD$) *c* therefore $AB \perp FD$. *b* 17. 10. *c* 13. 10. *c* cor. 17. Which was to be dem.

Secondly, If $AB \perp FD$, then $AB \perp AB - FD$ ($2 DB$;) *d* wherefore $AB \perp DB$, *e* and consequently $AD \perp DB$. Which was to be dem. *d* 13. 10. *e* 17. 10.

PROP.

PROP. XX.




A rectangle BD comprehended under right lines BC, CD, rational and commensurable in length, according to one of the foresaid ways, is rational.

- Let A be given $\hat{p}$, and
 a 46. 1. the square BE described upon BC. Because DC,
 b 1. 6. CE (BC) $b :: BD$. BE. and DC $c \sqsubset BC$,
 c hyp. therefore shall the rectangle BD be $\sqsubset$ square
 d 10. 10. BE. wherefore seeing the square BE $e \sqsubset Aq$,
 e hyp. and shall also BD be $\sqsubset Aq$. and so the rectangle
 g def. 10. BD is $\hat{p}$. Which was to be dem.
 f 12. 12. Note, There are three kinds of lines rational

commensurable one to another. For either, of two lines rational commensurable in length one to the other, one is equal to the rational line propounded, or neither of them is equal to it, notwithstanding both of them are commensurable to it in length; or lastly, both of them are commensurable to the rational line given only in power. And these are the ways which the present Theorem speaks of.

In numbers, let there be BC $\sqrt{8}$ ($2\sqrt{2}$) and CD $\sqrt{18}$ ($3\sqrt{2}$) then shall the rectangle BD = $\sqrt{144} = 12$.

PROP. XXI.



- If a rational rectangle DB be applied to a rational line DC, it makes the breadth thereof CB rational, and commensurable in length to that
 a 1. 6. line DC, whereto DB is applied.
 b hyp. Let G be propounded $\hat{p}$, and the square DA described on BC. because BD. DA $:: a BC$. CA;
 c sch. 12. and BD, DA b are $\hat{p} a c$ and so $\sqsubset$. d therefore
 d 10. 10. BC $\sqsubset CA$. but CD (CA) is $\hat{p}$. e therefore BC
 e sch. 12. is $\hat{p}$. Which was to be dem.
 f 10.


In numbers, let there be rectangle DB. 12, and DC, $\sqrt{8}$. then shall CB, $\sqrt{18}$. but $\sqrt{18} = 3\sqrt{2}$. and $\sqrt{8} = 2 \times \sqrt{2}$.

Lemma.

A ——— To find out two right lines rational commensurable only in power.

C ——— Let A be propounded ρ . a a 11. 10.
Take B $\perp$ A, a and C $\perp$ B. b it is clear that b sch. 12.
B and C are the lines required. 10.

P R O P. XXII.



A rectangle DB, comprehended under right lines rational DC, CB, commensurable in power only, is irrational; and the right line H, which containeth that rectangle in power is irrational, and called a Medial line.

Let G be the propounded ρ , and the square DA described on DC, and let Hq = DB. Because AC. CB a :: DA. DB. b and AC $\perp$ CB, c shall a i. 6. be DA $\perp$ DB (Hq,) d but Gq $\perp$ DA. e b hyp. therefore Hq $\perp$ Gq. f wherefore H is ρ . Which c 10. 10. was to be dem. and let it be called a Medial line, d hyp. and because AC. H :: H. CB. 93. def. 10.

In numbers, let there be DC, 3. and CB, $\sqrt{6}$. e 13. 10. then shall the rectangle be DB (Hq) $\sqrt{54}$. f 11. 10. wherefore H is $\sqrt{54}$.


The note of a medial line is μ , of a medial rectangle $\mu\rho$, of more together $\mu\alpha$.

Schol.

Every rectangle that can be contained under two right lines rational commensurable only in power, is medial, altho' it be contained under two right lines irrational: and every medial rectangle may be contained under two right lines rational, commensurable only in power, as for example, the

the $\sqrt{24}$ is $\mu\nu$, because it is contained under $\sqrt{3}$, and $\sqrt{8}$, which are ρ , $\square$. altho' it may be contained under $v\sqrt{6}$, and $v\sqrt{96}$ irrationals; for $\sqrt{24} = v\sqrt{576} = v\sqrt{6 \times v\sqrt{96}}$.

P R O P. XXIII.




If the rectangle BD made of a medial line A, be applied on a rational line BC, it makes the breadth CD rational, and incommensurable in length to the line

BC, whereunto the rectangle BD is applied.

- a sch. 12. Because A is μ , a therefore shall Aq be equal to some rectangle (EG) contained under EF and 10.
 b 1. ax. 1. FG ρ $\square$. b therefore $BD = EG$. c whence BC. c 14. 6. $EF :: FG$. CD. d therefore $BCq \cdot EFq :: FGq \cdot CDq$.
 d 22. 6. But BCq and EFq e are $\rho\alpha$, f and so $\square$. g therefore $FGq \square CDq$. Wherefore being FG is ρ , f sch. 12. h therefore CD shall be ρ . Moreover, because 10.
 g 10. 10. EF. FG k :: $EFq \cdot EG$ (BD;) for that EF $\square$ FG, h sch. 12. therefore the rectangle BD $\square$ CDq. Whence 10.
 k 1. 6. being CDq. $BD o :: CD$. BC. p shall CD be $\square$ BC. therefore, &c.

- l 10. 10.
 m sch. 12.
 10.
 n 13. 10.
 o 1. 6.
 p 10. 10.

P R O P. XXIV.



A right line B commensurable to a medial line A is also a medial line.

- a 11. 6.
 b hyp.
 c 23. 10. rectangle $CF = Bq$. Because Aq (CE) is $\mu\nu$, b and d 1. 6. CD ρ . e therefore shall the latitude DE be ρ $\square$ CD. But for that CE. CF d :: ED. DF. and CE $\square$ CF,

CF, *f* therefore ED $\perp$ DF. *g* therefore DF is *f* 10. 10.
 $\perp$ CD. *h* whence the rectangle CF (Bq) is *g* 12. and
 μ , and so B is μ . Which was to be dem. 13. 10.

Obf. That the note $\perp$ for the most part signifies *h* 22. 10.
 commensurable in power only, as in this and the pre-
 cedent demonstrations, &c.

Coroll.

Hereby it is manifest that a space commensu-
 rable to a medial space, is also medial.

Lemma.


A ——— To find out two right lines medial
 B ——— A, B, commensurable in length, and
 C ——— also two, A, C, commensurable on-
 ly in power.

a Let A be any μ , *b* take B $\perp$ A, and *c* C a lem. 22.
 $\perp$ A. *d* it appears to be done. 10. and
 13. 6.

P R O P. XXV.

h 2 lem. 10.

10.




A rectangle DB contained un *c* 3 lem. 10.
 der DC, CB medial right lines 10.
 commensurable in length, is me- dconstr. &
 dial. 24. 10.

Upon DC describe the square
 DA. Being AC (DC.) CB *a* :: *a* 1. 6.
 DA. DB. and DC $\perp$ CB; *b* shall DA $\perp$ DB. *b* 10. 10.
c therefore DB is μ . Which was to be dem. *c* 24. 10.

O 2

PROP.

PROP. XXVI.



A rectangle AC comprehended under medial right lines AB, BC commensurable only in power, is either rational or medial.


246. 1. Upon the lines AB, BC, *a* describe the squares AD, CE; and upon FG *p* make the rectangles FH, b cor. 16.6. = AD, *b* and IK = AC. *b* and LM = CE.

The squares AD, CE, that is, the rectangles c hyp. & FH, LM, *c* are $\mu\alpha$ and $\square$. therefore GH, KM, 24. 10. having the same proportion *d* are ρ , *e* and $\square$. d 23. 10. f therefore GH $\times$ KM is $\rho\nu$. But because AD, e 10. 10. AC, CE. that is, FH, IK, LM, *g* are $\div\div$; *b* and f 20. 10. so GH, HK. KM also $\div\div$; *k* thence HK $\times$ GH g sch. 22.6. $\times$ KM. *l* therefore HK is ρ or $\square$, or $\square$ IH h 1. 6. (GF;) if $\square$, *m* then the rectangle IK or AC is k 17.6. $\rho\nu$. but if $\square$, *n* then AC is $\mu\alpha$. Which was to l 12. 10. be dem.

m 20. 10.

n 22. 10.

Lemma.




If A and E be $\square$ only, Then first shall Aq,

a hyp. and Eq, Aq + Eq, Aq - Eq a $\square$. And secondly Aq, Eq, 10. 10. Aq + Eq. Aq - Eq $\square$ AE and 2 E. For A.Eb:: b 1. 6. Aq. AE *b* :: AE. Eq. therefore seeing A c $\square$ E, c hyp. d shall Aq $\square$ AE, *e* and 2 AE. also Eq d $\square$ d 10. 10. AE, *e* and 2 AE. wherefore because Aq + Eq $\square$ e 14. 10. Aq and Eq; and Aq - Eq $\square$ Aq and Eq. f f 14. 10. therefore shall Aq + Eq. f and Aq - Eq be $\square$ AE, and 2 AE.

Hence

Hence also thirdly, $Aq \cdot Eq \cdot Aq + Eq \cdot Aq - Eq$,
 $2AE \cdot g \cdot Aq + Eq \cdot 2AE$; and $Aq + Eq \cdot g \cdot 14.16.$
 $-2AE \cdot g$ and $Aq + Eq + 2AE \cdot Aq + Eq \cdot 17.10.$
 $-2AE \cdot b(Q.A - E.)$ hcor.7.12.


PROP. XXVII.



A medial rectangle
 AB exceedeth not a me-
 dial rectangle AC by a
 rational rectangle DB.

Upon EF ρ , a make $a \cdot cor.16.6.$
 $EG = AB$, a and EH
 $= AC$. The rectangles
 AB, AC , i. e. EG, EH , b b hyp.
 are ua ; c therefore $FG \cdot c \cdot 23.10.$
 and FH are $\rho \cdot EF$. Whence, if KG , d i. e. $DB \cdot d \cdot 3.ax.1.$
 be $\rho \cdot e$ then shall HG be $\sqcap HK$; f wherefore e $21.10.$
 $HG \cdot FH$. g and consequently $FG \cdot FH \cdot q \cdot 13.10.$
 But FH is ρ . b therefore is $FG \cdot \rho$. but FG was $g \cdot lem.16.$
 before. Which is contradictory. $10.$
 $b \cdot sch.12.$
 $10.$

Schol.



1. A rational rectangle AE
 exceeds a rational rectangle AD
 by a rational rectangle CE.

For AE a $\sqcap AD$. b there-a hyp.
 fore AE $\sqcap CE$. c wherefore b $cor.16.$
 CE is $\rho \cdot v$. Which, &c. $10.$

2. A rational rectangle AD c $sch.12.$
 joynd with a rational rectangle $10.$
 CF makes a rational rectangle
 AF.

For AD a $\sqcap AF$. b where-a $sch.11.$
 fore AF $\sqcap AD$ and CF, c $10.$
 and so AF is $\rho \cdot v$. Which was to be dem. $b \cdot 16.11.$
 $c \cdot sch.12.$


O;

PROP. 10.

Schol.

A, 6. C, 12. To find out plane numbers, like
 B, 4. D, 8. or unlike.
 AB, 24. CD, 96. Take any four numbers
 A, 6. C, 5. proportional A. B :: C. D. it
 B, 4. D, 8. is manifest that AB and CD
 AB, 24. CD, 40. are like plane numbers. And
 you may find out as many un-
 like plane numbers, as you
 please, by help of *Schol.* 27. 8.

Lemma.



To find out two square numbers (*DEq* and *CDq*)
 so that the number composed of them (*CEq*) be
 square also.

Take AD, DB like plane numbers (of which
 let both be equal, or both odd) viz. AD, 24.
 and DB, 6. The total of these (AB) is 30; the
 difference (FD) 18 half of which (CD) is 9.
 Now the like plane numbers AD, DB, have a 18. 8.
 one mean number proportional, namely DE.
 therefore it is evident that every of those num-
 bers CE, CD, DE, are rational, and by conse-
 quence *CEq* (*b CDq* + *DEq*) is the square b 47. 1.
 number required.

Whereby it will be easy to find out two square
 numbers, the excess of which is a square or not
 a square number, namely by the same constru-
 ction c shall *CEq* — *CDq* be = *DEq*.

c 3. ax. 1.

But if AD, DB be plane numbers unlike, the
 medial

medial proportional line (DE) shall not be a rational number, and so neither shall the excess (DEq) of the square numbers, CEq, CDq, be a square number.

Lemma 2.

2. To find out two such square numbers B, C, as the number compounded of them D is not square. Also to divide a square number A into two numbers B, C, not squares.

A, 3. B, 9. C, 36. D, 45.

1. Take any square number B, and let C be $= 4 B$, and $D = B + C$. I say the matter is done.

For B is Q. by the constr. likewise because
 a 24. 8. B.C :: 1. 4 :: Q. Q. a therefore C also shall be a square number. But because $B + C$. (D) C ::
 b cor. 24. 8. 5. 4 :: not Q. Q. b therefore shall not D be a square number. Which was to be done.

A, 36. B, 24. C, 12. D, 3. E, 2. F, 1.

2. Let A be some square number. Take D, E, F, plane numbers alike, and let D be $= E + F$. make D.E :: A. B. and D.F :: A. C. I say the thing required is done.

For because D.E + F :: A. B + C. and D =
 a 14. 5. E + F, a therefore shall $A = B + C$. Now sup-
 b 21. def. 7. pose B to be square, b then A and B, c and con-
 c 26. 8. sequently D and E are like plane numbers.
 Which is contrary to the Hyp.

The same absurdity will follow if C be supposed a square number. Therefore, &c.

PROP.

b 9. 14

For, as above, AB, AF, are $\hat{p}$ $\square$. also ABq. BFq :: CD. ED. therefore being CD is not Q. AB, BF b shall be $\square$. Which was to be done.

In numbers, let there be AB, 5. CD, 45. CE = 36. ED = 9. Make 45. 9 :: 25 (ABq.) 5 (AFq.) therefore AF = $\sqrt{5}$. consequently BFq = 45 - 25 = 20. wherefore BF = $\sqrt{20}$.

P R O P. XXXII.

A ————— To find out two medial
B ————— lines C, D, commensurable
C ————— only in power, comprehending
D ————— a rational rectangle
CD, so that the greater

C be more in power than the lesser D by the square of a right line commensurable in length to the greater.

- a 30. 10. a Take A and B $\hat{p}$ $\square$; so as $\sqrt{Aq} - Bq \square$
b 13. 6. A. b and make A. C :: C. B. c and A. B :: C.D.
c 12. 6. I say the thing is done.
d constr. For because A and d B are $\hat{p}$ $\square$. e therefore
e 22. 10. shall C (f $\sqrt{AB}$) be μ . g and thence also C $\square$ D.
f 17. 6. b therefore D is likewise μ . Furthermore, where-
g 10. 10. as A.B d :: C.D; and inversely A.C :: B.D :: C.B;
h 24. 10. and Bq is $\hat{p}$. therefore shall CD (k Bq) be $\hat{p}$.
k 17. 6. Lastly, because $\sqrt{Aq} - Bqd \square$ A, l shall $\sqrt{Cq}$
l 15. 10. - Dq be $\square$ C. therefore, &c. But if $\sqrt{Aq} - Bq$
" $\square$ Aq, then shall $\sqrt{Cq} - Dq$ be $\square$ C.

In numbers; let there be A 8, B $\sqrt{48}$ ($\sqrt{64 - 16}$) therefore C = $\sqrt{AB} = v \sqrt{3072}$. and D = $v \sqrt{1728}$. wherefore CD = $v \sqrt{5308416} = \sqrt{2304}$.

P R O P. XXXIII.

A ————— To find out two medial
D ————— lines D, E, commensurable
B ————— in power only, comprehending
C ————— a medial rectangle DE,
E ————— so far that the greater D
shall be more in power than

the less E, by the square of a right line commensurable to the greater in length.

a Take


a Take A and C ρ $\perp$, so that $\sqrt{Aq - Cq} \sqsupseteq a$ 30. 10.
A. b take also B $\perp$ A and C, and make A. D $\perp$ *lem. 21.*
c :: D. B d :: C. E. then D and E are the lines *10.*
sought for. c 13. 6.

For because A and C e are ρ , e and B $\perp$ A d 12. 6.
and C, f therefore shall B be ρ , and D ($\sqrt{AB}$) g *constr.*
shall be μ . But because A. D :: C. E, therefore in f *sch. 12.*
versely A. C :: D. E. wherefore seeing A $\perp$ C, *10.*
therefore D shall be $\perp$ E. therefore E is μ . *g 22. 10.*
Furthermore, l being D. B :: C. E. and BC is μ . *h 10. 10.*
also DE, equal to it, is μ . Lastly, because A. *k 24. 10.*
C :: D. E. e seeing $\sqrt{Aq - Cq} \sqsupseteq A$. therefore l 22. 10.
 $\sqrt{Dq - Eq} \sqsupseteq D$. therefore, &c. But if $\sqrt{Aq} m$ 16. 6.
 $- Cq \sqsupseteq A$. then $\sqrt{Dq - Eq} \sqsupseteq Eq$. n 15. 5.

In numbers, let there be A 8, C $\sqrt{48}$. B $\sqrt{8}$.
then D $\sqrt{3072}$. and E $\sqrt{588}$. wherefore
D. E :: 2. $\sqrt{3}$. and DE = $\sqrt{1344}$.

PROP. XXXIV.

To find out two right
lines AF, BF, incommen-
surable in power, whose
squares added together
make a rational figure,
and the rectangle contain-
ed under them medial. a 31. 10.
b 10. 1.



a Let there be found AB, CD, ρ $\perp$; so that $\sqrt{ABq - CDq} \sqsupseteq AB$. divide CD equally in *d 12. 6.*
G. c make the rectangle AEB = GCq. Upon *e cor. 8. 6.*
AB the diameter draw the semicircle AFB, e $\perp$ *17. 6.*
rect the perpendicular EF, and draw AF, BF. *f 7. 5.*
these are the lines required. g 19. 10.


For AE. BE d :: BA $\times$ AE. AB $\times$ BE. But BA h 10. 10.
 $\times$ AE e = AFq; and AB $\times$ BE = FBq. *f there- k 31. 3. &*
fore AE. EB :: AFq. FBq. therefore being AE g 47. 1.
 $\perp$ EB, h AFq shall be $\perp$ FBq. Moreover ABq l *constr.*
(k AFq + FBq) l is ρ . Lastly, EFq l = AE l = m 1. ax. 1.
CGq. m therefore EF = CG. therefore CD $\times$ AB n 22. 10.
= 2 EF $\times$ AB. But CD $\times$ AB n is μ . *o there- o 24. 10.*
fore AB $\times$ EF, p or AF $\times$ FB is μ . Which was p *sch. 22. 6.*
to be dem. The

The Explication of the same by numbers.

Let AB be 6. CD $\sqrt{12}$. then $CG = \sqrt{12} = \sqrt{3} \cdot \sqrt{4} = 2\sqrt{3}$. but $AE = 3 + \sqrt{6}$. and $EB = 3 - \sqrt{6}$. whence AF shall be $\sqrt{18 + 216}$. and $FB \sqrt{18 - 216}$. Also $AEq + FBq$ is 36, and $AF \times FB = \sqrt{108}$.

But AE is found in this manner. Because BA (6.) $AF :: AF.AE$. therefore $6 AE = AFq = AEq + 3 (EFq.)$ therefore $6 AE - AEq = 3$. Put $3 + e = AE$. then $18 + 6e - 9 - 6e - ee$, that is, $9 - ee = 3$. or $ee = 6$. wherefore $e = \sqrt{6}$ and so $AE = 3 + \sqrt{6}$.

PROP. XXXV.




To find out two right lines AE, EB, incommensurable in power, whose squares added together make a medial figure, and the rectangle contained under them rational.

a 32.10. a Take AB and CF $\mu \sqsupset$, so that $AB \times CF$ be ρv , and $\sqrt{ABq - CFq} \sqsupset AB$. and let the rest be done as in the prec. Prop. AE, EB are the line required.

For, as it is shewn there, $AEq \sqsupset EBq$ also b constr. $ABq (AEq + EBq)$ is μv , and lastly, $AB \times CF$ c sch. 12. b is ρv , e therefore also $AB \times DE$, that is, $AE \times d sch. 22.6. EB$, is ρv . therefore, &c.

PROP.

PROP. XXXVI.




To find out two right lines BA, AC, incommensurable in power, whose squares added together make a medial figure, and the rectangle

also contained under them medial, and incommensurable to the figure composed of the squares.

a Take BC and EF μ $\square$, so that BC $\times$ EF a 33. 10. be $\mu\nu$. and $\sqrt{BCq - EFq} \square$ BC. and so forward, as in the prec. BA, AC, shall be the lines sought for.

For (as above) BAq $\square$ ACq. also BAq + ACq is $\mu\nu$. and BA $\times$ AC is $\mu\nu$. Lastly, BC b $\square$ EF, and b const. c so BC $\square$ EG; likewise BC.EG d :: BCq. BC $\times$ c 13. 10. EG (BC $\times$ AD, or BA $\times$ AC) e therefore BCq. d 1. 6. (ABq + ACq) $\square$ BA $\times$ AC. therefore, &c. e 14. 10.

Schol.




To find out two medial lines incommensurable both in length and power.

a Take BC μ and let BA $\times$ AC be $\mu\nu$, and $\square$ a 36. 10. BCq (BAq + ACq b make BA. H :: H. AC. then b 33. 6. I say BC and H are μ $\square$. For BC is μ . a and c 17. 6. BA $\times$ AC (c Hq) is $\mu\nu$. wherefore H is also μ . d 14. 10. d Likewise BA $\times$ AC $\square$ BCq; therefore Hq $\square$ BCq. therefore, &c.

More

Here begin the senaries of lines irrational
by composition.

PROP. XXXVII.


 If two rational lines AB , BC , commensurable only in power, be added together, the whole line AC is irrational, and is called a binomial line, or of two names.

a hyp. For because AB a $\square$ BC , thence b shall AC q
b lem. 26. be $\square$ AB q. But AB a is a $\dot{p}$, c therefore AC
10. is $\dot{p}$. Which was to be dem.

c II. def.

10.

PROP. XXXVIII.

 If two medial lines AB , BC , only in power commensurable, be compounded, and contain a rational rectangle, the whole line AC is irrational, and called a first binomial line.

a hyp. For being that AB a $\square$ BC , b shall AC q be
b lem. 26. $\square$ $AB \times BC$, $\dot{p}v$. c therefore AC is $\dot{p}$. Which
10. was to be dem.

c II. def.

10.

Lemma.




A rectangle AC , contained under a rational line AB and an irrational line BC , is irrational.

For if the rectangle AC be affirmed,
a hyp. $\dot{p}v$, a then being AB is $\dot{p}$, b the breadth BC
b 21. 10. shall be also $\dot{p}$. against the Hyp.

PROP.

PROP. XXXIX.




If two medial lines AB, BC , commensurable only in power, containing a medial rectangle, be compounded, the whole line AC shall be irrational, and is called a second binomial line.

Upon the propounded line DE ρ make the rectangle $DF = ACq$; b and $DG = ABq + BCq$. a cor. 10. 6. b 47. 1. 83

Because $ABq \perp BCq$, d therefore $ABq + BCq$ $\perp$ 12. 6. d
 BCq , i. e. DG , $\perp$ ABq : but ABq is $\mu\nu$, e c hyp. e
 therefore DG is $\mu\nu$, but the rectangle ABC is d 16. 10. d
 taken $\mu\nu$; e and consequently $2 ABC$ (f HF) is e 24. 10. e
 $\mu\nu$. g therefore EG and GF are ρ . Being also that f 4. 2. f
 $DG \perp HF$; and $DG. HF :: k EG. GF$; g 27. 10. g
 therefore $EG \perp GF$, m therefore the whole EF h lem. 26. h
 is ρ . n wherefore the rectangle DF is $\rho\nu$, o there- o
 fore $\sqrt{DF}$, i. e. AC , is ρ . Which was to be dem. k 1. 6. k
 l 10. 10. l
 m 37. 10. m
 n lem. 38. n

PROP. XL.



If two right lines AB, BC commensurable only in power, be added together,

making that which is composed of their squares rational, and the rectangle contained under them medial, the whole right line AC is irrational, and is called a Major line.

For whereas $ABq + CBq$ a is $\rho\nu$. and $b \perp 2 b$ a hyp. a
 ABC c $\mu\nu$; and so ACq (d $ABq + BCq + 2 ABC$) b sch. 12. b
 $c \perp ABq + BCq$ $\rho\nu$, f therefore shall AC be ρ . c hyp. and c
 Which was to be dem. d 4. 2. d
 e 17. 10. e
 f 11. def. f
 10 . 10

PROP. f 11. def. 10.

PROP. XLI.



If two right lines AC , CB , incommensurable in power, be added to-

gether, having that which is made of their squares added together medial, and the rectangle contained under them rational, the whole right line AB shall be irrational, and is called *A line containing in power a rational and a medial rectangle.*

a hyp. and For 2 rectangles ACB a $\mu\nu$, $b \sqsupset ACq + CBq$ *sch. 12. 10. c* $\mu\nu$. d therefore 2 ACB $d \sqsupset ABq$. wherefore b *sch. 12. AB* is ρ . Which was to be dem.

10.


c hyp.

d 17. 10.

e 11. def.

10.

PROP. XLII.




If two right lines GH , HK , incommensurable in power be added together, having both that which is composed of their squares medial, and the rectangle contained under them medial, and incommensurable to that which is composed of their squares, the whole right line GK is irrational, and is called *a line containing in power two medial figures.*

a hyp. Upon the propounded line FB ρ make the rect-
b 23. 10. angles $AF = GKq$, and $CF = GHq + HKq$. Being
c 4. 2. $GHq + HKq$ (CF) a is $\mu\nu$, the breadth CB b shall
d 2. 6. be ρ . Also because 2 rectangles GHK (c AD) a
e 20. 10. is $\mu\nu$, therefore AC b shall be ρ . Moreover be-
f 37. 10. cause the rectangle AD $a \sqsupset CF$, d and $AD.CF$
g lem. 38. $AC.CB$. e thence shall e AC be $\sqsupset CB$. f
10. wherefore AB is g ρ . therefore the rectangle AF .
h 21. def. e . GKq is ρ ; b and consequently GK is ρ .
10. Which was to be dem.

PROP.

Because the rectangles ADB, AEB *a* are μa ; *a* 37. 10.
a and each of ADq, DBq, AEq, EBq is $\dot{p} x$. *b* and *b* sch. 27.
 so ADq + DBq *b* and AEq + EBq are also $\dot{p} x$. *b* there- 10.
 fore ADq + DBq = AEq + EBq *i. e.* 2 AEB = 2 *c* sch. 5. 2.
 ADB is $\dot{p} y$. *d* therefore AEB - ADB is $\dot{p} y$. therefore *d* sch. 12.
 μv exceeds μv by $\dot{p} y$. *e* Which is absurd. 10.

PROP. XLIV.




Conceive AB to be divided into other names
 AE, EB, whereupon every one ADq, DBq, EBq, a 38. 10.
 will be a *ma.* and the rectangles ADB, AEB, b *sch.* 27.
 and the doubles of them p 2. b therefore 2 AEB 10.
 - 2 ADB. c i. e. ADq + DBq - : AEq + EBq c *sch.* 5.2.
 is p. Which is absurd. d 27. 10.

P

PROP.

PROP. XLV.




A second bimedral line AB , is divided into its names AC , CB , only at one point C .

Suppose there were other names AD , DB . Upon the propounded line EF $\dot{\rho}$ make the rectangles $EG = ABq$, and

$EH = ACq + CBq$, as also $EK = ADq + DBq$.

- a 39. 10. Because ACq , BCq are $\mu\alpha$ $\square$; b $ACq +$
 b 16. and CBq (EH) shall be $\mu\nu$. c therefore the breadth
 24. 10. FH is $\dot{\rho}$. a moreover the rectangle ACB , d and so
 c 23. 10. $2 ACB$ (e IG) is $\mu\nu$. c therefore HG is also $\dot{\rho}$.
 d 24. 10. And since EH is f $\square$ IG , g and $EH:IG :: FH$.
 e 4. 2. HG . b therefore FH , HG shall be $\square$. k there-
 f lem. 26. fore FG is a binomial, whose names are FH ,
 10. HG . By the same reason FG is binomial, and
 g 1. 6. the names of it FK , KG : contrary to the 43. of
 h 10. 10. this Book.
 k 37. 10.

PROP. XLVI.




A Major line AB is at one point only D divided into its names, AD , DB .

Imagine other names AE , EB . whereupon the

- a 40. 10. rectangles ADB , AEB , are a $\mu\alpha$. a and as well
 b sch. 27. $ADq + DBq$, as $AEq + EBq$ are $\dot{\rho}\alpha$. b therefore
 10. $ADq + DBq = AEq + EBq$, c i. e. $2 AEB =$
 c sch. 5. 2. $2 ADB$ is $\dot{\rho}\nu$. Which is impossible.
 d 27. 10.

PROP.

PROP. XLVII.




A line *AB* containing in power a rational and a medial figure

is divided at one point only *D* into its names *AD*, *DB*.

Conceive other names *AE*, *EB*. then both *AEq* + *EBq*, and *ADq* + *DBq* are $\mu\alpha$. a and the a 41. 10. rectangles *AEB*, *ADB* are $\rho\alpha$. b the ϵ 10. 2 *AEB* b sch. 17. - 2 *ADB*, c i. e. *ADq* + *DBq* = *AEq* + *EBq* 10. is $\rho\mu$. d Which is absurd. c sch. 5. 2. d 27. 10.

PROP. XLVIII.



A line *AB* containing in power two medial rectangles, is at one point only *C* divided into its names *AC*, *CB*.

If you will divide *AB* into other names *AD*, *DB*. draw upon the line propounded *EF* ρ the rectangles *EG* = *ABq*, and *EH* = *ACq* + *CBq*, and *EK* = *ADq* + *DBq*. then because *ACq* + *CBq*, namely, *EH*, a is $\mu\nu$, b the breadth *FH* shall be ρ . Also because 2 *ACB*, c that is, *IG*, is a $\mu\nu$, *HG* b shall be b 23. 10. likewise ρ . Therefore, whereas *EH* a $\sqsupset$ *IG*. and c 4. 2. *FH*. *IG* d :: *FH*. *HG*, thence *FH* e shall be $\sqsupset$ d 1. 6. *HG*. f therefore *FG* is a binomial, and the names e 10. 10. of it *FH*, *HG*. In like manner *FK*, *KG* shall be f 37. 10. the names of it, against the 43. of this Book.

Second Definitions.

A Rational line being propounded, and the binomial divided into its names, the greatest of whose names is more in power than the less by the square of a right line commensurable to the greater in length; then

P 2

1. If

I. If the greater name be commenfurable in length to the rational line propounded, the whole line is called a first binomial line.

II. But if the lesser name be commenfurable in length to the rational line propounded, the whole line is called a second binomial.

III. If neither of the names be commenfurable in length to the rational line propounded, it is called a third binomial.

Furthermore if the greater name be more in power than the less by the square of a right line incommenfurable to the greater in length, then

IV. If the greater name be commenfurable to the propounded rational line in length, it is called a fourth binomial.

V. If the lesser name be so, a fifth.

VI. If neither, a sixth.

P R O P. XLIX.

A....4C.....; B To find out a first binomial
D ———— line, EG.

a sch. 29. E ———— G a Take AB, AC, square
10. numbers, whose excels CB
b 2. lem. H ———— F is not Q. let D be propound-
10. 10. ed p. b Take EF $\sqsupset$ D, and c make AB. CB ::
c 3. lem. EFq, FGq. then EG shall be a 1. bin.
10. 10. For EF d $\sqsupset$ D. e therefore EF is p. falso
d constr. EFq $\sqsupset$ FGq. g therefore FG is also p. likewise
e 6.def.10. d be EFq. FGq :: AB.CB :: Q. not Q. h therefore
f 6. 10. EF $\sqsupset$ FG. Lastly, because by conversion of pro-
g sch. 12. portion, EFq, EFq - FGq :: AB.AC :: Q.Q. thence
10. EF k shall be $\sqsupset$ $\sqrt{EFq - FGq}$. l therefore EG
h 9. 10. is a first binomial. Which was to be done.
k 9. 10. In numbers thus; let there be D 8. EF 6.
l 1.def.48. AB 9. CB 5. wherefore because 9. 5 :: 36. 25,
10. therefore FG is $\sqrt{20}$. and consequently EG is
6 + $\sqrt{20}$.

P R O P.

PROP. L

A...4 C...5 B

D —————

E ————— G

H ————— F

To find out a second binomial line, EG.

Take AB and AC square numbers, the excess of which is CB not Q. Let D

Prove it as the prec.

be the line propounded $\dot{p}$. take FG $\perp$ D, and make CB. AB :: FGq. EFq. then EG will be the line desired.

For FG $\perp$ D. wherefore FG is $\dot{p}$. Also EFq $\perp$ FGq. therefore EF is $\dot{p}$. Likewise because FGq. EFq :: CB. AB :: not Q. Q. thence FG is $\perp$ EF. Lastly, seeing CB. AB :: FGq. EFq. and inversely AB. CB :: EFq. FGq. therefore as in the foregoing Prop. EF $\perp$ $\sqrt{\text{EFq} - \text{FGq}}$. a where- a 2. def. 48. by EG is a 2. binomial. Which was to be done. 10.

In numbers; let there be D 8, FG 10, AB 9, CB 5, then EF is $\sqrt{180}$. wherefore EG is $10 + \sqrt{180}$.

PROP. LI.

A...4 C...5 B

L...6

G —————

D ————— F

H ————— E

To find out a third binomial line, DF.

a Take A B, A C, a sch. 19. square numbers, the 10.

excess of which B C is not Q. and let L be a number not Q next

greater than CB, viz. by a unit or two. Let G be the line propounded $\dot{p}$. b Make L. AB :: Gq, DEq. b and AB. CB :: DEq. EFq. then DF shall b 3. lem. be a 3. bin. 10. 10.

For because DEq $\perp$ Gq, d DE is $\dot{p}$. also Gq. c constr. 6. DEq :: L. AB :: not Q. Q. e therefore G $\perp$ DE. 10.

Likewise being that DEq $\perp$ EFq, d also EF is d sch. 12. $\dot{p}$. Moreover because DEq. EFq :: AB. CB :: Q. not 10.

Q. f is DE $\perp$ EF. and being that by constr. and e 6. 10.

P 3

of

f 9. 10. of equality Gq. EFq :: L. CB :: not Q. Q. (for g
g sch. 27. 8. L and CB are not like plane numbers) h there-
h 9. 10. fore shall G be also $\square$ EF. Lastly, as in the
prec. Prop. $\sqrt{DEq} - EFq \square DE$. k therefore

k 3. def. 48. DF is a 3 bin. Which was to be done.

10.

In numbers; let there be AB, 9. CB, 5. L, 6.
G, 8. then shall be DE $\sqrt{96}$, and EF $\sqrt{48}$.
wherefore DF = $\sqrt{96} + \sqrt{48}$.

P R O P. LII.

A ... 3 C 6 B
G _____
a sch. 29. D _____ F
10. H _____ E
b 2. lem. line propounded p. b take DE $\square$ G. c and make
10. AB. CB :: DEq. EFq. then DF shall be a 4 bin.
c 3. lem. For, as in the 49 of this Book, DF may be
10. 10. shewn to be a binom. and also because by constr.
d 9. 10. and conversion of proportion DEq. DEq - EFq
e 4. def. 48. :: AB. AC :: Q not Q. d shall DE be $\square \sqrt{DEq}$
10. - EFq. e therefore DF is a 4 bin.
In numbers, let G be 8, DE, 6. then EF shall
be $\sqrt{24}$. therefore DF is $6 + \sqrt{24}$.

P R O P. LIII.

A ... 3 C 6 E
G _____
D _____ F
H _____ E
To find out a fifth bino-
mial line, DF.
Take any square num-
ber AB, whole segments
AC, CB are not Q. Let G
be the line propounded p.
take EF $\square$ G. and make CB. AB :: EFq. DEq.
then shall DF be a 5 bin.

For DF shall be a bin. as in the 50. of this Book, and because by construction, and inversion DEq. EFq :: AB. CB. and so by conversion of proportion, DEq. DEq - EFq :: AB. AC :: Q. not Q. *a* therefore shall DE be $\sqrt{\square} \sqrt{\text{DEq}} - a$ 9. 10. EFq. *b* therefore DF is the 5 bin. Which was to *b* 5.def.48. *a* 10. be done.

In numbers, let there be G, 7. EF, 6. then DE shall be $\sqrt{54}$. wherefore DF is $6 + \sqrt{54}$.

P R O P. LIV.


A.....5 C.....7 B To find out a sixth binomial
L.....9 line.
G ———— Take AC, CB, prime num-
D. ———— F bers, so that AC + CB (AB)
H ———— E be not Q. take also any
number square L. Let G be
the line propounded *p*. *a* 3. lem.
and make L. AB :: Gq. DEq. and AB. CB :: 10. 10.
DEq. EFq. then DF shall be a 6 binomial.

For DF may be demonstrated bin. as in the 51. of this Book, and also by reason that DE and EF $\square$ G. lastly, likewise because by constr. and conversion of proportion DEq. DEq - EFq :: AB. AC :: not Q. Q. (For AB is prime to AC, *b* and so unlike to it) *c* therefore DE $\square \sqrt{\square}$ b sch. 27. 8. DEq - EFq. *d* therefore DF is a 6 bin. Which *c* 9. 10. *d* 6.def.48. was required.

In numbers, let there be G 6. DE $\sqrt{48}$. then 10. EF shall be $\sqrt{28}$. wherefore DF is $\sqrt{48} + \sqrt{28}$.

Lemma.

a 28. 6.
b 31. 1.
c 14. 2.



Let AD be a rectangle, and the sides thereof AC divided unequally in E ; also let the lesser portion EC be equally divided in F , upon the line AE make the rectangle $AGE = EFq$. and from the points G, E, F draw GH, EI, FK , parallel to AB . Let the square LM be made equal to the rectangle AH , and upon OM produced the square $MN = GI$, and let the right lines LQ, TS, NRS, NPT be produced.

I say 1. MS, MT , are rectangles. For by reason of the right angles of the squares OMQ, RMP , a shall QMR be a right line. b therefore RMQ, QEP , are right angles. wherefore the parallelograms MS, MT are rectangles.

c 2. ax. 21. 2. Hence it is plain that $LS = LT$, and consequently that LN is a square.

3. The rectangles SM, MT, EK, FD are equal. For because the rectangle $AGE = EFq$. e thence shall $AE.EF :: EF.GE$. f and so $AH.EK :: EK.GI$. that is by const. $LM.EK :: EK.MN$. g sch. 22. 6. g but $LM.SM :: SM.MN$. therefore $EK = SM$. h 9. 5. $k = FD$ i = MT .

k 36. 1. 3. Hence $LN = AD$.

l 43. 1. 5. Being that EC is equally divided in F , it is m 2. ax. 1. plain that EF, FC, EC are $\square$.

n 16. 10. 6. If $AE = EC$, and $AE = \sqrt{AEq - ECq}$, o 18. and then shall AG, GE, AE , be $\square$ also, because 16. 10. $AG.GE :: AH.GI$. p therefore shall AH, GI , p 10. 10. i.e. LM, MN , be $\square$. Likewise thereupon,

7. OM $\square$ MP. For by the Hyp. AE $\square$ EC.
therefore BC $\square$ GE. q wherefore EF $\square$ q 14. 10.
GE. but EF. GE :: EK. GI. r therefore EK $\square$ r 10. 10.
GI, that is, SM $\square$ MN. but SM. MN :: OM.
MP. r therefore OM $\square$ MP.

8. If AE be supposed $\square$ $\sqrt{AEq - ECq}$, it is s 19. and
apparent that AG, GE, AE, are $\square$. whence 17. 10.
LM $\square$ MN. for AG.GE :: AH.GI :: LM.MN.

*These being well considered, we shall easily dispatch
the six following Propositions.*

PROP. LV.

*If a space AD be contained under a rational line
AB, and a first binomial line AC (AE + EC) the
right line OP which containeth that space in power
is irrational, and called a binomial line.*

All that being supposed which is described
and demonstrated in the next foregoing Lemma,
it is manifest that the right line OP containeth
in power the space AD. a Likewise AG, GE, a hyp. and
AE are $\square$. therefore seeing AE is ρ $\square$ AB. lem. 54. 10.
c shall also AG and GE be ρ $\square$ AB. d there- b hyp.
fore the rectangles AH, GI, that is the squares c sch. 12.
LM. MN are ρa . therefore OM, MP are ρe $\square$. 10.
f and consequently OP is a binomial. Which d 20. 10.
was to be dem. e lem. 54.

In numbers, let there be AB 5. AC $4 + \sqrt{12}$. 10.
wherefore the rectangle AD = $20 + \sqrt{300}$ = f 37. 10.
to the square LN. therefore OP is $\sqrt{15 + \sqrt{3}}$
5. namely a 6 binomial.

PROP. LVI.


*If a space AD be comprehended under a rational
line AB, and a second binomial AC (AE + EC) the
right line OP, which contains that space AD in power,
is irrational, and called a first medial line.*

The

The foresaid Lemma of the 54 of this Book be
 a hyp. anding again supposed, then shall OP be $= \sqrt{AD}$
 lem. 54. 10. a also AE, AG, GE are $\square$ therefore since AE
 b hyp. is $\square \sqsubset AB$, likewise AG, GE $\square$ shall be $\square \sqsubset$
 c sch. 12. AB. therefore the rectangle AH, GI, i. e. OM
 10. MPq. d are μa . e Moreover OM $\square$ MP. Lem.
 e lem. 54. ly, EF $\square$ EC, and EC $\square$ AB, g wherefore
 10. EK. i. e. SM, or OMP is μr . h Consequently OP
 f hyp. 12. is a first binomial. Which was to be dem.
 10. In numbers, let there be AB, 5; and AC
 g 20. 10. 48: +6. then the rectangle AD $= \sqrt{1200} + 30$
 h 38. 10. $= OPq$. therefore OP is $v \sqrt{675} + v \sqrt{75}$. viz.
 a first binomial.

See Scheme 57.

PROP. LVII.



If a space AD be contained
 under a rational line AB and
 a third binomial line AC
 (AE + EC) the right line OP
 which contains in power
 the space AD, is irrational,
 and called a second binomial
 line.


As above, $OPq = AD$.
 p also the rectangles AH,
 GI, that is OMP, MPq
 are μa . a Likewise EK
 or OMP is μr . b there-
 fore OP is a second bi-
 medial.

a hyp. and
 22. 10.
 b 39. 10.

In numbers; let there be AB 5. AC $\sqrt{32} + \sqrt{34}$. wherefore AD is $\sqrt{800} + \sqrt{600} = OPq$.
 and so OP is $v \sqrt{450} + v \sqrt{50}$. that is a 2.
 bimed.

PROP.

PROP. LVIII.



If a space AD be comprehended under a rational line AB and a fourth binomial AC ($AE + EC$) the right line OP containing the space AD in power, is that irrational line which is called a Major line.

For again, OMq a $\square$ a lem. 34. MPq ; and the rectangle IO . AI , i. e. $OMq + MPq$ b b hyp. and is ρv . c also EK or OMP 20. 10. is μv . d therefore OP ($\sqrt{c}$ hyp. and AD) is a Major line. 22. 10. Which was to be dem. d 40. 10.

In numbers; let there be AB 5. and AC $4 + \sqrt{8}$. then the rectangle AD is $20 + \sqrt{200}$. wherefore OP is $\sqrt{20 + \sqrt{200}}$.

PROP. LIX.

If a space AD be contained under a rational line AB , and a fifth binomial AC , the right line OP which contains the space AD in power, is that irrational line, which is a line containing a rational and a medial rectangle in power.

Again OMq $\square$ MPq . and the rectangle AI or $OMq + MPq$ is μv . a Likewise the rectangle a As in the EK or OMP is ρv , b therefore OP ($\sqrt{AD}$) contains in power ρv and μv . Which was to be dem. b 41. 10.

In numbers, let there be AB 5. and AC $2 + \sqrt{8}$. then the rectangle $AD = 10 + \sqrt{200} = OPq$, wherefore OP is $\sqrt{10 + \sqrt{200}}$.

PROP.


PROP. LX.

If a space AD be contained under a rational line AB and a sixth binomial AC ($AE + EC$) the line OP containing the space AD in power is irrational, which contains in power two medial rectangles.

As often before, $OMq \sqsupset MPq$, and $OMq + MPq$ is μv . and also the rectangle $(EK) OMP$ is μv . a therefore $OP = \sqrt{AD}$ contains in power 2 μa . Which was to be dem.

In numbers, let there be AB 5. $AC \sqrt{12} + \sqrt{8}$. therefore the rectangle AD or OPq is $\sqrt{300} + \sqrt{200}$. and so OP is $\sqrt{\sqrt{300} + \sqrt{200}}$.

Lemma.



Let a right line AB be unequally divided in C , and let AC be the greater portion, and upon some line DE apply the rectangles $DF = ABq$, and $DH = ACq$, and $IK = CBq$. and let LG be divided equally in

M , and also MN drawn parallel to GF .

a 4.2. and I say 1. The rectangle ACB is $= LN$ or MF .

3. ax. 1. a For 2 $ACB = LF$.

b 7. 2. 2. $DL \sqsubset LG$. for $DK (ACq + CBq) \sqsubset LF$

c 1. 6. (2 ACB) therefore being DK , LF are of equal

d 16. 10. altitude, c DL shall be $\sqsubset LG$.

3. If $AC \sqsupset CB$, d then shall the rectangle DK be $\sqsupset ACq$ and CBq .

e lem. 26. 4. Also $DL \sqsupset LG$. for $ACq + CBq \sqsupset 2$

10. ACB , i. e. $DK \sqsupset LF$. but $DK. LF \sqsupset DL$.

f 10. 10. LG . f therefore $DL \sqsupset LG$.

g 1. 6. 5. Moreover $DL \sqsupset \sqrt{DLq - LGq}$. For

h 17. 6. $ACq. ACB \sqsupset ACB. CBq$. that is, $DH. LN \sqsupset$

k hyp. $LN. IK$. e wherefore $DI. LM \sqsupset LM. IL$. h

l 10. 10. therefore $DI \times IL = LMq$. therefore seeing ACq

m 18. 10. k $\sqsupset CBq$, that is, $DH \sqsupset IK$. and l so DI

$\sqsupset IL$, m shall DL be $\sqsupset \sqrt{DLq - LGq}$. Which was to be dem.

c. But

6. But if ACq be put $\square$ CB, then shall DL be $\sqrt{DLq - LGq}$.

This Lemma is preparatory to the following Propositions.

PROP. LXI.

The square of a binomial line (AC + CB) applied unto a rational line DE, makes the breadth DG a first binomial line.

Those things being supposed, which are described and demonstrated in the next preceding Lemma; because AC, CB are $\dot{p}$ $\square$, $\dot{b}$ the rect- angle DK shall be $\square$ ACq. $\dot{c}$ and so DK is $\dot{p}\dot{v}$. $\dot{c}$ therefore DL $\square$ DE $\dot{p}$. but the rectangle ACB, and so $\dot{z}$ ACB (LF) $\dot{e}$ is $\dot{p}\dot{v}$. $\dot{f}$ therefore the latitude LG is $\dot{p}$ $\square$ DE. $\dot{g}$ therefore also DL $\square$ LG also DL $\square$ $\sqrt{DLq - LGq}$. from whence it follows that DG is a first binomial. Which was to be dem.

$\dot{a}$ hyp. $\dot{b}$ lem. 60. $\dot{c}$ sch. 12. $\dot{d}$ 21. 10. $\dot{e}$ 22. and $\dot{f}$ 23. 10. $\dot{g}$ 13. 10. $\dot{h}$ lem. 60. $\dot{i}$ 10. $\dot{k}$ 1. def. 10. $\dot{l}$ 10.

PROP. LXII.

The square of a first bimedial line (AC + CB) being applied to a rational line DE, makes the breadth DG a second binomial line.

The aforesaid Lemma being again supposed; the rectangle DK $\square$ ACq. $\dot{a}$ therefore DK is $\dot{p}\dot{v}$. $\dot{b}$ therefore the breadth DL is $\dot{p}$ $\square$ DE. But because the rectangle ACB, and so LF ($\dot{z}$ ACB) $\dot{c}$ is $\dot{p}\dot{v}$, $\dot{d}$ shall LG be $\dot{p}$ $\square$ DE. $\dot{e}$ therefore DL, LG are $\square$ $\dot{f}$ also DL $\square$ $\sqrt{DLq - LGq}$. from whence it is clear that DG is a second binomial. Which was to be dem.

$\dot{a}$ 24. 10. $\dot{b}$ 23. 10. $\dot{c}$ hyp. and sch. 22. 10. $\dot{d}$ 2. 10. $\dot{e}$ 13. 10. $\dot{f}$ lem. 10. $\dot{g}$ 10. $\dot{h}$ 10. $\dot{i}$ 2 def. 48. $\dot{j}$ 10.

PROP.

PROP. LXIII.

The square of a second binomial line ($AC+CB$) applied to a rational line DE makes the breadth DG a third binomial line.

a hyp. and As in the prec. DL is $\dot{p} \sqsubset DE$. Further-
 24. 10. more because the rectangle ACB , and so LF (2
 b 23. 10. ACB) a is $\mu\nu$. b therefore shall LG be $\dot{p} \sqsubset DE$.
 c lem. 60. c Moreover $DL \sqsubset LG$. and also $DL \sqsubset \sqrt{\quad}$
 10. $DLq - LGq$. d therefore DG is a third bino-
 d 3. def. 48. mial. Which was to be dem.
 10.

PROP. LXIV.

The square of a Major line ($AC+CB$) applied to a rational line DE , makes the breadth DG a fourth binomial line.

a hyp. and Again $ACq+CBq$. i. e. DK a is $\dot{p}\nu$ b therefore
 sch. 12. 10. DL is $\dot{p} \sqsubset DE$, also ACB , and so LF (2 ACB)
 b 21. 10. c is $\mu\nu$. d therefore LG is $\dot{p} \sqsubset DE$, e and con-
 c hyp. and sequently $DL \sqsubset LG$. Lastly, because $AC \sqsubset$
 24. 10. BC . f shall DL be $\sqsubset DLq - LGq$. g whence DG
 d 23. 10. is a fourth binomial. Which was to be dem.
 e 13. 10.
 f lem. 60.

PROP. LXV.

The square of a line containing in power a rational and a medial rectangle ($AC+CB$) applied to a rational line DE makes the breadth DG a fifth binomial.

a 23. 10. Again, DK is $\mu\nu$. a therefore DL is $\dot{p} \sqsubset DE$.
 b 21. 10. also LF is $\dot{p}\nu$. b therefore LG is $\dot{p} \sqsubset DE$. c
 c 13. 10. therefore $DL \sqsubset LG$. d likewise $DL \sqsubset \sqrt{\quad}$
 d lem. 60. $DLq - LGq$. e and so by consequence DG is a
 10. fifth binomial. Which was to be dem.
 e 5. def. 48.
 10.


PROP. LXVI.

The square of a line containing in power two medial rectangles ($AC+CB$) applied to a rational line DE , makes the breadth DG a sixth binomial line.

As

As before, DL and LG are $\perp$ DE. Buta *hyp.*
for that ACq + CBq (DK) a $\perp$ ACB, b and b 14. 10.
so DK $\perp$ LF (2 ACB) and also DK. LF c :: c 1. 6.
DL. LG. d therefore shall DL be $\perp$ LG. e d 10. 10.
Lastly, DL $\perp$ $\sqrt$ DLq - LGq. f by which it c *lem.* 60.
appears that DG is a sixth binomial. 10.
f 6. def. 48. 10.

Lemma.



Let AB, DE be $\perp$. and make AB. DE :: AC. DF.

I say 1. AC $\perp$ DF. as appears by 10. 10.

also CB $\perp$ FE. a because AB. DE :: CB. FE. a 19. 5.

2. AC. CB :: DF. FE. For AC. DF :: AB. DE :: CB. FE. therefore inversely AC. CB :: DF. FE.


3. Therefore angle ACB $\perp$ DFE. For ACq. ACB b :: AC. CB c :: DF. EF :: DFq. DFE. wherefore b 1. 6.
by inversion ACq. DFq :: ACB. DFE. therefore c *before*
being ACq $\perp$ DFq. d shall ACB be $\perp$ DFE. d 10. 10.

4. ACq + CBq $\perp$ DFq + EFq. For because ACq. CBq e :: DFq. FEq. therefore by addition e 22. 6.
ACq + CBq. CBq :: DFq + FEq. FEq. there-
fore being CBq $\perp$ FEq, f shall also ACq + f 10. 10.
CBq be $\perp$ DFq + FEq.

5. Hence, If AC $\perp$ or $\perp$ CB. g then like- g 10. 10.
wise shall DE be $\perp$ or $\perp$ EF.

PROP.

PROP. LXVII.



A line DE, commensurable in length to a binomial line (AC+CB) is itself a binomial line, and of the same order.

Make AB. DE :: AC. DF. *a* then are AC, DF
a lem. 66. $\square$ *a* and CB, FE $\square$. whence being that AC
 10. and CB *b* are ρ $\square$, *c* thence DF, FE ρ $\square$ there-
b hyp. fore DE is a binomial. But for that AC. CB *a* ::
c lem. 66. DF. FE. if AC $\square$ or $\square$ $\sqrt{ACq - BCq}$. *d*
 10. and sch. then in like manner DF $\square$ or $\square$ $\sqrt{DFq -$
 12. 10. FEq. also if AC $\square$ or $\square$ ρ propounded, *e*
d 15. 10. then shall DF be $\square$ or $\square$ ρ propounded. But
e 12. 10. if CB $\square$ or $\square$ ρ . likewise FE $\square$ or $\square$ ρ .
 and 14. 10. If both AC, CB, $\square$ ρ . *g* that is, whatsoever
f by def. 48. binomial AB is, DE shall be of the same order.
 10. Which was to be dem.

g 14. 10.

PROP. LXVIII.


A line DE commensurable in length to a binomial line (AC+CB) is also a binomial line, and of the same order.

Make AB. DE :: AC. DF. *b* therefore AC $\square$
a 12. 6. DF and CB $\square$ FE. therefore seeing AC and
b lem. 66. CB *c* are μ , *d* also DF, and FE shall be μ , and
 10. for that AC *e* $\square$ CB. *e* therefore FD $\square$ FE.
c hyp. *f* therefore DE is 2μ . Wherefore if the rectan-
d 24. 10. gle ACB be $\rho\nu$. because DFE *b* $\square$ ACB, *g* like-
e 10. 10. wise DFE is $\rho\nu$. and if that be $\mu\nu$. *h* this shall
f 38. 10. be $\mu\nu$ too. *k* That is, whether AB be 1 bimed.
g sch. 12. or 2 bimed. DF shall be of the same order.
 10. Which was to be dem.
k 38, or 39
 10.

PROP.

e 14. 10. e shall DFq . FEq be $\sqsupset$ DFE. f from whence
 f 42. 10. it follows that DE contains in power 2 μ .
 Which was to be dem.

P R O P. LXXII.




If a rational rect-
 angle A, and a me-
 dial B, be composed
 together, these four
 irrational lines will
 be made; either a
 binomial, or a first
 bimedial, or a ma-
 jor, or a line containing in power a rational and a
 medial rectangle.

Namely, if $Hq = A \mid B$. then H shall be one of
 the four lines which the Theorem mentions. For
 a cor. 16.6. upon CD the propounded ρ , a make the rectan-
 b 2. ax. 1. gle $CE = A$, and $FI = B$. b and so $CI = Hq$.
 c 21. 10. Whereas then is $A \rho v$, likewise CE is ρv . c there-
 d 23. 10. fore the latitude CF is $\rho \sqsupset$ CD. and because B
 e 13. 10. is μv , also FI shall be μv . d therefore FK is $\rho \sqsupset$
 f 37. 10. CD. e therefore CF, FK are $\rho \sqsupset$. and so the
 g 1. 6. whole CK f is binom. wherefore if $A \sqsubset B$, i. e.
 h 1. def. 48. $CE \sqsubset FI$, g then $CF \sqsubset FK$. therefore if $CF \sqsupset$
 10. $\sqrt{CFq} - FKq$, h likewise CK shall be a 1 bin.
 k 55. 10. and consequently $H = \sqrt{CI}$ k is a bin. If CF
 l 4. def. 48. be supposed $\sqsupset \sqrt{CFq} - FKq$, l then shall CK
 10. be a 4 bin. wherefore H ($\sqrt{CI}$) m is a major
 m 58. 10. line. But if $A \sqsupset B$, g then shall CF be $\sqsupset$ FK,
 n 2. def. 48. consequently if $FK \sqsupset \sqrt{FKq} - CFq$, n then
 10. shall CK be a 2 bin. o wherefore H is a first 2 μ .
 o 56. 10. lastly, if $FK \sqsupset \sqrt{FKq} - CFq$; p then CK
 p 5. def. 48. shall be a fifth binom. q whence H shall contain
 10. in power ρv and μv . Which was to be dem.
 q 59. 10.

P R O P.

PROP. LXXIII.



If two medial rect-
angles A, B , incommensurable to one
another be composed
together, the two re-
maining irrational
lines are made, either
a second binomial, or

a line containing in power two medial rectangles.

As H containing in power $A + B$ is one of the
said irrational lines. For upon CD propounded
draw the rectangle $CE = A$, and $FI = B$. whence
 $HQ = CI$. Therefore because CE and FI are μ . a hyp.
the latitudes CF, FK , shall be ρ $\square$ CD . also b 23. 10.
because CE $\square$ FI , and $CE \cdot FI$ $c :: CF \cdot FK$, d 1. 6.
therefore CF $\square$ FK . e therefore CK is a 3. bin. d 10. 10.
namely, if CF $\square$ $\sqrt{CFq} - FKq$. whence H e 3. def. 48.
 $= \sqrt{CI}$ f shall be a second z μ . But if CF $\square$ f 57. 10.
 $\sqrt{CFq} - FKq$, g then CK shall be a 6 binom. g 6. def. 48.
and consequently H contains in power 2 μ . h 60. 10.
Which was to be dem.

Here begin the Senaries of lines irrational by
Subtraction.

PROP. LXXIV.

If from a rational line DF a
rational line DE , commensura-
ble in power only to the whole
 DF , be taken away, the residue EF is irrational,
and is called an Apotome or residual line.

For EFq a $\square$ DEq ; b but DEq is ρ ; c there- a leni. 26.
fore EF is ρ . Which was to be dem. $10.$

In numbers, let there be $DF, 2$. $DE, \sqrt{3}$. then b hyp.
 EF shall be $2 - \sqrt{3}$. c 10. $\&$ 11. $def. 10.$

Q 2

PROP.

PROP. LXXV.

D E F If from a medial line DF , a medial line DE commensurable only in power to the whole DF , and comprehending with the whole DF a rational rectangle be taken away, the remainder EF is irrational, and is called a first residual line of a medial.

a sch. 16. For EFq a $\square$ to the rectangle FDE . therefore seeing FDE b is ρv , c BF shall be ρ . Which was to be dem.

b hyp. In numbers, let DF be $v \sqrt{54}$, and DE $v \sqrt{24}$. therefore EF is $v \sqrt{54} - v \sqrt{24}$.

PROP. LXXVI.

D E F If from a medial line DF , a medial line DE be taken away being incommensurable only in power to the whole DF , and comprehending together with the whole line DF a medial rectangle, the remainder EF is irrational, and is called a second residual of a medial line.

a hyp. Because DFq and DEq a are μv $\square$, b therefore shall $DFq + DEq$ be $\square$ DEq . c wherefore b 16. 10. $DFq - DEq$ is μv . also the rectangle FDE , c 24. 10. $DFq \cdot DEq$ is μv . also the rectangle FDE , c d cor. 7. 1. and so $2 FDE$, a is μv . therefore EFq (d $DFq + DEq - 2 FDE$) e is ρv . wherefore EF is ρ . Which was to be dem.

In numbers, let DF be $v \sqrt{18}$. and DE $v \sqrt{8}$. then EF $v \sqrt{18} - v \sqrt{8}$.

PROP. LXXVII.

A B C If from a right line AC be taken away a right line AB being incommensurable in power to the whole BC , and making with the whole AC that which is composed of their squares rational, and the rectangle contained under them medial, the remainder BC is irrational, and is called a Minor line.

a hyp. For $ACq + ABq$ a is ρv , but the rectangle ACB b is μv . b therefore $2 CAB \square ACq + ABq$ ($2c$ CAB

b sch. 12. 10. 6 7. 2.

CAB+BCq.) d therefore ACq+ABq $\sqsupset$ BCq. ed 17. 10.
therefore BC is p. Which was to be dem. e 11. def.

In numbers, let AC be $\sqrt{18} + \sqrt{108}$; AB 10.
 $\sqrt{18} - \sqrt{108}$. then BC is $\sqrt{18} + \sqrt{108} -$
 $\sqrt{18} - \sqrt{108}$.


PROP. LXXVIII.

D——E——F If from a right line DF
be taken away a right line
DE, being incommensurable in power to the whole
line DF, and with the whole DF making that which
is composed of their squares medial, and the rectangle
contained under the same lines rational, the line re-
maining EF is irrational, and is called a line making
a whole space medial with a rational space.

For 2 FDE a is p. b and DFq+DEq is μ . c a hyp. and
therefore 2 FDE $\sqsupset$ DFq+DEq d (2 FDE+EFq) sch. 12. 10.
e therefore EF is p. Which was to be dem. b hyp.

In numbers, let DF be $\sqrt{216} + \sqrt{72}$; DE c sch. 12.
 $\sqrt{216} - \sqrt{72}$. therefore EF is $\sqrt{216} + 10$.
 $\sqrt{72} - \sqrt{216} - \sqrt{72}$. d 7. 2.

PROP. LXXIX.



If from a right line DF be taken
away a right line DE incommensu-
rable in power to the whole DE, and
which together with the whole makes that which is
composed of their squares medial, and the rectangle
contained under them, also medial and incommensu-
rable to that which is composed of their squares, the
remainder is irrational, and is called a line making
a whole space medial with a medial space.


For 2 FDE, and FDq+DEq a are μ a; b there- a hyp. and
fore EFq (c DFq+DEq - 2 FDE) is p. d and 24. 10.
so consequently EF is p. Which was to be dem. b 27. 10.

In numbers, let DF be $\sqrt{180} + \sqrt{60}$. DE c cor. 7. 2.
 $\sqrt{180} - \sqrt{60}$. then EF shall be $\sqrt{180} + 10$. d 11. def.
 $\sqrt{60} - \sqrt{180} - \sqrt{60}$. 10.

Q 3

Lemma.

Lemma.



If there be the same excess between the first magnitude BG, and the second C (MG) as is between the third magnitude DF and the fourth H (EF;) then alternately, the same excess shall be between the first magnitude BG and the third DF, as is between the second C and the fourth H.

a hyp.

For because that *a* to the equals BM, DE, are added the unequals MG, EF, that is, CH; *b* 15. *bx.1.* the excess of the wholes BG, DF, *b* shall be equal to the excess of the parts added C, H. Which was to be dem.

Coroll.

Hence, Four magnitudes Arithmetically proportional, are alternately also Arithmetically proportional.

PROP. LXXX.

A ——— *B D C* To an Apotome or residual line *AB* only one rational right line *BC*, be-

a 22. 10. ing commensurable in power only to the whole *AB*,
b 22. 10. is congruent, or can be joined.

c cor. 7. 2. If it be possible, let some other line *BD* be added to it; *a* then the rectangles *ACB*, *ADB*, *b*
10. and so consequently the double of them are *ua.*

e hyp. and wherefore seeing $ACq + BCq - 2ACBc = ABqc$
27. 10. $= ADq + DBq - 2ADB.$ therefore alternately

f sch. 12. $AGq + BCq - ADq + DBqd = 2ACB - 2ADB.$

10. But $ACq + BCq - ADq + DBqe$ is *py.* therefore

g 27. 10. $2ACB - 2ADB$ is *py.* Which is absurd.


PROP.

PROP. LXXXI.

To a first medial residual line
 $A \quad B \quad D \quad C$ AB only one medial right line
 BC , being commensurable only
in power to the whole, and comprehending with the
whole line a rational rectangle, can be joined.

Conceive BD to be such a line as may be
joined to it; then because ACq and BCq , as well
as ADq and BDq are $ua \sqcap .b$ also $ACq + BCq$, a hyp.
and $ADq + BDq$ shall be $ua . c$ but the rectangles b 10. and
 ACB, ADB, d and so $2 ACE$ and $2 ADB$ are $p\alpha . 24. 10.$
therefore $2 ACB = 2 ADB$, f that is, $ACq + c$ hyp.
 $BCq = ADq + BDq$ is $py . g$ Which is absurd. d sch. 12.
10.

PROP. LXXXII.




Unto a second medial
residual line AB only
one medial right line
 BC , commensurable only
in power to the whole,
and with it containing
a medial rectangle, can
be joined.

If it be possible, let
some other line BD be
added to it; and upon EF p , make the rectangle
 $EG = ACq + BCq$; as also the rectangle $EL = a$ 4. 2. and
 $ADq + BDq$. likewise $EI = ABq$. Now $2 ACB$ 3. ax. 1.
 $+ ABq = ACq + BCq = EG$. therefore seeing EI b hyp.
 $= ABq$, a also GK shall be $= 2 ACB$. moreover c 24. 10.
 ACq and BCq b are $ua \sqcap . c$ therefore $EG (ACq$ d 23. 10.
 $+ CBq)$ is $uv . d$ therefore the breadth EH is p e hyp.
 $\sqcap EF$. e Further, the rectangle ACB f and so f 24. 10.
 $2 ACB (KG)$ is $uv . d$ therefore KH is also B , g lem. 26.
 EF . lastly, because $ACq + BCq (EG)$ g $\sqcap 2 ACB$ h 1. 6.
 (KG) and $EG . KG$ $b :: EH . KH$. k therefore h 1. 6.
 $EH \sqcap HK$. l therefore EK is a residual line, k 10. 10.
where HK is congruent. by the same reason l 74. 10.

also shall KM be congruent to the said EK, which is repugnant to the 80. Prop. of this Book.

P R O P. LXXXIII.

 To a Minor line AB only one right line BC can be joined being incommensurable in power to the whole; and making together with the whole line that which is composed of their squares rational, and the rectangle which is contained under them medial.


Conceive any other BD to be congruent to it; therefore whereas $ACq + BCq$, and $ADq + DBq$ are $\rho\alpha$, their excesses ($2b$ ACB $:-$ 2 ADB) a hyp. DBq a are $\rho\alpha$, their excesses ($2b$ ACB $:-$ 2 ADB) b lem. 97. c is $\rho\gamma$, Which is absurd; because ACB and ADB $10.$ are $\mu\alpha$ by the Hyp.

c sch. 27.

$10.$

d 27. $10.$

P R O P. LXXXIV.

 Unto a line (AB) making with a rational space a whole space medial only one right line BC can be joined, being incommensurable in power to the whole, and making together with the whole that which is composed of their squares medial, and the rectangle which is contained under them rational.


Suppose some other BD to be congruent also to it; a then the rectangles ACB, ADB, b and b sch. 12. so 2 ACB and 2 ADB are $\rho\alpha$. therefore 2 ACB $10.$ $:-$ 2 ADB, c that is, $ACq + BCq$ $:-$ $ADq + BDq$ c lem. 79. d is $\rho\gamma$. Which is absurd. since $ACq + BCq$, and $10.$ $ADq + BDq$ are $\mu\alpha$ by the Hyp.

d sch. 27.

$10.$

P R O P.

PROP. LXXXV.



To a line AB , which with a medial space makes a whole space medial, can be joined only one right line BC , incommensurable in power to the whole, and making with the whole both that which is composed of their squares medial, and the rectangle

which is contained under them medial and incommensurable to that which is composed of their squares.

Those things being supposed which are done and shewn in the 82. Prop. of this Book; it is clear that EH and KH are $\perp$ EF . Besides, being that $ACq + CBq$, that is, the rectangle EG , a is $\perp$ ACB . b and so $EG \perp 2 ACB$ (KG ;) a hyp. and $EG.KGc :: EH.KH$; shall EH be $\perp$ KH . b 14. 10. therefore EK is a residual line, and the line congruent to it is KH . In like manner may KM be shewn to be congruent to the said residual EK , against the 80. Prop. of this Book.

Third Definitions.

A Rational line and a residual being propounded, if the whole be more in power than the line joined to the residual, by the square of a right line commensurable unto it in length; then

I. If the whole be commensurable in length to the rational line propounded, it is called a first residual line.

II. But if the line adjoined be commensurable in length to the rational line propounded, it is called a second residual line.

III. If neither the whole nor the line adjoined be commensurable in length to the rational line propounded; it is called a third residual line.

More-

Moreover, if the whole be in more power than the line adjoined by the square of a right line incommensurable to it in length, then

IV. If the whole be commensurable in length to the rational line propounded, it is called a fourth residual line.

V. But if the line adjoined be commensurable in length to the rational line propounded, it is a fifth residual.


VI. If neither the whole nor the line adjoined be commensurable in length to the rational line propounded, it is termed a sixth residual line.

PROP. LXXXVI, 87, 88, 89, 90, 91.

A....4 C....5 E To find out a first, second, third, fourth, fifth, and sixth residual line.

H ——— G Residual lines are found out by subducting the less names or parts of binomials from the greater. *ex. gr.* Let $6 + \sqrt{20}$ be a first binom. then shall $6 - \sqrt{20}$ be a first residual. So that it is not necessary to repeat more concerning the finding of them out.

Lemna.



Let AC be a rectangle contained under the right lines AB, AD. Let AD be drawn forth to E, and DE equally divided in F. and let the rectangle AGE be = FEq. and the rectangles AI, DK, FH, be finished. Then let the square LM = AH be made, and the square NO = GI; and the lines NSR, OST, produced.


I say 1. the rectangle AI = LM + NO = TOq + SOq, which appears by the constr.

2. The

2. The rectangle $DK = LG$. For because the rectangle AGE $a = FEq$. b thence are AG, FE , a *constr.* GE , c and so AH, FI, GI d , that is, LM, FI , b 17. 6. NO e ; but LM, LO, NO d are e ; therefore c 1. 6. $FI = e$ $LO f = DK = g$ NM . *d sch. 2. 6.*
3. Hence, $AC = AI - DK - FI = LM + NO - LO$ e 9. 5. $-NM = TR$. f 36. 1.
4. b It is manifest that DF, FE, DE , are $\perp$. g 43. 1.
5. If $AE \perp DE$, and $AE \perp \sqrt{AEq - DEq}$, h 16. 10. k then shall AG, GE, AE be $\perp$. k 18. and
6. Also, because $AE \perp DE$, m thence shall AE, FE be $\perp$. n and so AI, FI , that is, LM l hyp. NO and LO are $\perp$. m 13. 10.
7. Because $AG * \perp GE$, n shall AH, GI , that is, LM, NO be $\perp$. n 1. 6. and $10. 10.$
8. But because $AE \perp DE$, o therefore shall FE, GE be $\perp$, n and so the rectangle $FI \perp GI$, o 14. 10. that is, $LO \perp NO$. wherefore seeing LO, NO $p :: p$ 2. 6. $TS. SO$. q therefore shall TS, SO be $\perp$. q 10. 10.
9. If AE be put $\perp \sqrt{AEq - DEq}$, r then shall AG, GE, AE be $\perp$. r 19. 10. $17. 10.$
10. f Wherefore the rectangles AH, GI , that is, TOq, SOq shall be $\perp$. f 1. 6. and $10. 10.$

PROP.

PROP. XCII.



a hyp.
b 12. 10.
c 20. 10.
d lem. 91.
10.
e 74. 10.

If a space AC be contained under a rational line AB , and a first residual line AD ($AE - DE$) the right line TS , which contains the space AC in power, is a residual line.

Use the foregoing Lemma for a preparatory to the demonstration of this Prop. Therefore $TS = \sqrt{AC}$. Also AG, GE, AE , are $\square$; therefore since $AE \square a AB$ ρ , b also AG and GE shall be $\square AB$. c therefore the rectangles AH and GI , that is, TOq and SOq are ρa . d Likewise TO, SO , are $\rho \square$. e and consequently TS is a residual line. Which was to be dem.

PROP. XCIII.

See the preceding Scheme.

If a space AC be contained under a rational line AB and a second residual AD ($AE - DE$) the right line TS , containing the space AC in power, is a first medial residual line.

Again, by the foregoing Lemma, AG, GE, AE are $\square$. therefore a since AE is $\rho \square AB$, b 13. 10. b also AG, GE , shall be $\rho \square AB$. c therefore c 22. 10. the rectangles AH, GI , that is, TOq, SOq are ρa . d lem. 74. d likewise $TO \square SO$. Lastly, because DE e 10. $\square AB$ ρ , f the right angle DI , and the half e hyp. thereof DK or LO , that is, TOS shall be ρv . g f 20. 10. from whence it follows that TS ($\sqrt{AC}$) is a g 75. 10. first medial residual. Which was to be dem.

PROP.

PROP. XCIV.

See Scheme 92.

If a space AC be contained under a rational line AB and a third residual AD ($AE - DE$) the right line TS containing in power the space AC is a second medial residual line.

As in the former, TO and SO are μ . Therefore because DE is $\rho \sqsubset AB$. b the rectangle a hyp. DL , c and so DK , or TOS , shall be $\mu\nu$. d there- b 21. 10. c 24. 10. d 76. 10. $TS = \sqrt{AC}$ is a second medial residual. *Which was to be dem.*

PROP. XCV.

See Scheme 92.

If a space AC be contained under a rational line AB and a fourth residual AD ($AE - DE$) the right line TS containing the space AC in power, is a minor line.

As before, TO is $\sqsupset SO$. Therefore because a lem. 91. AE is $\rho \sqsubset AB$, c shall AI ($TOq + SOq$) be ρ . b but, as before, the rectangle TOS is $\mu\nu$. d b hyp. c 20. 10. d 77. 10. $TS = \sqrt{AC}$ is a Minor line. *Which was to be dem.*

PROP. XCVI.

See Scheme 92.


If a space AC be contained under a rational line AB and a fifth residual AD ($AE - DE$) the right line TS containing in power the space AC , is a line which makes with a rational space the whole space medial.

For again TO is $\sqsupset SO$. therefore since AE is a hyp. $\rho \sqsubset AB$. b also AI , that is, $TOq + SOq$, shall be ρ . c 22. 10. d be $\mu\nu$. But, as in the 93. the rectangle TOS is $\rho\nu$. c 78. 10. d whence $TS = \sqrt{AC}$ is a line which with $\rho\nu$ makes a whole $\mu\nu$. *Which was to be dem.*


PROP.

PROP. XCVII.

A D FGE



B L C K H I



Q R M

If a space AC be contained under a rational line AB, and a sixth residual AD (AE-DE) the right line TS containing in power the space AC, is a line making with a medial rectangle, a whole space medial.

As often above, TO $\sqsupset$ SO. also, as in 96. TOq + SOq is $\mu\nu$. but the rectangle TOS is $\rho\nu$, as in 94. a Lastly, TOq + SOq $\sqsupset$ TOS. b therefore TS = $\sqrt{AC}$ is a line which with $\mu\nu$ makes

a whole $\mu\nu$. Which was to be dem.]

a lem. 91.

20.

b 79. 10.

Lemma.

* cor. 16. 6.

A B C

D G M IL



E N HK

Upon a right line DE* apply the rectangles DF=ABq, and DH=ACq, and IK=BCq. and let GL be bisected in M, and the line MN drawn parallel to GF.

Then 1. The rectangle DK is = ACq + BCq. as the construction manifests.


a constv. 2. The rectangle ACB = GN or MK. For DK b 7. 2. a = ACq + BCq b = 2 ACB + ABq. but ABq c 3. ax. 1. a = DF. therefore GK c = 2 ACB. and d con- d 7. ax. 1. sequently GN or MK = ACB.

e 1. 6. 3. The rectangle DIL = MLq. For because f i 7. 6. ACq. ACB e :: ACB. BCq, that is, DH, MK :: MK. IK. e thence is DI.ML :: ML.IL. f there- fore DIL = MLq.

4. If

4. If AC be taken $\sqsupset BC$, then DK shall be $\sqsupset ACq$. For $ACq + BCq$ (DK) $g \sqsupset ACq$. g 16. 10.
 5. Likewise $DL \sqsupset \sqrt{DLq - GLq}$. For because DH (ACq) $\sqsupset IK$ (BCq) h thence shall DI be h 10. 10.
 $\sqsupset IL$. k therefore $\sqrt{DLq - GLq} \sqsupset DL$. k 18. 10.
 6. Also $DL \sqsupset GL$. For $ACq + BCq \sqsupset l$ lem. 26.
 $2 ACB$. that is, $DK \sqsupset GK$. m therefore DL 10.
 $\sqsupset GL$. m 10. 10.
 7. But if AC be taken $\sqsupset BC$, n then DL shall n 19. 10.
 be $\sqsupset \sqrt{DLq - GLq}$.

PROP. XCVIII.



The square of a residual line AB ($AC - BC$) applied to a rational line DE , makes the breadth DG a first residual line. a hyp. b lem. 97.
 Do as is enjoined in the Lemma next preceding. Then because AC, BC , a are $\dot{p} \sqsupset b$ 10. c sch. 12.
 also DK ($ACq + BCq$) shall be $\sqsupset ACq$. c there- d 21. 10.
 fore DK is $\dot{p}v$. d wherefore DL is $\dot{p} \sqsupset DE$. e 22. and
 Likewise the rectangle GK ($2 ACB$) is μv . f 24. 10.
 therefore GL is $\dot{p} \sqsupset DE$. g and consequently f 23. 10.
 $DL \sqsupset GL$. h But $DLq \sqsupset GLq$. k therefore g 13. 10.
 DG is a residual, l and that of the first order h sch. 12.
 (because $m AC \sqsupset BC$, and therefore $DL \sqsupset$ 10.
 $\sqrt{DLq - GLq}$) Which was to be dem. k 74. 10.
 l 1. def. 85.
 10 .
 m lem. 97.
 10 .

PROP.

PROP. XCIX.

See the following Scheme.

The square of a first medial residual line AB ($AC - BC$) applied to a rational line DE , makes the breadth DG a second residual line.

Supposing the foregoing Lemma; because AC and BC are μ . $\square$, b thence shall DK ($AC - BC$) be $\square$ AC . c wherefore DK is μ . d therefore DL is ρ $\square$ DE . e also GK ($2 ACB$) is ρ . f therefore GL is ρ $\square$ DE , g wherefore DL $\square$ GL . h But DLq $\square$ GLq . k therefore DG is a residual line: and because DL is $\square$ $\sqrt{DLq}$ $\square$ GLq . m therefore shall DG be a second residual. Which was to be dem.

g 13. 10.

h sch. 12.

10.

k 74. 10.

l lem. 97.

10.

m 2. def.

85. 10.

a 23. 10.

b lem. 26.

10.

c 1. 6. and

10. 10.

d sch. 12.

10.

e 74. 10.


f 3. def. 85.

10.

g lem. 97.

10.

PROP. C.



The square of a second medial residual line AB ($AC - BC$) applied to a rational line DE makes the breadth DG a third residual line.

Again DK is μ . a wherefore DL is ρ $\square$ DE . also DGK is μ . a

whence GL is ρ $\square$ DE . b likewise DK $\square$ GK . c wherefore DL $\square$ GL . d but DLq $\square$ GLq . e therefore DG is a residual line, and that of the third order, g because DL $\square$ $\sqrt{DLq}$ $\square$ GLq . Which was to be dem.

PROP. CI.

See the foregoing Scheme.

The square of a Minor line AB ($AC - BC$) applied

to a rational line DE, makes the breadth DG a fourth residual.

As before, ACq + BCq, that is, DK is $\rho\nu$. *aa* 21. 10. therefore DL is $\rho \sqsupset$ DE. but the rectangle ACB, * *hyp.* and so GK (2 ACB) * is $\mu\nu$. *b* wherefore GL is $\rho \sqsupset$ DE. *c* therefore DL $\sqsupset$ GL. *b* but *c* 13. 10. DLq $\sqsupset$ GLq, and because * ACq $\sqsupset$ BCq, *ed sch.* 12. thence shall DL be $\sqsupset \sqrt{DLq - GLq}$. *f* there-10. fore DG has the conditions required to a fourth *lem.* 97. residual. Which was to be dem. 10. f 4.def.85. 10.

PROP. CII.

See Scheme 100.

The square of a line AB (AC - BC) which makes with a rational space the whole space medial, applied to a rational line DE, makes the breadth DG a fifth residual line.

For, as above, DK is $\mu\nu$. *a* wherefore DL is ρ^a 23. 10. $\sqsupset$ DE. also GK is $\rho\nu$. *b* whence GL is $\rho \sqsupset$ DE. *c* therefore DL $\sqsupset$ GL. *d* but DLq $\sqsupset$ GLq. *c* 13. 10. Moreover DL $\sqsupset \sqrt{DLq - GLq}$. wherefore *d sch.* 12. DG *f* is a fifth residual. Which was to be dem. 10. e *lem.* 97. 10.

PROP. CIII.

See the last Scheme.

The square of a line AB (AC - BC) making with a medial space the whole space medial, applied to a rational line DE, makes the breadth DG a sixth residual line.

As above DK and GK are μa ; *a* wherefore *a* 23. 10. DL and GL are $\rho \sqsupset$ DE. also DK *b* $\sqsupset$ GK. *b hyp.* and *c* whence DL $\sqsupset$ GL. *d* therefore DG is a re-*lem.* 97. 10. sidual. *b* And whereas ACq $\sqsupset$ BCq. and so DL *c* $\sqsupset \sqrt{DLq - GLq}$, *e* therefore DG shall be *d* 74. 10. sixth residual. Which was to be dem. *e* 6.def.85. 10.

R

PROP.

PROP. CVI.

$\begin{array}{c} A \qquad B \qquad C \\ \text{---} 1 \text{---} \\ \text{---} 1 \text{---} \\ D \qquad E \qquad F \end{array}$
*A right line DE commen-
surable to a Minor line AB
(AC - BC) is it self also a
Minor line.*

Make AB. DE:: AC. DF. *a* then is AC + BC *a* lem. 103.
 $\square$ DF + EF. But AC + BC *b* is a Major line; 10.
 therefore DF + EF is also a Major line; *d* and *b* hyp.
 consequently DF — EF is a Minor line. Which *c* 69. 10.
 was to be dem. *d* 77. 10.

PROP. CVII.



A right line DE commensurable to a line AB (AC — BC) which makes with a rational space the whole space medial, is it self also a line making with a rational space the whole space medial.

For, accordingly as in the former, we may
shew $DF + EF$ to contain in power $\rho\gamma$ and $\mu\gamma$. a 78. 10.
whence $DF - EF$ is a line making, &c.


PROP. CVIII.

$\begin{array}{ccc} A & B & C \\ \hline & 1 & \\ \hline D & E & F \end{array}$

A right line DE commensurable to a line AB (AC - BC) which with a medial space makes the whole space medial, is it self a line making with a medial space the whole space medial.

For according to the preceding $DF + EF$ shall contain in power 2 *ua.* and therefore $DF - EF$ shall be, as in the Prop. 79. 16

The tenth Book of
PROP. CIX.



A medial rectangle B being taken from a rational rectangle $A+B$, the right line H which contains in power the space remaining A , is one of

those two irrational lines, viz. either a residual line, or a Minor line.

Upon CD ρ make the rectangles $CI=A+B$.
a 5. ax. 1. and $FI=B$. whence $CE=A=Hq$. wherefore
b hyp. and because CI is ρv . c therefore CK is $\rho \sqsubset CD$.
constr. but being FI is μv , d shall FK be $\rho \sqsubset CD$.
c 21. 10. whence $CK \sqsubset FK$. f therefore CF is a residual
d 23. 10. line. Wherefore if CK be $\sqsubset \sqrt{CKq - FKq}$,
e 13. 10. g then CF shall be a first residual, b therefore $\sqrt{}$
f 74. 10. $CE (H)$ is a residual line. But if $CK \sqsubset \sqrt{}$
g 1. def. 85. $CKq - FKq$. k then CF shall be a fifth residual;
10. and consequently $H (\sqrt{CE})$ l shall be a
h 92. 10. Minor line. Which was to be dem.
k 4. def. 85.

10.

1 95. 10.

PROP. CX.

See the prec. Scheme.

A rational rectangle B being taken away from a medial rectangle $A+B$, other two irrational lines are made, namely either a first medial residual line,

a 3. ax. 1. or a line making with a rational space the whole
b hyp. and space medial.

Upon CD the propounded ρ make the rectangles $CI=A+B$, and $FI=B$. a whence $CE=A$
e 23. 10. $=Hq$. Therefore because CI is μv , c shall CK
d 21. 10. be $\rho \sqsubset CD$. but because FI is ρv , d thence FK
e 13. 10. $\rho \sqsubset CD$. e whence $CK \sqsubset FK$. f therefore CF
f 74. 10. is a residual, g and that a second. If $CK \sqsubset \sqrt{}$
g 2. def. 85. $\sqrt{CKq - FKq}$, b then $H (\sqrt{CE})$ is a first medial residual.
10. But if $CK \sqsubset \sqrt{CKq - FKq}$. k
h 93. 10. then shall CF be a fifth residual; and l consequently $H (\sqrt{CE})$ shall be a line making μv
k 5. def. 85. with ρv . Which was to be dem.
10. 1 96. 10.

PROP.

PROP. CXI.

See the same Scheme.

A medial space B being taken away from a medial space $A + B$, which is incommensurable to the whole $A + B$, the other two irrational lines are made, viz. either a second medial residual line, or a line making with a medial space the whole space medial.

Upon CD ρ make the rectangles $CI = A + B$, a 3. ax. 1. and $FI = B$. a wherefore $CE = A = Hq$. Because b 23. 10. therefore CI is $\mu\nu$, b thence CK is $\rho \sqsubset CD$, c hyp. and in like manner $FK \rho \sqsubset CD$. Likewise be- d 10. 10. cause CI c $\sqsubset FI$, d therefore $CK \sqsubset FK$. e e 74. 10. wherefore CF is a residual, f namely a third. If f 3. def. 85. $CK \sqsubset \sqrt{CKq - FKq}$, g whence $H (\sqrt{CE})$ shall 10. be a second medial residual, but if $CK \sqsubset \sqrt{g}$ 94. 10. $CKq - FKq$ b then shall CF be a sixth residual. h 6. def. 85. k wherefore A shall be a line making $\mu\nu$ with μ . 10. Which was to be dem. k 97. 10.

PROP. CXII.

A $\overline{\quad}$ A residual line A is not
F D E the same with a binomial
line.



Upon BC propounded ρ make the rectangle $CD = Aq$. Therefore seeing A a 98. 10. is a residual, a BD shall be a first residual, to which let DE be the line con. b 74. 10. gruent, or that may be adjoined. b wherefore BE , c 1. def. 85. DE , are $\rho \sqsubset$. c and $BE \sqsubset BC$. If you con- 10. ceive A to be a binomial, then BD is a first bin, whose names let be BF , FD ; and let BF be $\sqsubset$ d 37. 10. FD . d therefore BF , FD are $\rho \sqsubset$; and BF e e 1. def. 48. $\sqsubset BC$. therefore since $BC \sqsubset BE$ f shall BE be 10. $\sqsubset BF$. g and thence $BE \sqsubset FE$. b therefore f 12. 10. FE is ρ . Likewise because $BE \sqsubset DE$, k shall g cor. 16. 10. FE be $\sqsubset DE$. l wherefore FD is a residual, h sch. 12. and so FD is ρ . but it was shewn ρ , which are 10. repugnant. Therefore A is falsely conceived to k 14. 10. be a binomial. Which was to be dem. 1 74. 10.

R 3


The

The names of the 13 irrational lines differing one from another.

1. A Medial line.
2. A binomial line ; of which there are fix species.
3. A first bimedial line.
4. A second bimedial.
5. A Major line.
6. A line containing in power a rational superficies, and a medial superficies.
7. A line containing in power two medial superficies.
8. A residual line; of which there are also 6 kinds.
9. A first medial residual line.
10. A second medial residual line.
11. A Minor line.
12. A line making with a rational superficies the whole superficies medial.
13. A line making with a medial superficies the whole superficies medial.

Being the differences of breadths do argue differences of right lines, whose squares are applied to some rational line, and it is demonstrated in the pre. Propositions that the breadths which arise from applying of the squares of these 13 lines do differ one from another, it evidently follows that these 13 lines do also differ one from another.

P R O P. CXIII.




The square of a rational line A applied to a binomial BC ($BD + DC$) makes the breadth EC a residual line, whose names EH , CH , are commensurable to the names BD , DO , of the binomial line, and in

in the same proportion ($EH.BD :: CH.DC$) and moreover the residual line EC which is made, is of the same order with BC the binomial.

Upon DC the less name a make the rectangle a *cor.* 16.6.
 $DF = Aq = BE$. whence $BC.CD b :: FC.CE.b$ 14. 6.
 therefore by division, $BD.DC :: FE.EC$. And
 whereas $BD c \sqsubset DC$, d thence FE shall be $\sqsubset c$ *hyp.*
 EC . Take $EG = EC$, and make FG . $GE :: EC.d$ 14. 5.
 CH . Then EH , and CH shall be the names of
 the residual EC , whereunto all is agreeable that
 is propounded in the Theorem. For being that
 by addition $FE.GE (EC) :: EH.CH$. therefore e 12. 5.
 $FH, EH e :: EH.CH f :: FE.EC f :: BD.BC, f$ *before*.
 wherefore since $BD g \sqsubset DC$. h thence shall $EH g$ *hyp.*
 be $\sqsubset CH$. h and $FHq \sqsubset EHq$. Therefore be- h 10. 10.
 cause $FHq.EHq :: FH.CH$. h shall FH be $\sqsubset k$ *cor.* 20.6.
 CH, l and so $FC \sqsubset CH$. Moreover $CD g$ is $\dot{p}$, l 16. 10.
 and $DF (Aq) g$ is $\dot{p}v$. m therefore FC is $\dot{p}$, $\sqsubset m$ 21. 10.
 CD . whence also CH is $\dot{p} \sqsubset CD$. n therefore n *sch.* 12.
 EH, CH are $\dot{p}$ and $\sqsubset$. as before, o therefore EC 10.
 is a residual line, to which CH may be joined. o 74. 10.
 Furthermore $EH.CH f :: BD.DC$. and so by
 inversion $EH.BD :: CH.DC$. whence because
 $CH f \sqsubset DC$, p shall EH be $\sqsubset BD$. But sup- p 10. 10.
 pose $BD \sqsubset \sqrt{BDq - DCq}$. q then shall EH be q 15. 10.
 $\sqsubset \sqrt{EHq - CHq}$. Also if $BD \sqsubset \dot{p}$ propound- r 12. 10.
 ed, then shall EH be $\sqsubset$ to the same $\dot{p}$. f that f 1. *def.* 48.
 is, if BC be a first binomial, t EC shall be a first 10.
 residual. In like manner, if DC be to the $\sqsubset t$ 1. *def.* 85.
 propounded $\dot{p}$. r then is $CH \sqsubset$ to the same $\dot{p}$. u 10.
 that is, if BC be a second binomial, x EC shall u 2. *def.* 48.
 be a second residual; and if this be a third binom. 10,
 then that shall be a third residual, &c. But if x 2. *def.* 85.
 BD be $\sqsubset \sqrt{BDq - DCq}$, y then shall EH be 10.
 $\sqsubset \sqrt{EHq - CHq}$. therefore if BC be a 4th, y 15. 10.
 5th, or 6th binomial, EG shall be likewise a 4th,
 5th or 6th residual. Which was to be dem,

PROP. CXIV.




The square of a rational line A applied to a residual line BC ($BD - CD$) makes the breadth BE a binomial; whose names BE , GE are commensurable to the names BD , BC of the residual line BC , and in the same proportion, and moreover, the binomial line which is made (BE) is of the same order with the residual line (BC).

- a cor. 16. 6. a Make the rectangle $DF = Aq.$ and $BF. FE$
b 12. 6. $b :: EG. GF.$ whence for that $DF = Aq = CE.$
c 14. 6. therefore $BD. BC :: BE. BF,$ therefore by con-
d 19. 5. version of proportion $BD. CD :: BE. FE :: EG.$
e hyp. $GF :: d BG. EG.$ but $BD e \sqsubset CD.$ f therefore
f 10. 10. $BG \sqsubset GE.$ therefore because $BGq. GEqg ::$
g cor. 20. 6. $BG. GF. b$ shall BG be $\sqsubset GF.$ k and so BG
h 10. 10. $\sqsubset BF.$ moreover $BD e$ is $\rho,$ and the rectangle
k cor. 16. $DF (Aq) e$ is $\rho.$ l therefore BF is $\rho \sqsubset BD.$ m
10. therefore also BG is $\rho \sqsubset BD.$ n therefore $BG,$
l 21. 10. GE are $\rho \sqsubset.$ o wherefore BE is a binomial.
m 12. 10. Lastly, because $BD. CD :: BG. GE.$ and inverse-
n sch. 12. ly $BD. BG :: CD. GE.$ and $BD \sqsubset BG.$ p thence
10. shall CD be $\sqsubset GE.$ therefore if CB be a first
o 37. 10. residual, BE shall be a first binomial, &c. as in
p 10. 10. the prec. therefore, &c.

PROP.

PROP. CXV.




If a space AB be contained under a residual line AC ($CE - AE$) and a binomial CB , whose names CD, DB are commensurable to the names CE, AE , of the residual line, and in the same proportion ($CE : AE :: CD : DB$) then the right line F which contains in power that space AB , is irrational.

Let G be ρ . and make the rectangle $CH = Gq$; a then shall BH ($HI - IB$) be a residual line, a 113. 10. and HI $a \sqsupset CD$ $b \sqsupset CE$. a and BI $\sqsupset DB$ $a b$ hyp. and $HI.BI :: CD.DB$ $b :: CE.EA$. therefore by inversion $HI.CE :: BI.EA$. c therefore $BH.AC :: c$ 19. 5. $HI.CE :: BI.EA$. c therefore $BH.AC :: HI.CE :: d$ 12. 10. $BI.EA$. wherefore since d $HI \sqsupset CE$, e thence e 10. 10. $BH \sqsupset AC$. f therefore the rectangle $HC \sqsupset BA$. f 1. 6. and But HC (Gq) b is ρv , g therefore BA (Fq) is $\rho v : g$ sch. 12. and consequently F is ρ . Which was to be dem. 10.

Coroll.

Hereby it appears that a rational superficies may be contained under two irrational right lines.

PROP. CXVI.




Of a medial line AB are produced infinite irrational lines $BE, EF, \&c.$ whereof none is of the same kind with any of the precedent.

Let AC be propounded ρ . and AD a rectangle cqu .

contained under AC, AB. *a* therefore AD is $\sqrt{p}$.
a lem. 38. Take $BE = \sqrt{AD}$. *b* then BE is $\sqrt{p}$, and the
 10. same with none of the former. For no square of
b 11. 10. any of the former being applied to $\sqrt{p}$, makes the
 breadth medial. Let the rectangle DE be finish-
 ed, *a* then DE shall be $\sqrt{p}$. and *b* consequently EF
 ($\sqrt{DE}$) shall be $\sqrt{p}$, and not the same with any
 of the former, for no square of the former be-
 ing applied to $\sqrt{p}$, makes the breadth BE. there-
 fore, &c.

P R O P. CXVII.



a 47. 1.
b cor. 24. 8.
c 9. 10.

B C

Let it be required to shew that
 in square figures BD. the diameter
 AC is incommensurable in length
 to the side AB.

For ACq. ABq *a* :: 2. 1 *b* ::
 not Q. Q. *c* therefore AC $\nparallel$

AB. Which was to be dem. This Theorem was
 of great note with the ancient Philosophers;
 so that he that understood it not was esteemed
 by Plato undeserving the name of a Man, but
 rather to be reckoned among Brutes.

The End of the tenth Book.

THE

THE

EUC

A



A



also n
 CD,


THE ELEVENTH BOOK OF EUCLIDE'S ELEMENTS.

Definitions.


A

Solid is that which hath length, breadth, and thickness.

II. The term, or extreme of a solid is a Superficies.




III. A right line AB is perpendicular to a Plane CD, when it makes right angles ABD, ABE, ABF, with all the right lines BD, BE, BF, that touch it, and are drawn in the said Plane.




IV. A Plane AB, is perpendicular to a Plane CD, when the right lines FG, HK, drawn in one Plane AB to the line of common section of the two Planes EB, and making right angles therewith, do

also make right angles with the other Plane CD.

V. The



V. The inclination of a right line AB to a Plane CD, is, when a perpendicular AE is drawn from A the highest point of that line AB to the plane CD, and another line EB drawn from the point E, which the perpendicular AE makes in the Plane CD, to the end B of the said line AB which is in the same Plane, whereby the angle is acute ABE which is contained under the inclining line AB, and the line drawn in the plane EB.



VI. The inclination of a Plane AB, to a Plane CD, is an acute angle FGH contained under the right lines FH, GH, which being drawn in either of the Planes AB, CD to the same point H of the common section BE, make right angles FHB, GHB, with the common section BE.

VII. Planes are said to be inclined to other Planes in the same manner, when the said angles of inclination are equal one to another.

VIII. Parallel Planes are those which being prolonged never meet.

IX. Like solid figures are such as are contained under like Planes equal in number.

X. Equal and like solid figures are such as are contained under like Planes equal both in multitude and magnitude.

XI. A solid angle is the inclination of more than two right lines which touch one another, and are not in the same superficies.

A solid more than superficies

XII. A solid under (which is) thered to

XIII. A solid under pl equal, li parallel

XIV. the diam semicirc to the f moved.

Hence to the f themfel

XV. line, ab

XVI. point w

XVI. line dra

either

XVI. side of

that co

triangl

the pl

fixed

tains t

gled C

Cone

XII. abou

Or thus;

A solid angle is that which is contained under more than two plane angles not being in the same superficies, but consisting all at one point.

XII. A Pyramide is a solid figure comprehended under divers planes set upon one plane, (which is the base of the Pyramide) and gathered together to one point.

XIII. A Prisme is a solid figure contained under planes, whereof the two opposite are equal, like, and parallel; but the others are parallelograms.

XIV. A Sphere is a solid figure made when the diameter of a circle abiding unmoved, the semicircle is turned round about, till it return to the same place from whence it began to be moved.

Coroll.

Hence, all the rayes drawn from the center to the superficies of a sphere, are equal amongst themselves.

XV. The Axis of a sphere, is that fixed right line, about which the semicircle is moved.

XVI. The Center of a sphere, is the same point with that of the semicircle.

XVII. The Diameter of a sphere, is a right line drawn thro' the center, and terminated on either side in the superficies of the sphere.

XVIII. A Cone is a figure made, when one side of a rectangled triangle (*viz.* one of those that contain the right angle) remaining fixed, the triangle is turned round about till it return to the place from whence it first moved. And if the fixed right line be equal to the other which contains the right angle, then the Cone is a rectangled Cone: but if it be less, it is an obtuse-angled Cone; if greater an acute-angled Cone.

XIX. The Axis of a Cone is that fix'd line about which the triangle is moved.

XX. The

XX. The Base of a Cone is the circle, which is described by the right line moved about.

XXI. A Cylinder is a figure made by the moving round of a right-angled parallelogram, one of the sides thereof, (namely which contain the right angle) abiding fix'd, till the parallelogram be turned about to the same place, where it began to move.

XXII. The Axis of a Cylinder is that quiescent right line, about which the parallelogr. is turned.

XXIII. And the Bases of a Cylinder are the circles which are described by the two opposite sides in their motion.

XXIV. Like Cones and Cylinders, are they, both whose Axes and Diameters of their Bases are proportional.

XXV. A Cube is a solid figure contained under six equal squares.

XXVI. A Tetraedron is a solid figure contained under four equal and equilateral triangles.

XXVII. An Octaedron is a solid figure contained under eight equal and equilateral triangles.

XXVIII. A Dodecaedron is a solid figure contained under twelve equal, equilateral and equiangular Pentagones.

XXIX. An Icosaedron is a solid figure contained under twenty equal and equilateral triangles.

XXX. A Parallelepipedon is a solid figure contained under six quadrilateral figures, whereof those which are opposite are parallel.

XXXI. A solid figure is said to be inscribed in a solid figure, when all the angles of the figure inscribed are comprehended either within the angles, or in the sides, or in the planes of the figure wherein it is inscribed.

XXXII. Likewise a solid figure is then said to be circumscribed about a solid figure, when either the angles, or sides, or planes of the circumscribed figure touch all the angles of the figure which it contains,

PROF.



rectly to
strait from
one com



For
to be in
another
in a pla
a Which
is in o
the rig
lines A
to be de



in the
in the
theref
superf


PROP. I.



One part *AC* of a right line cannot be in a plane superficies, and another part *CB* elevated upward.

Produce *AC* in the plane directly to *F*. If you conceive *CB* to be drawn straight from *AC*, then two right lines *AB*, *AF*, have one common segment *AC*. *a Which is impossible.* a 10. ax. 1.


PROP. II.



If two right lines *AB*, *CD*, cut one another, they are in the same plane: And every triangle *DEB* is in one and the same plane.

For imagine *EFG*, part of the triangle *DEB*, to be in one plane, and the part *FDGB* to be in another, then *EF* part of the right line *ED* is in a plane; and the other part elevated upwards. *a Which is absurd.* Therefore the triangle *EDB* a 1. 11. is in one and the same plane; and so also are the right lines *ED*, *EB*; *a wherefore the whole lines AB*, *DC*, are in one plane. *Which was to be demonstrated.*


PROP. III.



If two planes *AB*, *CD*, cut one the other, their common section *EF* is a right line.

If *EF* the common section be not a right line, *a* then in the plane *AB* draw the right line *EGF*, *a* and *a* 1. post. 1. in the plane *CD* draw the right line *EHF*. therefore two right lines *EGF*, *EHF*, include a superficies. *b Which is absurd,* b 14. ax. 1.

PROP.




If a right line EF be at right angles erected upon two lines AB, CD , cutting one the other, at the common section E ; it shall also be at right angles to the plane $ACBD$ drawn by the said lines.

- Take EA, EC, EB, ED , equal one to the other, and join the right lines AC, CB, BD, AD , draw any right line GH thro' E , and join FA, FC, FD, FB, FG, FH . Because AE is $a = EB$, and DE $a = EC$, and the angle AED $b = CEB$. therefore AD is $= CB$, c and likewise $AC = DB$. d therefore AD is parallel to CB , d and AC to DB . e wherefore the angle $GAE = EBH$, and the angle $AGE = EHB$. But also $AE = EB$. f therefore $GE = EH$, g and $AG = BH$. whence by reason of the right angles, by the hyp. and so equal, at E , h the bases FA, FC, FB, FD , are equal. Therefore the triangles ADF, FBC , are equilateral one to another. k and thence the angle $DAF = BCF$. Therefore in the triangles AGF, FBH , the sides FG, FH l are equal; and so by consequence the triangle FEG and FEH are mutually equilateral. m therefore the angles FEG, FEH are equal, and n so right angles. In like manner, FE makes right angles with all the lines drawn thro' E in the plane $ADBC$, o and is therefore perpendicular to the said plane.

PROP.


PROP. V.



If a right line AB be erected perpendicular to three right lines AC , AD , touching one the other at the common section, those three lines are in the same plane.

For AC , AD , a are in one a 2. 11. plane FC ; a and AD , AE , are in one plane BE . which if you conceive to be several planes, then let their intersection b be b 3. 11. the right line AG ; therefore because BA by the Hypoth. is perpendicular to the right lines AC , AD . c and so to the plane FC , d it is also c 4. 11. perpendicular to the right line AG . therefore d 3. def. 11. (since a that AB is in the same plane with AC , AE) the angles BAG , BAE , are right angles, and consequently equal, the part and the whole. Which is absurd.

PROP. VI.



If two right lines AB , DC , be erected perpendicular to one and the same plane EF ; those right lines AB , DC are parallel one to the other.

Draw AD , whereunto let $DG = AB$ be perpendicular in the plane EF , and join BD , BG , AG . Being in the triangles BAD , ADG , the angles BAD , ADG a a hyp. are right angles, and AB $b = DG$, and AD is b constr. common, c therefore BD is $= AG$. whence in the c 4. 1. triangles AGB , BGD , equilateral one to the other, the angle BAG is $d = BDG$; of which d 8. 1. being BAG is a right angle, BDG shall be so also. but the angle GDC is supposed right, therefore the right line GD is perpendicular to the three lines DA , DB , CD . e which are therefore e 5. 11. in the same plane f wherein AB is. Wherefore f 2. 11. since AB and CD are in the same plane, and the internal angles BAD , CDA , are right angles, g g 28. 1. AB and CD shall be parallels. Which was to be dem.

S

PROP.

PROP. VII.




If there be two parallel right lines AB , CD , and any points E , F , be taken in both of them, the line EF which is joined at these points, is in the same plane with the parallels $ABCD$.

Let the plane in which AB , CD are, be cut by another plane at the points E , F . then if EF is not in the plane $ABCD$, it shall not be the common section. Therefore let EGF be the common section; which *a* then is a right line. therefore two right lines EF , EGF , include a superficies. *b* Which is absurd.

a 3. II.

b 14. ax. I.

PROP. VIII.



If there be two parallel right lines AB , CD , whereof one AB is perpendicular to a plane EF . then the other CD shall be perpendicular to the same plane EF .

The preparation and demonstration of the sixth of this Book being transfer'd hither; the angles GDA , and GDB are right angles: *a* therefore GD is perpendicular to the plane, wherein are AD , DB (*b* in which also AB , CD are,) *c* therefore GD is perpendicular to CD . but the angle CDA is also *d* a right angle. *e* therefore CD is perpendicular to the plane EF . Which was to be dem.

a 4. II.

b 7. II.


c 3. def. II.

d 29. I.

e 4. II.

PROP.

P R O P. IX.



Right lines (AB, CD) which are parallel to the same right line EF , but not in the same plane with it, are also parallel one to the other.

In the plane of the parallels AB, EF , draw HG perpendicular to EF ; also in the plane of the parallels EF, CD , draw IG perpendicular to EF . *a* therefore EG is perpendicular to the plane *a* 4. 11. wherein HG, GI are; and AH, CI are perpen- *b* 8. 11. dicular to the same plane. *c* therefore EH and *c* 6. 11. CI are parallels. Which was to be dem.


P R O P. X.



If two right lines AB, AC , touching one another be parallel to two other right lines ED, DF , touching one another, and not being in the same plane, those right lines contain equal angles, BAC, EDF .

Let AB, AC, DE, DF , be equal one to the other, and draw AB, BC, EF, BE, CF . Being AB, DE , *a* are parallels, and equal, *b* also BE, AD , are paral- *a* hyp. and *constr.* els and equal. In like manner CF, AD , are pa- *b* 33. 1. rallels and equal; *c* therefore also BE, FC , are *c* 2. ax. 1. parallels and equal. *d* Therefore BC, EF are e- *d* 33. 1. qual. Wherefore since the triangles BAC, EDF , and *e* 8. 1. are of equal sides one to the other, the angles BAC, EDF *e* shall be equal. Which was to be dem.

P R O P. XI.



From a point given on high A , to draw a right line AI perpendicular to a plane below BC .

In the plane BC draw any line S

- a 12. I. line DE; to which from the point A *a* draw the
 b 11. I. perpendicular AF, and *b* likewise FH in the
 plane BC cutting the said line DE at F; *a* then
 let fall AI perpendicular to FH. Which AI
 shall be perpendicular to the plane BC.
- c 31. I. For thro' I *c* let KIL be drawn parallel to DE.
 d constr. Because DE *d* is perpendicular to AF, and FH, *e*
 e 4. II. therefore DE shall be perpendicular to the plane
 f 8. 21. IFA. and so also KL *f* is perpendicular to the
 g 3. def. 11. same plane. *g* therefore the angle KIA is a right
 h constr. angle. but the angle AIF is also *b* a right angle.
 l 4. II. therefore AI is perpendicular to the plane BC.
Which was to be done.

PROP. XII.



a 11. II.

b 31. I.


c 8. II.

In a plane given BC, at a point
 given therein A, to erect a per-
 pendicular line AF.

From some point without the
 plane D, *a* draw DE perpendi-
 cular to the said plane BC, and joining the
 points A, E, by a line AE, *b* draw AF parallel
 to DE. *c* it is apparent that AF is perpendicu-
 lar to the plane BC. *Which was to be done.*

This and the preceding Problem are practi-
 cally performed by applying two squares to the
 point given; as appears by 4. II.

PROP. XIII.




a 6. II.

As a point given C in a
 plane given AB, two right lines
 CD, CE, cannot be erected per-
 pendicular on the same side.

For both CD, and CE, *a*
 should then be perpendicular to the plane AB,
 and consequently parallels; which is repugnant
 to the definition of parallel lines.

PROP.


PROP. XIV.



Planes CD, FE, to which the same right line AB is perpendicular, are parallel.

If you deny this; then let the planes CD, FE, meet, so that their common section be the right line GH, in which take any point I, draw to it the right lines IA, IB, in the said planes, whereby in the triangle IAB, two angles IAB, IBA *a* *hyp. and 3. def. 11.* are right angles. *b* Which is absurd. *b* 17.1.


PROP. XV.



If two right lines AB, AC, touching one the other, be parallel to two other right lines DE, DF, touching one the other, and not being in the same plane with them, the planes BAC, EDF, drawn by those right lines are parallel one to the other.

From A *a* draw AG perpendicular to the plane EF. *b* and let GH, GI be parallel to DE, *b* 11. 11. DF. *c* these also shall be parallel to AB, AC. *c* 31. 1. Therefore since the angles IGA, HGA, *d* are right angles, also CAG, BAG, *e* shall be right angles. *f* therefore GA is perpendicular to the plane BC; but the same is perpendicular to the plane EF. *g* therefore the planes BC, EF, are parallel. *h* 14. 11. Which was to be dem. *g* 30. 1. *d* 3. def. 11. *e* 29. 1. *f* 4. 11. *h* 14. 11.

PROP. XVI.




If two parallel planes *AD*, *CD*, be cut by some other plane *HEIGF*, their common sections *EH*, *GF* are parallel one to the other.

For if they be conceived to be otherwise; being in the same plane that cuts them, they will meet some-

where, if produced; suppose in *I*; wherefore since the whole lines *HEI*, *FGI* are in the planes *AD*, *CD*, being produced, the planes also shall meet. *contrary to the Hyp.*

a 1. 11.

PROP. XVII.



If two right lines *ALB*, *CMD*, be cut by parallel planes *EF*, *GH*, *IK*; they shall be cut proportionally, (*AL*. *LB* :: *CM*. *MD*).

Let the right lines *AC*, *BD*, be drawn in the planes *EF*, *IK*; as also *AD* passing thro' the plane *GH* in the point *N*. and join *NL*,


LM the planes of the triangles *ADC*, *ADB*, make the sections *DB*, *LN*, and *AC*, *NM* & parallels. Therefore *AL*. *LB* :: *AN*. *ND* :: *CM*. *MD*. Which was to be dem.

a 16. 11.

b 2. 6,

PROP.


PROP. XVIII.



If a right line AB be perpendicular to some plane CD , all the planes extended by that right line AB (EF , &c.) shall be perpendicular to the same plane CD .

Let there be some plane EF drawn by AB , making the section EG ^{a 31. 1.} with the plane CD ; from some point whereof ^{b 8. 11.} H , draw HI parallel to AB in the plane EF ; ^{c 4. def. 11.} then shall HI be perpendicular to the plane CD , and so likewise any other lines, that are perpendicular to EG , therefore the plane EF is perpendicular to the plane CD ; and by the same reason any other planes drawn by AB shall be perpendicular to EF . Which was to be dem.

PROP. XIX.



If two planes AB , CD , cutting one the other, be perpendicular to some plane GH . their line of common section EF shall be perpendicular to the same plane (GH).

Because the planes AB , CD , are taken perpendicular to the plane GH , it appears by 4. def. 11. ^{a 13. 11.} that out of the point F there may be drawn in both planes AB , CD , a perpendicular to the plane GH . which shall be a but one; and therefore the common section of the said planes. Which was to be dem.

PROP. XX.




If a solid angle $ABCD$ be contained under three plane angles, BAD , DAC , BAC , any two of them howsoever taken are greater than the third.

If the three angles are equal, the assertion is evident; if unequal, then let the greatest be BAC ; from whence a take away $BAE = BAD$, and make $AD = AE$; and also draw BEC , BD , DC .

Because the side BA is common, and $AD = AE$; and the angle $BAE = BAD$. c thence is c 4. I. $BE = BD$. but $BD + DC$ is $d < BC$. e therefore d 20. I. $DC < EC$. Wherefore since $AD = AE$, and the e ax. I. side AC is common, and $DC < EC$. f the angle f 25. I. CAD shall be $< EAC$. g therefore the angle g 4. ax. I. $BAD + CAD < BAC$. Which was to be dem.

PROP. XXI.



Every solid angle A is contained under less angles than four plane right angles.

For let a plane any-wise cutting the sides of the solid angle A make a many-sided figure $BCDE$, and as many triangles ABC , ACD , ADE , AEB . I denote all the angles of the polygone by X ; and I term the sum of the angle at the bases of the a 32. I. Σ triangles Y . wherefore $X + 4$ right angles $a =$ a 31. I. $Y + A$. but being that, (of the angles at B) b 20. II. the angle $ABE + ABC$ is $< CBE$, and the same c 5. ax. I. is true also of the angles at C , at D , and at E , c it is manifest that Y is $< X$. and consequently A shall be < 4 right angles. Which was to be dem.

PROP.

PROP. XXII.




If there be three plane angles A, B, HCI , whereof two howsoever taken are greater than the third, and the right lines which contain them be equal AD, AE, FB , &c. then of the right lines DE, FG, HI , coupling those equal right lines together, it is possible to make a triangle.

A triangle may be made of them, if any two be greater than the third: but they are so. For make the angle $HCK = B$, and $CK = b$ 23. 1. CH , and draw HK, IK c thence $HK = FG$. and c 4. 1. because the angle $KCI = A$. e therefore $KI = d$ hyp. DE . but $KI = f = HI + KH$ (FG .) therefore $DE = 24. 1. = HI + FG$. By the like argument any two f 20. 1. may be proved greater than the third; and consequently a it is possible to make a triangle of them. Which was to be dem.

PROP.

PROP. XXIII.




To make a solid angle $MHIK$ of three plane angles A, B, C , whereof two howsoever taken are greater than the third. * But it is necessary that those three angles be less than four right angles.

Make AD, AE, BE, BF, CF, CG , equal one to the other; and of the subtended lines DE, EF, FG (that is, of the equal lines HI, IK, KH) a
 a 22. 11. make the triangle HKI ; about which b describe
 and 22. 1. the circle $LHKI$. * But because AD is $\perp HL$.
 b 5. 4. * See Cla- c let ADq be $= HLq + LMq$. d and let LM
 vna. be perpendicular to the plane of the circle HKI .
 c sch. 47. 1. and draw HM, KM, IM . wherefore since the
 d 12. 11. angle HLM is a right angle, f thence is MHq
 e 3. def. 11. $= HLq + LMq$ $g = ADq$. therefore $MH =$
 f 47. 1. AD . By the same reason MK, MI, AD (that is,
 g constr. $AE, EB, \&c.$) are equal; therefore since $HM =$
 h constr. AD , and $MI = AE$, and $DE = HI$. k the angle
 k 8. 1. A shall be $= HMI$, k as likewise the angle IMK
 $= B$, k and the angle $HMK = C$. wherefore a
 solid angle is made at M of the three given
 plane angles. Which was to be done. AD is assu-
 med to be $\perp HL$. But this is manifest. For if
 1 constr. $\&$ AD be $=$ or $\perp HL$, then is the angle $A =$
 8. 1. m or $\perp HLI$. In like manner shall B be $=$ or $\perp$
 m 22. 1. HLK , and $C =$ or $\perp KLI$. wherefore $A + B + C$
 * 4. cor. 13. * shall either equal or exceed four right angles.
 1, contrary to the Hypoth. therefore rather let AD
 be $\perp HL$. Which was to be dem.

PROP.

PROP. XXIV.



If a solid AB be contained under parallel planes, the opposite planes thereof ($AG, DB, \&c.$) are like and equal parallelograms.

The plane AC cutting the parallel planes AG, DB , a makes the sections AH, DC , parallel. and by the same reason AD, HC are parallel. Therefore $ADCH$ is a parallelogram. By the like argument the other planes of the parallelepipedon are b parallelograms. wherefore b 35. def. 1. being AF is parallel to HG , and AD to HC , c 10. 11. the angle FAD shall be $= CGH$. therefore be- d 34. 2. cause AF $d = HG$, and AD $d = HC$, and so e 7. 5. $AF. AD :: HG. HC$, the triangles FAD, GHC , g 6. 6. are like and h equal; and consequently the h 4. 1. parallelograms AE, HB are like and k equal. and k 6. ax. 1. the same may be shewn of the other opposite planes. therefore, $\&c.$

PROP. XXV.



If a solid Parallelepipedon $ABCD$ be cut by a plane EF , parallel to the opposite planes AD, BC ; then as the base AH is to the base BH , so shall solid AHD be to solid BHC .


Conceive the Parallelepipedon to be extended on either side, and take $AI = AE$, and $BK = EB$, a 36. 1. $\&c.$ and put the plane IQ, KP , parallel to the planes $1.$ def. 6. AD, BC ; then the pgrs. IM, AH , and a DL, DG , b 24. 11. and $IQ, AD, EF, \&c.$ are a like and equal. c 10. def. wherefore the Parallelepipedon AQ is $= AF$; and $11.$ by

by the same reason the Parallelepipedon BP = BF. therefore the solids IF, EP are as multiple of the solids AF, EC, as the bases IH, KH, are of the bases AH, BH. And if the basis IH be d 24. II. $\square, =, \square$ KH, & likewise shall the solid IE be and 9. def. $\square, =, \square$ EP. e consequently AH. BH :: AF. II. EC. Which was to be dem.
e 6. def. 5. The same may be accommodated to all sorts of Prisms, whence

Coroll.

If any Prism whatsoever be cut by a plane parallel to the opposite planes, the section shall be a figure equal and like to the opposite planes.

P R O P. XXVI.




Upon a right line given AB, and at a point given in it A, to make a solid angle AHIL equal to a solid angle given CDEF.

2 II. II. From some point F draw FG perpendicular to the plane DCE, and draw the right lines DF, FE, EG, GD, CG. Make AH = CD, and the angles HAI = DCE, and AI = CE; and in the plane HAI make the angle HAK = DCG, and AK = CG. then erect KL perpendicular to the plane HAI, and let KL be = GF. and draw AL: then AHIL shall be a solid angle equal to that given CDEF. For the construction of this does wholly resemble the framing of that, as may easily appear to any that examine it.

PROP.

PROP. XXVII.




Upon a right line given AB to describe a parallelepipedon AK like, and in like manner situate, with a solid parallelepipedon given CD .

Of the plane angles BAH , HAI , BAI , which are equal to FCE , ECG , FCG , a make the solid a 26. 11. angle A equal to the solid angle C . also b make b 12. 6. $FC.CE :: BA.AH$. b and $CE.CG :: AH.AI$ (c 22. 5. whence of equality $FC.CG :: BA.AI$) and finish the parallelepipedon AK , which shall be like to that which is given.

For by the construction, the Parallelogram d d 1. def. 6. BH is like FE , and d HI to EG , and d BI to FG , and e so the opposites of these to the opposites e 14. 11. of them: therefore the six planes of the solid AK are like to the six planes of the solid CD , f and consequently AK , CD , are like solids. f 9. def. 11. Which was to be dem.

PROP. XXVIII.




If a solid parallelepipedon AB be cut by a plane $FGCD$ drawn by the diagonal lines DF , CG , of the opposite planes AE , HB , that solid AB shall be equally bisected by the plane $FGCD$.

For because DC , FG , are a equal and parallels, a 24. 11. b the plane $FGCD$ is a Pgr. and being a the Pgrs. b 34. 1. AE , HB , are equal and like, b also the triangles AFD , HGC , CGB , DFE are equal and like. But the Pgrs. AC , AG , are equal and like to FB and FD . therefore all the planes of the prisme $FGC-DAH$ are equal and like to all the planes of the prisme $FGCDEB$, and c consequently this prisme c 9. def. 11. is equal to that. Which was to be dem.

PROP.

The eleventh Book of
PROP. XXIX.



Solid Parallelepipedons $AGHEFBCD$, $AGHEMLKI$, being constituted upon the same base $AGHE$, * i. e. be- and * in the same height, whose insisting lines AF , AM , are placed in the same right lines AG , FL , are parallel equal one to the other.

For a if from the equal prismes $AFMEDI$, $HEFL-GBLHCK$, the common prisme $NBMPCI$ be KD , and so taken away, and the solid $AGNEHR$ be added, understand the Parallelepipedon $AGHEFBCD$ shall be = it in the $AGHEMLKI$. Which was to be dem. following.

a 10. def.


ii. and 35.

i.

b 3. and i.

ax. i.

PROP. XXX.




Solid parallelepipedons $ADBCHEFG$, $ADCBIMLK$ being constituted upon the same base $ADBC$, and

in the same height, whose insisting lines AH , AI are not placed in the same right lines, are equal one to the other.

For produce the right lines HEO , GFN , and LMO , KIP ; and draw AP , DO , BQ , CN . *a* then *a* 34. 1. shall DC , AB , HG , EF , PQ , ON be as well equal and parallel one to the other as AD , HE , GF , BC , KL , IM , QN , PO . *b* wherefore the parallelepipedon $ADCBPONQ$ shall be equal to either parallelepipedon $ADCBHEFG$, $ADCBIMLK$; and *c* consequently these two are equal *c* 1. 28. 1. one to the other. Which was to be dem.

P R O P. XXXI.



* by height
understand

Solid parallelepipeds, $ALEKGMBI$, $CPWOH2$ the perpendicular, being constituted upon equal bases $ALEK$, $CPWO$, and * in the same height are equal, one to the other. drawn from


First, let the parallelepipeds AB , CD , have the plane of the sides perpendicular to the bases, and at the the base to side CP being produced, *a* make the parallelo- the opposite gram $PRTS$ equal and like to the parallelogram plane.

$KELA$. *b* and so the parallelepipedon $PRTSQ$ - *a* 18. 6. VYX equal and like to the parallelepipedon AB . *b* 27. 11. Produce OE , ND , $WPZDQF$, ERB , SV , and *10. def.* TSZ , YXF , and draw ES , BZ , ZF . *11.*

The planes $OEAN$, $CRVHZTYF$ *c* are paral- *c* 30. def. lels one to the other; *d* and the pgrs. $ALEK$, *11.* $CDWO$, $PRTS$, $PRBZ$ are equal. Therefore since *d* hyp. and the parallelepipedon $CD.PV$ $\Delta ::$ pgr. CO ($PRBZ$) *35. 1.* P $c ::$ parallelepipedon $PRBZQVYF$. PV $\Delta ::$ the *e* 25. 11. paral-

- f 9. 5. parallelepipedon CDf shall be = PRBZVQ. Fig.
 g 29. II. PRVQSTYX $b = AB$. Which was to be dem.
 h constr. But if the parallelepipedons AB, CD, have
 sides oblique to the bases, then on the same ba-
 ses and in the same heighth place parallelepipe-
 k 29. II. dons whose sides are perpendicular to the bases
 & They shall be equal to one another, and those
 m 1. ax. I. that are oblique; m whence also the oblique
 parallelepipedons AB, CD are equal. Which
 was to be dem.


PROP. XXXII.



Solid parallelepipedons $ABCD$, $EFGL$, of the same
 heighth, are one to the other, as their bases AB , EF .

- a 45. I. Produce EHI , a and make the pgr. $FI = AB$
 b 31. I. and b compleat the parallelepipedon $FINM$. It
 c 31. II. is clear that the parallelepipedon $FINM$ (of
 d 25. II. $ABCD$.) $EFGL$ $d :: FI$ (AB .) EF . Which was
 to be dem.

PROP. XXXIII.



Like solid parallelepipeds, $ABCD$, $EFCH$, are
 in triplicated proportion one to the other, of that in which
 their homologous sides are in like proportion AI , ER
 are.

Produce the right line
 AIL , DIO , BIN , and
 make IL , IO , IN , equal


a 3. I.

to EK, KH, KF, *b* and so the parallelepipedon *b* 27. 11.
 IXMT equal and like to the parallelepipedon *c* 31. 1.
 EFGH. *c* Let the paralleps. IXPB, DLYQ be fi- *d* hyp.
 nished. *d* Then shall be AL. IL (EK) :: DL. IO *e* 1. 6.
 (HK) :: BI. IN. KF. *e* that is, the Pgr. AD. DL *f* 32. 11.
 :: DL. IX :: BO. IT. *f* i. e. the parallepp. ABCD. *g* constr.
 DLQY :: DLQY. IXPB :: IXPB. IXMT. (*g* EFGH.) *h* 10. def. 5.
b therefore the proportion of ABCD to EFGH *k* 1. 6.
 is triple of the proportion of ABCD to DLQY,
k, or of AI to EK. Which was to be dem.

Coroll.

Hence it appears, that if four right lines be continually proportional, as the first is to the fourth, so is a parallelepipedon described on the first to a parallelepipedon described on the second, being like and in like manner described.

PROP. XXXIV.



In equal solid parallelepipedons ADCB, EHGF, the bases and altitudes are reciprocal (AD. EH :: EG. AC.) And solid parallelepipedons, ADCB, EHGF, whose bases and altitudes are reciprocal,

are equal.

First, let the sides CB, GE be perpendicular to *a* 3. 1.
 the bases; then if the altitudes of the solids are *b* 31. 1.
 equal, the bases also shall be equal, and the thing *c* 32. 11.
 is clear. But if the altitudes are unequal, from *d* 17. 5.
 the greater EG *a* take EI = AC, and at *I* *b* draw *e* 1. 6.
 the plane IK parallel to the base EH. then *f* constr.
 1. Hyp. AD. EH *c* :: parallepp. ADCB. EHIK *g* 11. 5.
d :: parallepp. EHGF. EHIK *c* :: GL. IL *e* :: GE. *h* 32. 11. 1
 IE. (*f* AC.) *g* it is plain therefore that AD. EH *k* hyp.
 :: GE. AC. Which was to be dem. 1 1. 6.

2. Hyp. ADCB. EHIK *b* :: AD. EH *k* :: EG. EI *m* 32. 11.
 :: GL. IL *m* :: parallepp. EHGF. EHIK. *n* where- *n* 9. 5.

T

fore

fore the parallelepipedon $ADCB = EHGF$.
Which was to be dem.


Moreover, let the sides be oblique to the bases, and erect right parallelepipedons upon the same bases in the same altitude; the oblique parallelepipedons shall be equal to them. Wherefore since by the first part, the bases and altitudes of those be reciprocal, the bases and altitudes of these also shall be reciprocal. Which was to be dem.

Coroll:

All that hath been demonstrated of parallelepipedons in the 29, 30, 31, 32, 33, 34 Prop. does also agree in triangular prismes, which are half parallelepipedons, as appears by Prop. 28. Therefore,

1. Triangular prismes are of equal height with their bases.
2. If they have the same or equal bases and the same altitude, they are equal.
3. If they be like, their proportion is triple to that of their sides of like proportion.
4. If they be equal, their bases and altitudes are reciprocal; and if their bases and altitudes be reciprocal, they are also equal.

P R O P: XXXV.



If there be two plane angles BAC , EDF , equal, and from the points of those angles two right lines AG , DH be elevated on high,

containing equal angles with the lines first given, each to his correspondent angle (the angle $GAB = HDE$, and $GAC = HDF$.) and if in those elevated lines AG , DH , some points be taken, G , H ; and from these points perpendicular lines GI , HK , drawn to the planes BAC , EDF , in which the angles first given are, and right lines

lines *AI, DK*, be drawn to the angles first given from the points *I, K*, which were made by the perpendiculars in the planes; those right lines with the elevated lines *AG, DH* shall contain equal angles *GAM, HDK*.

Make *DH, AL*, equal; and *GI, LM* parallels, and *MC* to *AC*, *MB* to *AB*, *KF* to *DF*, *KE* to *DE* perpendicular; and draw the right lines *BC, LB, LC*, and *EF, HF, HE*; *a* and *LM* is perpendicular to the plane *BAC*; *b* wherefore the angles *LMC, LMA, LMB*; and by the same reason the angles *HKF, HKD, HKE* are right angles. Therefore $ALq\ c = LMq + AMq\ c = LMq + CMq + ACq\ c = LCq + ACq$. *d* therefore the angle *ACL* is a right angle. Again $ALq\ e = LMq + MAq\ e = LMq + BMq + BAq\ e = BLq + BAq$. *d* therefore the angle *ABL* is also a right angle. By the like inference the angles *DFH, DEH* are right angles; *f* therefore *AB = DE*, *f* and *BL = EH*, *f* and *AC = DF*, and *CL = FH*. *g* wherefore also *BC = EF*; *g* and the angle *ABC = DEF*, *g* and the angle *ACB = DFE*. whence the other right angles *CBM, BCM*, are equal to the other *FEK, EFK*. *k* therefore *CM = FK*, *l* and so also *AM = DK*. therefore if from *LAq\ m = HDq* be taken away *AMq = DKq*, *n* there remains *LMq = HKq*. wherefore the triangles *LAM, HDK* are equilateral one to the other; therefore the angle *LAM = HDK*. Which was to be dem.

Coroll.

Therefore, if there be two plane angles equal, from whose points equal right lines be elevated on high, containing equal angles with the lines first given, each to each; perpendiculars drawn from the extreme points of those elevated lines to the planes of the angles first given, are equal one to the other; viz. *LM = HK*.

T 2

PROP.

P R O P. XXXVIII.




If a plane *AB* be perpendicular to a plane *AC*, and a perpendicular line *EF* be drawn from a point

E in one of the planes (*AB*) to the other plane *AC*, that perpendicular *EF* shall fall upon the common section of the planes *AD*.

If it be possible, let *F* fall without the intersection *AD*, and in the plane *AC* draw *FG* ^{a 12. 1.} perpendicular to *AD*, and join *EG*. The angle *FGE* ^{b 4. and 3.} is a right angle, and *EFG* is supposed to be such also; therefore two right angles are in ^{def. 11.} the triangle *EFG*. ^{c 17. 1.} Which is absurd.

P R O P. XXXIX.



If the sides (*AE*, *FC*, *AF*, *EC*, and *DH*, *GB*, *DG*, *HB*) of the opposite planes *AC*, *DB*, of a solid parallelepipedon *AB*, be divided into two equal parts and planes *ILQO*, *PKMR*, be drawn thro' their sections, the common section of the planes *ST*, and the diameter of the

solid parallelepipedon *AB* shall divide one the other into two equal parts. ^{a 34. 1.}

Draw the right lines *SA*, *SC*, *TD*, *TB*. Because ^{b 29. 1.} the sides *DO*, *OT* are equal to the sides *BQ*, *QT*. ^{c 4. 1.} and the alternate angles *TOD*, *TQB* equal ^{al-d sch. 15. 1.} so, ^e the base *DT*, *TB*, and the angles *DTO*, ^{e 34. 1.} *BTQ* are equal. ^d therefore *DTB* is a right line, ^{f 9. 11. &} and so in like manner is *ASC*. Moreover ^{e as 1. ax.} well *AD* is parallel and equal to *FG* ^{e as FG to g 33. 1.} *CB*, and ^f thence *AD* is parallel and equal to *C*; ^{h 7. 11.}

T 3


and

- g 33. I. g and consequently AC to DB. *h* wherefore AB
 h 7. II. and ST are in the same plane ABCD. Therefore
 since the vertical angles AVS, BVT, and the al-
 k 7. ax. I. ternate angles ASV, BTV are equal; *k* and AS
 l 26. I. = BT; therefore shall AV be = BV. *l* and SV
 = VT. Which was to be dem.

Coroll.

Hence in every parallelepipedon all the diame-
 ters bisect one another in one point, V.

PROP. XL.



If two Prisms *ABCFED*, *GHMLIK*, be of equal
 altitude, whereof one hath its base *ABCF* a paralle-
 logram, and the other *GHM*, a triangle; and if the
 parallelogram *ABCF* be double to the triangle *GHM*;
 these Prisms *ABCFED*, *GHMLIK* are equal.

- a 31. II. For if the parallelepipedons *AN*, *GQ* be com-
 b 34. I. & pleted, *a* they shall be equal, because of the equal-
 7. ax. lity *b* of the bases *AC*, *GP*, and *c* of the altitudes.
 c hyp. *d* therefore also the prisms, *e* the halves thereof
 d 28. II. shall be equal. Which was to be dem.

Coroll.

From the preceding demonstrations, the dimension
 of triangular Prisms, and quadrangular, or paralle-
 lepidons, is learnt; viz. by multiplying the alti-
 tude into the base.

As if the altitude be 10 feet, and the base 100
 square feet (the base may be measured by sch. 35.
 I. or by 41. I.) then multiply 100 by 10, and

1000

1000 cubic feet shall be produced, for the solidity of the prism given.

For as a rectangle, so also is a right parallelepipedon produced of the altitude multiplied into the base. Therefore every parallelepipedon is produced of the altitude multiplied into the base, as appears by 31. of this Book.

Moreover, since the whole parallelepipedon is produced of the altitude drawn into the base, the half thereof (that is, a triangular Prism) shall be produced of the altitude drawn into half the base, namely the triangle.

An Advertisement.

Observe, That of those letters which denote a solid angle, the first is always at the point in which the angle is; but of those letters which denote a Pyramide, the last is at the supreme point thereof.

Ex. gr. The solid angle ABCD is at the point A; and the supreme point of the Pyramide BCDA is at the point A. and the base is the triangle BCD.


The End of the eleventh Book.

- g 33. I. g and consequently AC to DB. *h* wherefore AB
 h 7. II. and ST are in the same plane ABCD. Therefore
 since the vertical angles AVS, BVT, and the al-
 k 7. ax. I. ternate angles ASV, BTV are equal; *k* and AS
 l 26. I. = BT; therefore shall AV be = BV. *l* and SV
 = VT. Which was to be dem.

Coroll.

Hence in every parallelepipedon all the diame-
 ters bisect one another in one point, V.

PROP. XL.



If two Prisms $ABCFED$, $GHMLIK$, be of equal
 altitude, whereof one hath its base $ABCF$ a paralle-
 logram, and the other GHM , a triangle; and if the
 parallelogram $ABCF$ be double to the triangle GHM ;
 these Prisms $ABCFED$, $GHMLIK$ are equal.

- a 31. II. For if the parallelepipedons AN, GQ , be com-
 b 34. I. & pleted, *a* they shall be equal, because of the equa-
 7. ax. lity *b* of the bases AC, GP , and *c* of the altitudes.
 c hyp. *d* therefore also the prisms, *e* the halves thereof
 d 28. II. shall be equal. Which was to be dem.

Coroll.

e 7. ax. I. From the preceding demonstrations, the dimension
 Andr. of triangular Prisms, and quadrangular, or paralle-
 Tacq. lepipedons, is learnt; viz. by multiplying the alti-
 tude into the base.

As if the altitude be 10 feet, and the base 100
 square feet (the base may be measured by sch. 35.
 1. or by 41. 1.) then multiply 100 by 10, and

1000

1000 cubic feet shall be produced, for the solidity of the prism given.

For as a rectangle, so also is a right parallelepipedon produced of the altitude multiplied into the base. Therefore every parallelepipedon is produced of the altitude multiplied into the base, as appears by 31. of this Book.

Moreover, since the whole parallelepipedon is produced of the altitude drawn into the base, the half thereof (that is, a triangular Prism) shall be produced of the altitude drawn into half the base, namely the triangle.

An Advertisement.


Observe, That of those letters which denote a solid angle, the first is always at the point in which the angle is; but of those letters which denote a Pyramide, the last is at the supreme point thereof.

Ex. gr. The solid angle ABCD is at the point A; and the supreme point of the Pyramide BCDA is at the point A. and the base is the triangle BCD.

The End of the eleventh Book.

THE TWELFTH BOOK OF EUCLIDE'S ELEMENTS.

PROP. I.




L *Ike polygonous figures $ABCDE, FGHK$, described in circles ABD, FGI , are one to another, as the squares described of the diameters of the circles AL, FM .*
 Draw AC, BL, FH, GM . Because
 a 1. def. 6. a the angle $ABC = FGH$, a and $AB. BC :: FG.$
 b 6. 6. $GH.$ b therefore shall the angle ACB (c ALB)
 c 21. 3. be $= FHG$ (c FMG .) but the angles $ABL,$
 d 31. 3. FGM d are right and so equal; e therefore the
 e 32. 3. triangles ABL, FGM are equiangular. f where-
 f cor. 4. 6. fore $AB. FG :: AL. FM.$ g therefore $ABCDE.$
 g 22. 6. $FGHIK :: ALq. FMq.$

Coroll.

Hence (because $AB. FG :: AL. FM :: BC. GH,$
 $\&c.$) the contents of like polygonous figures de-
 h 1. 12. $\&c$ scribed in a circle are in b proportion as the
 12. 5. diameters,


PROP.

PROP. II.



Circles *ABT, EFN*, are in proportion one to another, as the squares of their diameters *AC, EG* are.

Suppose $ACq : EGq ::$ the circle *ABT*. *I*. I say then *I* is equal to the circle *EFN*.



For first, if it be possible, let *I* be less than the circle *EFN*, and let *K* be the excess or difference. Inscribe the square *EFGH* in the circle *EFN*, *a* it being the half of a circ^a *sch.7. 4.* cumscribed square, and so greater than the semi-circle. *b* Divide equally in two the arches *EF, FG, GH, HE*, and at the points of the divisions join the right lines *EL, LF, &c.* at *L* draw the tangent *PQ* (*c* which is parallel to *EF*) and produce *HEP, GFQ*. then is the triangle *ELF* *sch.27. 3.* *dd* the half of the parallelogram *EPQF*, and so greater than the half of the segment *ELF*; and in like sort the rest of those triangles exceed the halves of the rest of the segments. And if the arches *EL, LF, FM, &c.* be again bisected, and the right lines joined, the triangles will likewise exceed the half of the segments. Wherefore if the square *EFGH* be taken from the circle *EFN*, and the triangles from the other segments, and this be done continually, at length *e* there will remaine *I. 10.* some magnitude less than *K*. Let us have gone so far, namely to the segments *EL, LF, FM, &c.* taken

f hyp. and taken together less than K. Therefore I (f the
 3. ax. circle FENK) $\supset$ the polyg. ELFMNHO (the
 g 30.3. & circle FEN — the segm. EL + LF, &c.) In the
 i. post. 1. circle ABT g conceive a like polygonon AKBS.
 h i. 12. CTDV inscribed. therefore since AKBSCTDV.
 k hyp. ELFMGNHO $b :: ACq. EGq k ::$ the circle
 l 9. ax. 1. ABT. I. and the polyg. AKBSCTDV $l \supset$ the
 m 14. 5. circle ABT. the polyg. ELFMGNHO m shall
 be $\supset$ I. but before, I was $\supset$ ELFMGNHO.
 which is repugnant.


Again, if it be possible, let I be $\supset$ the circle
 n hyp. EFN. Therefore because ACq. EGq $n ::$ the circle
 ABT. I; and inversely I. the circle ABT. EGq.
 ACq. suppose I. the circle ABT $::$ the circle
 o 14. 5. EFN. K. o therefore the circle ABT $\supset$ K. p and
 p 11. 5. EGq. ACq $::$ the circle EFN. K. which is shewn
 to be repugnant.

Therefore it must be concluded, that I is $=$
 to the circle EFN. Which was to be dem.

Coroll.

Hence it follows, that as a circle is to a cir-
 cle, so is a polygonon described in one to a like
 polygonon described in the other.

PRO P. III.




Every Pyramide ABDC
 having a triangular base, may
 be divided into two Pyramides
 AEGH, HIKC, equal, and
 like one to the other, having
 bases triangular, and like to
 the whole ABDC; and into
 two equal Prisms, BFGEIH,
 EGDHIK; which two Prisms are greater than the
 half of the whole Pyramide ABDC.

Divide the sides of the pyramide into two
 parts at the points E, F, G, H, I, K, and join the right
 lines EF, FG, GE, EI, IF, FK, KG, GH, HE. Because
 the sides of the pyramide are proportionally
 cut,

cut, *a* thence HI, AB; and GF, AB; and IF, DC; *a* 2. 6.
 and HG, DC, &c. are parallels, and consequently
 HI, FG; and GH, FI are also parallels. therefore
 it is apparent that the triangles ABD, AEG, EBF,
 FDG, HIK, *b* are equiangular, and that the four *b* 29. 1.
 last are *c* equal: in like manner the triangles *c* 26. 1.
 ACB, AHE, EIB, HIC, FGK are equiangular; and
 the four last are equal one to the other. Also the
 triangles BFI, FDK, IKC, EGH; and lastly, the
 triangles AHG, GDK, HKC, EFI are like and
 equal. Moreover the triangles HIK to ADB, and
 EGH to BDC, and EFI to ADC, and FGK to
 ABC, *d* are parallel. From whence it evidently *d* 15. 11.
 follows, first that the pyramids AEGH, HIKC
 are equal, and *e* like to the whole ABDC, and to *e* 10. *def.*
 one another. Next, that the solids BFGEIH, 11.
 FGDIHK, are prisms, and that of equal height,
 as being placed between the parallel planes ABD,
 HIK, but the base BFGE is *f* double of the base *f* 2. *ax.* 1.
 FDG, wherefore the said prisms are equal; *g* 40. 11.
 whereof the one BFGEIH is greater than the
 pyramide BFEI, that is, then AEGH, the whole
 than its part; and consequently the two prisms
 are greater than the two pyramids, and so ex-
 ceed the half of the whole pyramide ABDC.
Which was to be dem.

PROP. IV.



If there be two pyramides ABCD, EFGH, of the
 same altitude, having triangular bases ABC, EFG;
 and

f hyp. and taken together less than K. Therefore I (f the
 3. ax. circle FENK) $\supset$ the polyg. ELFMGNHO (the
 g 30.2. & circle FEN — the segm. EL + LF, &c.) In the
 1. post. 1. circle ABT g conceive a like polygonon AKBS-
 h 1. 12. CTDV inscribed. therefore since AKBSCTDV,
 k hyp. ELFMGNHO $b :: ACq. EGq$ k $::$ the circle
 l 9. ax. 1. ABT. I. and the polyg. AKBSCTDV $l \supset$ the
 m 14. 5. circle ABT. the polyg. ELFMGNHO m shall
 be $\supset$ I. but before, I was $\supset$ ELFMGNHO,
 which is repugnant.


Again, if it be possible, let I be $\supset$ the circle
 n hyp. EFN. Therefore because ACq. EGq $n ::$ the circle
 ABT. I; and inversely I. the circle ABT. EGq.
 ACq. suppose I. the circle ABT $::$ the circle
 o 14. 5. EFN. K. o therefore the circle ABT $\supset$ K. p and
 p 11. 5. EGq. ACq $::$ the circle EFN. K. which is shewn
 to be repugnant.

Therefore it must be concluded, that I is $=$
 to the circle EFN. Which was to be dem.

Coroll.

Hence it follows, that as a circle is to a cir-
 cle, so is a polygon described in one to a like
 polygon described in the other.

PROP. III.




Every Pyramide ABDC
 having a triangular base, may
 be divided into two Pyramides
 AEGH, HIKC, equal, and
 like one to the other, having
 bases triangular, and like to
 the whole ABDC; and into
 two equal Prisms, BFGEIH,
 EGDHIK; which two Prisms are greater than the
 half of the whole Pyramide ABDC.

Divide the sides of the pyramide into two
 parts at the points E, F, G, H, I, K, and join the right
 lines EF, FG, GE, EI, IF, FK, KG, GH, HE. Because
 the sides of the pyramide are proportionally
 cut,

cut, *a* thence HI, AB; and GF, AB; and IF, DC; *a* 2. 6.
 and HG, DC, &c. are parallels, and consequently
 HI, FG; and GH, FI are also parallels. therefore
 it is apparent that the triangles ABD, AEG, EBF,
 FDG, HIK, *b* are equiangular, and that the four *b* 29. 1.
 last are *c* equal: in like manner the triangles *c* 26. 1.
 ACB, AHE, EIB, HIC, FGK are equiangular; and
 the four last are equal one to the other. Also the
 triangles BFI, FDK, IKC, EGH; and lastly, the
 triangles AHG, GDK, HKC, EFI are like and
 equal. Moreover the triangles HIK to ADB, and
 EGH to BDC, and EFI to ADC, and FGK to
 ABC, *d* are parallel. From whence it evidently *d* 15. 11.
 follows, first that the pyramids AEGH, HIKC
 are equal, and *e* like to the whole ABDC, and to *e* 10. *def.*
 one another. Next, that the solids BFGEIH, 11.
 FGDIHK, are prisms, and that of equal height,
 as being placed between the parallel planes ABD,
 HIK, but the base BFGE is *f* double of the base *f* 2. *ax.* 1.
 FDG, wherefore the said prisms are equal; *g* 40. 11.
 whereof the one BFGEIH is greater than the
 pyramide BFEI, that is, then AEGH, the whole
 than its part; and consequently the two prisms
 are greater than the two pyramids, and so ex-
 ceed the half of the whole pyramide ABDC.
Which was to be dem.

PROP. IV.



If there be two pyramids ABCD, EFGH, of the
 same altitude, having triangular bases ABC, EFG;
 and

and either of them be divided into two pyramides (AILM, MNOD; and EPRS, STVH) equal one to the other, and like to the whole, and into two equal prisms (IBKLMN, KLCNMO; and PFQ RST, QRGTSV;) and if in like manner either of those pyramides made by the former division be divided, and this be done continually; then as the base of one pyramide is to the base of the other pyramide, so are all the prisms which are in one pyramide, to all the prisms which are in the other pyramide, being equal in multitude.


- a 15. 5. For (applying the construction of the prece-
 b 22. 6. dent Prop.) BC. KC $a ::$ FG. QG. *b* therefore the
 c 2. 6. & c. triangle ABC is to the like triangle LKC as
 d 16. 5. EFG is to *c* the like RQG. therefore by permuta-
 e sch. 34. tion ABCEFG $d ::$ LKC. RQG $e ::$ the prism
 ii. KLCNMO. QRGTSV (for these are of equal
 f 7. 5. altitude) $f ::$ IBKLMN. PFQ RST. *g* wherefore
 g 12. 5. the triangle ABC. EFG : the prism KLCMNO
 + IBKLMN. the prism QRGTSV + PFQ RST.
 Which was to be dem.

But if the pyramides MNOD, AILM, and EPRS, STVH, be further divided, in like manner the four new prisms made hereby shall be to the four produced before, as the bases MNO and AIL are to the bases STV, and EPR; that is, as LKC to RQG, or as ABC to EFG. *b* wherefore all the prisms of the pyramide ABCD are to all the prisms of the pyramide EFGH as the base ABC is to the base EFG.

PROP.

Pyra
 altitud
 one to
 Let
 say X
 be pos
 cels b
 prism
 in lik
 SVT
 the p
 that t
 are g
 mide
 then
 PFQ
 pyr.
 RST
 whi
 A
 mak
 :: EL
 7 t
 imp
 to t

PROP. V.




Pyramides $ABCD$, $EFGH$, being under the same altitude, having triangular bases ABC , EFG , are one to another as their bases ABC , EFG , are.

Let the triangles ABC . $EFG :: ABCD$. X . I say X is equal to the pyramide $EFGH$. For if it be possible, let X be $\sqsubset EFGH$. and let the excess be Y , divide the pyramide $EFGH$ into prisms and pyramides, and the other pyramides in like manner, *a* till the pyramides left $EPRS$, $SVTH$, be less than the solid Y . Therefore since the pyramide $EFGH = X + Y$, it is manifest that the remaining prisms $PFQRST$, $QRGTSV$ are greater than the solid X . Conceive the pyramide $ABCD$ divided after the same manner; *b* then will be the prism $IBKLMN + KLCNMO$. $PFQRST + QRGTSV :: ABC$. EFG *c* :: the *c* *hyp.* pyr. $ABCD$. X . *d* therefore $X \sqsubset$ the prism $PFQRST + QRGTSV$; which is contrary to that which was affirmed before.

Again, conceive $X \sqsubset$ the pyr. $EFGH$. and make the pyr. $EFGH$. $Y :: X$ the pyr. $ABCD$ *e e hyp. and* $:: EFG$. ABC . Because $EFGH$ *f* $\sqsubset X$, *g* thence Y *cor. 4. 5.* $\sqsubset$ the pyr. $ABCD$. which is shewn before to be *f* *suppos.* impossible. Therefore I conclude, that X is equal *g* *14. 5.* to the pyr. $EFGH$. Which was to be dem.

PROP.


PROP. VI.



Pyramides $ABCDEF$, $GHIKLM$, consisting under the same altitude, and having polygonous bases $ABCDE$, $GHIKL$, are to one another as their bases $ABCDE$, $FGHIKL$ are.

Draw the right lines AC, AD, GI, GK . then is the base $ABC.ACD$ *a* :: the pyr. $ABCF.ACDF$. *b* therefore by composition $ABCD. ACD$:: the pyr. $ABCDF.ACDF$. *a* but also $ACD. ADE$:: the pyr. $ACDF. ADEF$. *c* therefore of equality $ABCD. ADE$:: $ABCDF. ADEF$. and *b* thence by composition $ABCDE. ADE$:: the pyr. $ABCDEF. ADEF$. moreover $ADE. GKL$ *d* :: the pyr. $ADEF. GKLM$; and as before, and inverely $GKL. GHIKL$:: the pyr. $GKLM. GHIKLM$. *c* therefore again of equality $ABCDE. GHIKL$:: the pyr. $ABCDEF. GHIKLM$. Which was to be dem.

a 5. 12.
b 18. 5.
c 22. 5.
d 5. 12.




a 5. 12.

f 4. 5.

If the bases have not sides of equal multitude, the demonstration will proceed thus. The base $ABC. GHI$ *e* :: the pyr. $ABCF. GHIK$. *e* and $ACD. GHI$:: the pyr. $ACDF. GHIK$. *f* therefore the base $ABCD. GHI$:: the pyr. $ABCDF. GHIK$. *e* Moreover the base $ADE. GHI$:: the pyr. $ADEF. GHIK$. *f* therefore the base $ABCDE. GHI$:: the pyr. $ABCDEF. GHIK$.

PROP.


PROP. VII.



Every Prism, $ABCDEF$,
having a triangular base,
may be divided into three
Pyramides $ACBF$, $ACDF$,
 $CDFE$, equal one to the

other, and having triangular bases.

Draw the diameters of the parallelograms,
 AC , CF , FD . Then the triangle ACB is $a = a$ 34. 1.
 ACD . b therefore the pyramides of equal height b 5. 11.
 $ACBF$. $ACDF$. are equal. In like manner the
pyr. $DFAC =$ the pyr. $DFEC$. but $ACDF$ and
 $DFAC$ are one and the same pyramide. c there-
fore the three pyramides $ACBF$, $ACDF$, $DFEC$,
into which the Prism is divided, are equal one
to the other. Which was to be dem.




Hence, every pyramide is
the third part of the Prism
that has the same base and
height with it; or every
prism is treble of the pyra-
mide that has the same base
and height with it.

For resolve the polygonous
prism $ABCDEFGHIKF$ into
triangular Prisms; and the
pyramide $ABCDEH$ into triangular pyramides;
 a then all the parts of the prism shall be treble a 7. 12.
to all the parts of the pyramide, b consequent-
ly the whole prism $ABCDEFGHIKF$ is tre-
ble to the whole pyramide $ABCDEH$. Which
was to be dem.


PROP.

PROP. VI.



Pyramides $ABCDEF$, $GHIKLM$, consisting under the same altitude, and having polygonous bases $ABCDE$, $GHIKL$, are to one another as their bases $ABCDE$, $FGHIKL$ are.

Draw the right lines AC , AD , GI , GK . then is the base $ABC.ACD$ a :: the pyr. $ABCF.ACDF$. b therefore by composition $ABCD.ACD$:: the pyr. $ABCDF.ACDF$. a but also $ACD.ADE$:: the pyr. $ACDF.ADEF$. c therefore of equality $ABCD.ADE$:: $ABCDF.ADEF$. and b thence by composition $ABCDE.ADE$:: the pyr. $ABCDEF.ADEF$. moreover $ADE.GKL$ d :: the pyr. $ADEF.GKLM$; and as before, and inversely $GKL.GHIKL$:: the pyr. $GKLM.GHIKLM$. e therefore again of equality $ABCDE.GHIKL$:: the pyr. $ABCDEF.GHIKLM$. Which was to be dem.




a 5. 12.

f 4. 5.

If the bases have not sides of equal multitude, the demonstration will proceed thus. The base $ABC.GHI$ e :: the pyr. $ABCF.GHIK$. e and $ACD.GHI$:: the pyr. $ACDF.GHIK$. f therefore the base $ABCD.GHI$:: the pyr. $ABCDF.GHIK$. e Moreover the base $ADE.GHI$:: the pyr. $ADEF.GHIK$. f therefore the base $ABCDE.GHI$:: the pyr. $ABCDEF.GHIK$.

PROPT


PROP. VII.



Every Prism, $ABCDEF$,
having a triangular base,
may be divided into three
Pyramides $ACBF$, $ACDF$,
 $CDFE$, equal one to the

other, and having triangular bases.

Draw the diameters of the parallelograms,
 AC , CF , FD . Then the triangle ACB is $a = a$ 34. 1.
 CD . b therefore the pyramides of equal height b 5. 11.
 $ACBF$. $ACDF$. are equal. In like manner the
pyr. $DFAC =$ the pyr. $DFEC$. but $ACDF$ and
 $DFAC$ are one and the same pyramide. c there-
fore the three pyramides $ACBF$, $ACDF$, $DFEC$,
into which the Prism is divided, are equal one
to the other. Which was to be dem.




Hence, every pyramide is
the third part of the Prism
that has the same base and
height with it; or every
prism is treble of the pyra-
mide that has the same base
and height with it.

For resolve the polygonous
prism $ABCDEGHIKF$ into
triangular Prisms; and the
pyramide $ABCDEH$ into triangular pyramides;
then all the parts of the prism shall be treble a 7. 12.
to all the parts of the pyramide, b consequent-
ly the whole prism $ABCDEGHIKF$ is tre-
ble to the whole pyramide $ABCDEH$. Which
was to be dem.

PROP.

PROP. VIII.



Like Pyramides $ABCD, EFGH$, which have triangular bases ABC, EFG are in triple proportion of the sides in which their sides of like proportion AC, EG , are

- a 27. 11. a Compleat the parallelepipeds $ABICDM$
 b 9. def. 11. $KL, EFNGHQOP$, which b are like, and
 c 28. 11. sextuple of the pyramids $ABCD, EFGH$. d and
 and 7. 12. therefore in the same proportion with them one
 d 15. 5. to another, e that is, triple of that of the sides
 e 33. 11. of like proportion, &c.

Coroll.

Hence, also like polygonous pyramids have proportion tripled to that of the sides of like proportion; as may easily be proved by resolving the same into triangular pyramids.

PROP. IX.

See the preceding Scheme.

In equal pyramids $ABCD, EFGH$, having triangular bases ABC, EFG , the bases and altitudes are reciprocal; And pyramids having triangular bases, whose altitudes and bases are reciprocal, are equal.

1. Hyp. The compleated parallelepipeds $ABICDM$
 a 28. 11. $DMKL, EFNGHQOP$ are a sextuple of the equal
 and 7. 12. pyramids $ABCD, EFGH$ (either of either) and so
 equal

equal one to the other. therefore the altitude
(H.) the altitude (D) $b :: ABIC$. $EFNG$ $c :: b$ 34. 11.
 ABC . EFG . Which was to be dem. c 15. 5.

2. Hyp. The altitude (H.) the altitude (D) $d :: d$ hyp.
 ABC . EFG $e :: ABIC$. $EFNG$. f therefore the 15. 5.
parallelepipeds $ABICDMKL$, $EFNGHQOP$ 34. 11.
are equal. g consequently also the pyramides g 6. ax. 1.
 $ABCD$, $EFGH$ being subseptuple of the same,
are equal. Which was to be dem.

The same is applicable to polygonous pyramides;
for they may also in like manner be reduced to tri-
angulars.

Coroll.

Whatsoever is demonstrated of pyramides in Prop. 6.
8, 9. does likewise agree to any sort of prisms; seeing
they are triple of the pyramides that have the same
base and altitude with them. Therefore

1. The proportion of prisms of equal altitude
is the same with that of their bases.
2. The proportion of like prisms is triple
of that of the sides of like proportion.
3. Equal prisms have their bases and altitudes
reciprocal; and prisms which are so reciprocal,
are equal.

Schol.


From what is hitherto demonstrated the di-
mension of any prisms and pyramides may be
collected.

a The solidity of a prism is produced of the a 1. cor. 12.
altitude multiplied into the base; b and there- E sch. 40.
fore likewise that of a pyramide, of the third 11.
part of the altitude multiplied into the base. b 7. 12.

U

PROP:

PROP. X.



Every Cone is the third part of a cylinder having the same base with it $ABCD$, and the altitude equal.

If you deny it, then first let such cylinder be E . A prism described on a square in the circle $ABCD$ *a* is subduple of a prism described upon a square about the circle, being equal to it and the cylinder in the height. Therefore a prism upon the square $ABCD$ exceeds the half of the cyl. and likewise a prism upon the base AFB , of equal height to the cylinder, *b* is greater than the half of the segment of the cylinder AFB , continue an equal bisection of the arches, and subtract the prisms till the remaining segments of the cylinder, namely at $AF, FB, \&c.$ become less than the solid E . Therefore the cyl. — segm. $AF, FB, \&c.$ (the prism on the base $AFBGCHDI$) *c* is greater than the cylinder — E (*d* the triple of the cone,) therefore the pyramide, *e* a third part of the said prism (being placed on the same base, and of the same height) is greater than the cone of equal height on the base $ABCD$ *i. e.* the part greater than the whole. Which is absurd.


But if the cone be affirmed to be greater than the third part of the cylinder, then let the excess be E . Detract the pyramides from the cone, as you did in the first part the prisms from the cylinder,

der, til
ceive a
therefo
the pyr
&c.)
(viz. of
is great
the par
fore it
to trip

Cyl
under
bases
Le
ABC
Fo
HM
and a
pose
of th
the p
mak
caul
AT
the
the
to

der, till some segments of the cone remain, conceive at AF, FB, EG, &c. less than the solid E. therefore the cone — E (f + of the cylinder) $\supset$ f hyp. the pyr. AFBGCHDI (the cone — segm. AF, FB, &c.) therefore the prism triple to the pyramide (viz. of equal height, and on the same base) is greater than the cylinder, on the base ABCD, the part than the whole, *Which is absurd*. Wherefore it must be granted, that the cylinder is equal to triple of the cone.

PROP. XL.



Cylinders and Cones *ABCDK*, *EFGHM*, being under the same altitude, are to one another as their bases *ABCD*, *EFGH*.

Let the circle *ABCD*. the cir. *EFGH* :: the cone *ABCDK*. *N*. If say *N* is equal to the cone *EFGHM*.

For if it be possible, let *N* be $\supset$ the cone *EFGHM*, and let the excess be *O*. The preparation

and argumentation of the prec. Prop. being supposed; then shall *O* be greater than the segments

of the cone *EP*, *PF*, *FQ*, &c. and so the solid *N* $\supset$ the pyr. *EPFQGRHSM*.


in the circle *ABCD*, *a* 23 o. 3. and make a like polygonous figure *ATBV CXDY*. Be 1. post.

cause the pyr. *ABVYK*. the pyr. *EFQSM* *b* :: *b* 6. 12. *ATBVY*. the polyg. *EPFQSc* :: the cir. *ABCD*. *c* cor. 2. 22.

the cir. *EFGH* *d* :: the cone *ABCDK*. *N*. *e* thence *d* hyp. the pyr. *EPFQGRHSM* shall be $\supset$ *N*. contrary *e* 14. 5.

to what was affirmed before. Again conceive *N* $\supset$ the

PROP. X.



Every Cone is the third part of a cylinder having the same base with it $ABCD$, and the altitude equal.

If you deny it, then first let such cylinder be E . A prism described on a square in the circle $ABCD$ a is subduple of a prism described upon a square about the circle, being equal to it and the cylinder in the heighth. Therefore a prism upon the square $ABCD$ exceeds the half of the cyl. and likewise a prism upon the base AFB , of equal heighth to the cylinder, b is greater than the half of the segment of the cylinder AFB , continue an equal bisection of the arches, and substract the prisms till the remaining segments of the cylinder, namely at $AF, FB, \&c.$ become less than the solid E . Therefore the cyl. — segm. $AF, FB, \&c.$ (the prism on the base $AFBGCHDI$) c is greater than the cylinder — E (d the triple of the cone.) therefore the pyramide, e a third part of the said prism (being placed on the same base, and of the same heighth) is greater than the cone of equal heighth on the base $ABCD$ a circle *i. e.* the part greater than the whole. Which is absurd.

But if the cone be affirmed to be greater than the third part of the cylinder, then let the excess be E . Detract the pyramides from the cone, as you did in the first part the prisms from the cylinder,


der, ti
ceive a
therefo
the py
&c.) t
(viz. of
is grea
the pa
fore it
to trip

T
B

Cyl
under
bases
Le
ABC
Fo
HM
-and
pose
of th
the
mak
caul
AT
the
the
to

der, till some segments of the cone remain, conceive at AF, FB, EG, &c. less than the solid E. therefore the cone — E ($\frac{1}{3}$ of the cylinder) $\supset$ *hyp.* the pyr. AFBGCHDI (the cone — segm. AF, FB, &c.) therefore the prism triple to the pyramide (viz. of equal height, and on the same base) is greater than the cylinder, on the base ABCD, the part than the whole, *Which is absurd.* Wherefore it must be granted, that the cylinder is equal to triple of the cone.

PROP. XL.



Cylinders and Cones $ABCDK$, $EFGHM$, being under the same altitude, are to one another as their bases $ABCD$, $EFGH$.

Let the circle ABCD. the cir. EFGH :: the cone ABCDK. N. I say N is equal to the cone EFGHM.

For if it be possible, let N be $\neg$ the cone EFG-
HM, and let the excess be O. The preparation
and argumentation of the prec. Prop. being sup-
posed; then shall O be greater than the segments
of the cone EP, PF, FQ, &c. and so the solid N $\neg$
the pyr. EPFQGRHSM. in the circle ABCD, *a* 23 0.3. and
make a like polygonous figure ATBVXDY. Be 1. *post.*
cause the pyr. ABVYK. the pyr. EFQSM *b* :: *b* 6. 12.
ATBVY. the polyg. EPFQSc :: the cir. ABCD. *c* *cor. 2.22.*
the cir. EFGH *d* :: the cone ABCDK. N. *e* thence *d* *hyp.*
the pyr. EPFQGRHSM shall be $\neg$ N. contrary *e* 14. 5.
to what was affirmed before. Again conceive N
U 2 $\neg$ the

the cone EFGHM. and make the cone EFGH.
 f hyp. and MO :: N. the cone ABCDK f :: the circle EF.
 by inver- GH. ABCD. g therefore O $\supset$ the cone ABCDK;
 jon. which is absurd, as appears by what is shewn in
 g 14. 5. the first part.

Therefore rather admit ABCD. EFGH :: the
 cone ABCDK. EFGHM. Which was to be dem.

The same may be demonstrated of cylinders,
 if cylinders and prisms be conceived in the place
 of cones and pyramids. therefore, &c.


Schol.

Hence is gathered the dimension of all sorts of cy-
 linders and cones. The solidity of a right cylin-
 der is produc'd of the circular base (a the di-
 mens. whereof is to be learnt out of Archime-
 des) multiplied into the heighth; b whence

b II. 12: in like manner that of every cylinder.

Therefore the solidity of a cone is produced
 of the third part of the altitude multiplied
 into the base.

PROP. XII.



Like cones and cylinders ABCDK, EFGHM, are
 in triplicate proportion of that of the diameters TX,
 PR, of their bases ABCD, EFGH.


Let the cone A have to N triplicate proportion
 of TX to PR. I say N is = the cone EFGHM. For
 if it be possible, let N be $\supset$ EFGHM. and let the
 excess be O. therefore, N $\supset$ the pyr. EPFQGR-
 HSM. Let the axes of the cones be IK, LM, and
 join the right lines VK, CK, VI, CI, and GM,
 GM,

GM, QL, GL. Because the cones are like, *a a* 24. *def.*
 thence VI. IK :: QL. LM. but the angles VIK. *ii.*
 QLM *b* are right angles. *c* therefore the trian- *b* 18. *def.*
 gles VIK, QLM are equiangular. *d* whence VC, *ii.*
 VI :: QG. QL. also VI. VK :: QL. QM. there- *c* 6. 6.
 fore of equality VC. VK :: QG. QM. *e* moreover *d* 4. 6.
 VK. GK :: QM. MG. therefore again of equa- *e* 7. 5.
 lity VC. CK :: QG. GM. *f* therefore the trian- *f* 5. 6.
 gle VKC, QMG are like : and by the same rea-
 son the other triangles of this pyramide are like
 to the other of that ; *g* wherefore the pyramides *g* 9. *def. ii*
 themselves are like. *b* But they are in triplicate *h* *cor. 8. 12.*
 proportion of that of VC to QG. *k* that is, of *k* 4. 6.
 VI to RL, *l* or TX to PR. *m* therefore the pyr. *l* 15. 5.
 AIBVCXDYK. the pyr. EPFQGRHSM :: the *m* *hyp. and*
 cone of ABCDK. N. *n* whence the pyr. EPFQ- *ii.* 5.
 GRHSM $\supset$ N. which is repugnant to what *n* 14. 5.
 was affirmed before.

Again, take N $\sqsubset$ the cone EFGHM. make
 the cone EFGHM. O :: N. the cone ABCDK
 o :: the pyr. EPRM. ATCK *p* :: GQ. VC thrice *o* before,
 :: *q* PR. TX thrice. but O *r* is $\supset$ ABCDK. and in-
 which was before shewn to be repugnant. *versely.*
 Wherefore N = the cone EFGHM. Which *p* *cor. 8. 12.*
 was to be dem. *q* 4. 6.

But forasmuch as what proportion soever *r* 14. 5.
 cones have, also cylinders being triple of them,
 have the same ; therefore cylinder to cylinder
 shall have proportion triplicate of the diameters
 of the bases,

PROP. XIII.



a 3. 1.

b 11. 12.

c 11. 12.

d 6. def. 5.


If a cylinder $ABCD$, be divided by a plane EF parallel to the opposite planes BC, AD , then as one cylinder $AFED$ is to the other cylinder $EBCF$, so is the axis GI to the axis HI .

The axis being produced, a take $GK=GI$, and $HL=IH=LM$, and conceive planes drawn at the points K, L, M , parallel to the circles AD, BC , b therefore the cylinder $FD =$ the cyl. AN . and the

cyl. $EC b = BO b = OP$. therefore the cylinder EN is as multiple of the cylinder ED as the axis IK is of the axis IG . and in like manner the cylinder FP is as multiple of the cylinder BF , as the axis IM is of the axis IH . but as

IK is $=, <, >$ IM , c so is the cylinder $EN =, <, >$ EP . d therefore the cylinder $AEFD$. the cyl. $EBCF :: GI. IH$. Which was to be dem.

PROP. XIV.



Cones AEB, CFD , and cylinders AH, CK consisting upon equal bases AB, CD , are to one another as their altitudes ME, NF .

The cylinder AH , and the axis EM being produced, take $ML=FN$; and at the

a 11. 12. point L draw a plane parallel to the base AB , a


b 13. 12. then shall the cyl. AP be $= CK$. b but the cyl. AH .

$AP(CK) :: ME.ML (NP.)$ Which was to be dem.

* apply 9. The same may be affirmed of cones subtriple and 7. 12. of cylinders; * as also of prisms and pyramids.

PROP.

PROP. XV.



In equal cones BAC, EDF , and cylinders BH, EK , the bases and altitudes are reciprocal $BC.EF :: MD.LA$. And cones and cylinders, whose bases and altitudes are reciprocal, are equal one to another.

If the altitudes be equal then the bases are equal too, and the thing is evident. If unequal, then take away $MO = LA$.


1. Hyp. Then is $MD.MO$ (a LA) $b ::$ the a 14. 12. cyl. EK . (c BH) $EQ d ::$ the cir. $BC.EF$. Which b constr. was to be dem. c hyp.

2. Hyp. $BC.EF e :: DM.OM$ (LA) $f ::$ the d 11. 12. cyl. $EK.FQ g :: BC.EF h :: BH.EQ$. Therefore e hyp. the cylinder $EK = BH$. Which was to be dem. f 11. 12.

The same argument may be used for cones.

g 11. 5.
h 11. 12.
k 9. 5.

PROP. XVI.




Two unequal circles $ABCG, DEF$, having the same center M , to inscribe in the greater circle $ABCG$ a polygonous figure of equal, and even sides, which shall not touch the lesser circle DEF .

Through the center M draw the line AC cutting the circle DEF in F , from whence raise a perpendicular FH . a di-a 30. 3. vide the semicircle ABC into two equal parts; and the half thereof BC also; and so do continually. b 1. 10. till the arch IC become less than the arch HC , from I let fall the perpendicular IL . It is manifest that the arch IC measures the whole circle, and that the number of arches is even, and

c sch. 16.4. so that the subtended line IC is the side of the polygon that may be inscribed without touching the lesser circle DEF. For HG touches the circle DEF. e to which IK is parallel, and placed outwardly; f wherefore IK does not touch the circle DEF; much less do CI, CK, and the other sides of the polygon more remote from the center. Which was to be done.

PROP. XVII.



Two spheres $ABCV$, $EFGH$, consisting about the same center D , being given, to inscribe a solid of many sides (or Polyedron) in the greater sphere $ABCV$, which shall not touch the superficies of the lesser sphere $EFGH$.

Let

Let both the spheres be cut by a plane passing by the center making the circles EFGH, ABCV; and the diameters AC, BV drawn, cutting perpendicularly. In the circle ABCV, *b* inscribe the equilateral polygone VMLNC, &c. not touching the circle EFGH: then draw the diameter Na, and erect DO perpendicular to the plane ABC. by DO, and by the diameters AC, Na, conceive planes DOC, DON erected, which shall be perpendicular to the circle ABCV, and so in the superficies of the sphere make *c* the quadrants DOC, DON. In which let the right lines CP, PQ, QR, RO, NS, ST, Ty, yO *d* be fitted, equal, and of equal multitude with CN, NL, &c. make the same construction in the other quadrants OL, OM, &c. and in the whole sphere. Then I say the thing required is done.

From the points P, S, to the Plane ABCV draw the perpendiculars PX, SY, *e* which shall fall on the sections ACNa. Therefore because both the right angles PXC, SYN, *g* and PCX, SNY insisting on *b* equal circumferences, *f* are equal, the triangles also PCX, SNY *h* are equiangular. Wherefore being PC *k* = SN, *l* also is PX = SY, *l* and XC = YN; *m* whence DX = DY. *n* and therefore DX. XC :: DY. YN. *o* therefore YX, NC are parallels. but because PX, SY are equal, and since being perpendicular to the same plane ABCV, they are also *p* parallels, *q* therefore YX, SP shall be equal and parallel. *r* whence SP, NC, are parallel one to the other; and so the quadrilateral NCPS. and by the same reason SPQT, TORG, as also the *t* triangle y RO are so many planes. In like manner the whole sphere may be shewn full of such quadrilaterals and triangles. wherefore the figure inscribed is a polyedron.

From the center D *u* draw DZ perpendicular to the plane NCPS; and join ZN, ZC, ZS, ZP. Because DN. NC *x* :: DY. YX. thence NC is y *y* YX (SP)

(SP.) and likewise $SP \perp TQ$, and $TQ \perp R$.
 And because the angles DZC, DZN, DZS, DZP
 z 3. def. 11. z are right, and the sides DC, DN, DS, DP , a equal,
 a 15. def. and DZ common, b thence ZC, ZN, ZS, ZP are e-
 11. qual one to the other; and consequently about
 b 47. 1. the quadrilateral $NCPS$ c a circle may be descri-
 c 15. def. 1. bed, in which (because NS, NC, CP , are d equal,
 d constr. and $NC \perp SP$) NC e subtends more than the qua-
 e 28. 3. drant, f therefore the ang. NZC at the center is
 f 33. 9. obtuse, g therefore $NCq \perp z ZCq$ ($ZCq + ZNq$)
 g 12. 2. Let NI be drawn perpendicular to AC . there-
 h 32. 1. fore since the angle ADN (b $DNC + DCN$) k is
 k 9. ax. 1. obtuse, the half of it DCN shall be greater than
 l 5. 1. the half of a right angle; and so that which re-
 mains of the right ang. CNI shall be less than it
 n 19. 1. n whence $IN \perp IC$. therefore NCq ($NIq + ICq$)
 o 47. 1. o $\perp INq$. therefore $IN \perp ZC$. and consequently
 p 47. 1. $DZp \perp DI$. but the point I is q without the sphere
 q cor. 19. $EFGH$. and so much more the point Z . wherefore
 12. the plane $NCPS$, (whose r next point to the cen-
 r 47. 1. ter is Z) does not touch the sphere $EFGH$. And
 if a perpendicular Ds be drawn to the plane
 $SPQT$, the point s , and so also the plane $SPQT$
 is yet further removed from the center, which
 is also true of the other planes of the polyedron.
 Therefore the polyedron $ORPQCN$; &c. in-
 scribed in the greater sphere, does not touch the
 lesser. Which was to be done.

Coroll.

Hence it follows, That if in any other sphere a
 solid polyedron, like to the abovesaid solid polyedron,
 be inscribed, the proportion of the polyedron in one
 sphere to the polyedron in the other is triplicate of that
 of the diameters of the spheres.

For if right lines be drawn from the centers of
 the spheres to all the angles of the bases of the
 said polyedrons, then the polyedrons will be di-
 vided into pyramids equal in number and like;
 whole


whole
 the sph
 spheres
 er abo
 drawn
 gles of
 reason
 like py
 pyram
 it in th
 that o
 semidi
 ramide
 that i
 are to
 lyedro
 polyed
 tron
 that o
 meter

H

Sp
 to t
 EF.
 L
 prop
 dian
 sible
 G o
 ED
 sph

whose homologous sides are semidiameters of the spheres; as appears, if the lesser of these spheres be conceived described within the greater about the same center. For the right lines drawn from the center of the sphere to the angles of the bases will agree one to the other by reason of the likeness of the bases; and so will like pyramids be made. Wherefore since every pyramid in one sphere to every pyramid like it in the other sphere, *a* has proportion triple to *a* *cor. 8. 12.* that of the homologous sides, that is, of the semidiameters of the spheres; and *b* as one py- *b 12. 5.* ramide is to one pyramid, so all the pyramids, that is, the solid polyedron composed of these, are to all the pyramids, that is, the solid polyedron composed of the others; therefore the polyedron of one space shall have to the polyedron of the other sphere, proportion triple of that of the semidiameters, *c* and so of the dia- *c 15. 5.* meters of the spheres.

P R O P. XVIII.



Spheres *BAC, EDF*, are in triplicate proportion one to the other of that in which their diameters *BC, EF*.

Let the sphere *BAC* be to the sphere *G* in triple proportion of that of the diameter *BC* to the diameter *EF*. I say *G = DEF*. For if it be possible, let *G* be $\supset$ *EDF*. and conceive the sphere *G* concentric with *EDF*. In the sphere of *EDF* *a* inscribe a polyedron not touching the *a 17. 12.* sphere *G*, and a like polyedron in the sphere *BAC*.

(SP.) and likewise $SP \sqsubset TQ$, and $TQ \sqsubset \gamma R$.
 And because the angles DZC, DZN, DZS, DZP
 z 3. def. 11. z are right, and the sides DC, DN, DS, DP , a equal,
 a 15. def. 1. and DZ common, b thence ZC, ZN, ZS, ZP are e-
 11. qual one to the other; and consequently about
 b 47. 1. the quadrilateral $NCPS$ c a circle may be descri-
 c 15. def. 1. bed, in which (because NS, NC, CP , are d equal,
 d constr. and $NC \sqsubset SP$) NC subtends more than the qua-
 e 28. 3. drant, f therefore the ang. NZC at the center is
 f 33. 9. obtuse, g therefore $NCq \sqsubset 2 ZCq (ZCq + ZNq)$
 g 12. 2. Let NI be drawn perpendicular to AC . there-
 h 32. 1. fore since the angle ADN (b $DNC + DCN$) k is
 k 9. ax. 1. obtuse, the half of it DCN shall be greater than
 l 5. 1. the half of a right angle; and so that which re-
 mains of the right ang. CNI shall be less than it
 n 19. 1. n whence $IN \sqsubset IC$. therefore $NCq (NIq - ICq)$
 o 47. 1. $o \sqsubset INq$. therefore $IN \sqsubset ZC$. and consequently
 p 47. 1. $DZp \sqsubset DI$. but the point I is q without the sphere
 q cor. 19. $EFGH$. and so much more the point Z . wherefore
 12. the plane $NCPS$, (whose r next point to the cen-
 r 47. 1. ter is Z) does not touch the sphere $EFGH$. And
 if a perpendicular $D \delta$ be drawn to the plane
 $SPQT$, the point δ , and so also the plane $SPQT$
 is yet further removed from the center, which
 is also true of the other planes of the polyedron.
 Therefore the polyedron $ORPQCN$, &c. in-
 scribed in the greater sphere, does not touch the
 lesser. Which was to be done.

Coroll.

Hence it follows, That if in any other sphere a
 solid polyedron, like to the abovesaid solid polyedron,
 be inscribed, the proportion of the polyedron in one
 sphere to the polyedron in the other is triplicate of that
 of the diameters of the spheres.

For if right lines be drawn from the centers of
 the spheres to all the angles of the bases of the
 said polyedrons, then the polyedrons will be di-
 vided into pyramides equal in number and like;
 whose


whose
 the sph
 spheres
 er abo
 drawn
 gles of
 reason
 like py
 pyram
 it in th
 that o
 semidi
 ramide
 that is
 are to
 lyedron
 polyed
 dion
 that o
 meters

H

Sp
 to t
 EF.
 L
 prop
 dian
 fible
 G c
 ED
 sph

whose homologous sides are semidiameters of the spheres; as appears, if the lesser of these spheres be conceived described within the greater about the same center. For the right lines drawn from the center of the sphere to the angles of the bases will agree one to the other by reason of the likeness of the bases; and so will like pyramids be made. Wherefore since every pyramid in one sphere to every pyramid like it in the other sphere, *a* has proportion triple to *a* cor. 8. 12. that of the homologous sides, that is, of the semidiameters of the spheres; and *b* as one py- b 12. 5. ramide is to one pyramid, so all the pyramids, that is, the solid polyedron composed of these, are to all the pyramids, that is, the solid polyedron composed of the others; therefore the polyedron of one space shall have to the polyedron of the other sphere, proportion triple of that of the semidiameters, *c* and so of the dia- c 15. 5. meters of the spheres.

PROP. XVIII.



Spheres *BAC*, *EDF*, are in triplicate proportion one to the other of that in which their diameters *BC*, *EF*.

Let the sphere *BAC* be to the sphere *G* in triple proportion of that of the diameter *BC* to the diameter *EF*. I say $G = DEF$. For if it be possible, let *G* be $\neq EDF$. and conceive the sphere *G* concentric with *EDF*. In the sphere of *EDF* *a* inscribe a polyedron not touching the a 17. 12. sphere *G*, and a like polyedron in the sphere *BAC*.

b cor. 17. BAC. These polyedrons *b* are in triplicate proportion of the diameters BC, EF. *c* that is, of the sphere BAC to G. *d* Consequently the sphere G is greater than the polyedron inscribed in the sphere EDF, the part than the whole.

Again, if it be possible, let the sphere G be $\sqsubset$ EDF. and as the sphere EDF is to another sphere H, so let G be to BAC, *e* that is, in proportion of the diameter EF to BC. therefore since BAC $f \sqsubset$ H, we shall incur the absurdity of the first part. wherefore rather the sphere $G = EDF$. Which was to be dem.

Coroll.

Hence, As one sphere is to another sphere, so is a Polyedron described in that, to a like polyedron described in this.


The End of the twelfth Book.

THE

THE THIRTEENTH BOOK OF EUCLIDE'S ELEMENTS.

PROP. I.

If a right line z be divided according to extreme and mean proportion ($z. a :: a. e$) the square of the half of the whole line z , and of the greater segment a , as one line is quintuple to that which is described of half of that whole line z .



I say Q. $a + \frac{1}{2} z$
 $= 5 Q: \frac{1}{2} z. a$ that $a 4. 2.$
 is, $aa + \frac{1}{4} zz + za = zz.$ For $ze + c 2. 2.$
 $za c = zz.$ and $ze d = aa.$ e therefore $aa + za d$ hyp. and
 $= zz.$ Which was to be dem. 16. 6.
e z. ax. and
1. ax.

PROP. II.

See the 1. Scheme.

If a right line $\frac{1}{2} z + a$ be in power quintuple to a segment of it self $\frac{1}{2} z.$ the line double of the said segment (z) being divided according to extreme and mean proportion, the greater segment is (a) the other part of the right line at first given $\frac{1}{2} z + a.$


I say $z. a :: a. e.$ For because by the hyp. $* aa * 4. 2.$
 $+ \frac{1}{4} zz + za = zz + \frac{1}{4} zz;$ or $aa + za = zz a 2. 2.$
 $= ze + za, b$ thence shall aa be $= ze. c$ where- $b 3. ax. 1.$
 fore $z. a :: a. e.$ Which was to be dem. c 17. 6.

PROP.

PROP. III.

If a right line z be divided according to extreme and mean proportion ($z. a :: a. e.$) the line made of the less segment e and half of the greater segment a , is in power quintuple to the square, which is described of the half line of the greater segment a .

a 4. 2.
b 3. ax.
c 3. 2.
d hyp. and
17. 6.



I say $Q: e + \frac{1}{2} a = 5$
 $Q: \frac{1}{2} a. a$ that is $ee +$
 $\frac{1}{4} aa + ea = aa + \frac{1}{4} aa.$
 b or $ee + ea = aa.$ For

$ee + ea = ze d = aa.$ Which was to be dem.

PROP. IV.

If a right line z be cut according to extreme and mean proportion ($z. a :: a. e.$) the square made of the whole line z , and that made of the lesser segment e , both together, are triple of the square made of the greater segment a .


a 4. 1.
b 3. 2.
c 17. 6.
d 2. ax.



I say $zz + ee = 3$
 $aa. a$ or $aa + ee + 2$
 $ae + ee = 3 aa.$ For
 $ae + ee = ze c =$

$aa. d$ therefore $aa + 2 ae + 2 ee = 3 aa.$ Which was to be dem.

PROP. V.



If a right line AB be cut according to extreme and mean proportion in C , and a line AD , equal to the greater segment BC , added to it, the whole right line DB is divided according to extreme and mean proportion; and the greater segment the right line AB given at the beginning.

For because $AB. AD :: AC. CB.$ and by inversion $AD. AB :: CB. AC.$ therefore by composition $DB. AB :: AB. AC$ (AD.) Which was to be dem.

Schol.

Schol.


But if $BD.BA :: BA.AD$. then shall be $BA:AD :: AD.BA - AD$. For by division is $BD - BA (AD.) BA :: BA - AD.AD$. therefore inversely $BA.AD :: AD.BA - AD$.

PROP. VI.

$D \quad A \quad C \quad B$ If a rational right line
 $\text{---} 1 \text{---} 1 \text{---}$ AB be cut according to extreme and mean proportion
in C , either of the segments (AC, CB) is an irrational line of that kind which is called apotome or residual.

To the greater segment AC add $AD = \frac{1}{2} AB$. a 3. 1.
b therefore $DCq = 5 DAq$. c therefore $DCq \sqsupset b$ 1. 13.
 DAq . consequently d since AB , e and so the c 6. 10.
half thereof DA are ρ , likewise DC is ρ . But d hyp.
because $5. 1 :: \text{not } Q. Q$. f thence is $DC \sqsupset DA$. e sch. 12.
g therefore $DC - AD$, that is, AC , is a residual line. Further, because $ACq \cdot b = AB \times f$ 9. 10.
 BC , and AB is ρ . k likewise EC is a residual g 74. 10.
line. Which was to be dem. h 17. 6.
k 98. 10.

PROP. VII.



If three angles of an equilateral Pentagone $ABCDE$ whether they follow in order, (EAB, ABC, BCD) or not, (EAB, BCD, CDE) be equal, the Pentagone $ABCDE$ shall be equiangular.


Let

Let the right lines BE, AC, BD, be subtended to the equal angles in order.

- a hyp. Being the sides EA, AB, BC, CD, and the included angles *a* are equal, therefore shall the bases BE, AC, BD, *c* and the angles AEB, ABE, c 4. and 5. BAC, BCA, be equal. *d* wherefore $BF = FA$. e and consequently $FC = FE$; therefore the triangles FCD, FED, are equilateral one to the other: d 6. i. f whence the angle $FCD = FED$. g consequently e 3. ax. i. the angle $AED = BCD$. In like manner the ang. f 8. i. g 2. ax. i. CDE is equal to the rest; wherefore the pentagone is equiangular. Which was to be dem.

- h 4. i. But if the angles EAB, BCD, CDE, which are not in order, be supposed equal. *b* then shall k 5. i. the angle AEB be $= BDC$, and $BE = BD$. l and thence the angle $BED = BDE$. l consequently the whole angle $AED = CDE$. therefore because the angles A, E, D, in order, are equal, as before, the pentagone shall be equiangular. Which was to be dem.

P R O P. VIII.




If in an equilateral and equiangular Pentagone ABCDE, two right lines BD, CE, subtend two angles BCD, CDE, following in order, those lines do cut one another according to extreme and mean proportion, and their greater segments BF

or EF are equal to the side of the pentagone BC.

- a Describe about the pentagone the circle ABD. b 28. 3. The arch ED is $= BC$, *c* therefore the angle FCD $= FDC$, *d* therefore the ang. $BFC = 2 FCD$ (FCD + FDC. But the arch BAE *b* is $= 2 ED$, and consequently the angle BCF *e* $= 2 FCD = BFC$. *f* wherefore $BF = BC$. Which was to be dem. g 27. 3. Moreover because the triangles BCD, FCD, are h 4. 6. g equiangular, *h* therefore $BD : DC$ (BF.) $:: CD$. (BF)

(BF.) FD. and likewise EC.EF :: EF. FC. Which was to be dem.

PROP. IX.



If the side of an Hexagone BE, and the side of a Decagone AB, both described in the same circle ABC, be added together, the whole right line AE is cut according to extreme and mean proportion (AE. BE :: BE. AB.) and

the greater segment thereof is the side of the Hexagone BE.

Draw the diameter ADC, and join the right lines DB, DE. Because the angle BDC $a = 4a$ hyp. and BDA. and the angle BDC $b = 2$ DBA (DAC + 27.3. DBA) thence shall DBA (b DBE + BED) c be b 32. 1. $= 2$ BDA $d = 2$ BDE. whence the angle DBA c 7. ax. 1. or DAB $e =$ ADE. Therefore the triangles ADE, d 5. 1. ADB, are equiangular: f wherefore AE. AD (e 1. ax. 1. BE) :: AD. (BE.) AB. Which was to be dem. f 4. 6.

g cor. 1 5. 4.


Coroll.

Hence, If the side of an Hexagone in a circle be cut according to extreme and mean proportion; the greater segment thereof shall be the side of the Decagone in the same circle.

X

PROP.

PROP. X.



If an equilateral Pentagone $A-BCDE$ be described in a circle $ABCE$, the side of the Pentagone AB contains in power both the side of a hexagone FB , and the side of a decagone AH described in the same circle.

Draw the dia-

meter AG . and bisect equally the arch AH in K . and draw FK , FH , FB , BH , HM .

a 28.3. and The semicircle AG — the arch AC $a = AG$
3. ax. — AD . that is, the arch $CG = GD$ $b = AH$
b hyp. and $= HB$. therefore the arch $BCG = 2 BHK$; c
7. ax. and so the angle $BFG = 2 BFK$. d but the an-
c 33 6. gle $BFG = 2 BAG$. e therefore the angle BFK
d 20. 3. $= BAG$. Wherefore the triangles BFM , FAB , f
e 1. ax. 1. are equiangular. g whence $AB \cdot BF :: BF \cdot KM$. h
f 31. 1. therefore $AB \times BM = BF^2$. Moreover, the an-
g 4. 6. gle AFK $k = HFK$, and $FA = FH$. m where-
k 27. 3. fore $AL = LH$. m and the angles FLA , FLH
m 4. 1. are equal, and so right angles. therefore the an-
n 27. 3. gle LHM $m = LAM$ $n = HBA$. therefore the
o 32. 1. triangles AHB , AMH , o are equiangular. where-
p 4. 6. fore $AB \cdot AH :: AH \cdot AM$. q therefore $AB \times AM$
q 17. 6. $= AH^2$. So that seeing $AB^2 = AB \times BM +$
r 2. 2. $AB \times AM$, f thence $AB^2 = BF^2 + AH^2$. Which
f 2. ax. was to be dem.

Coroll.

1. Hence, a rightline (FK) which being drawn from the center (F) divides an arch (HA) into two equal segments, does also divide the right line


line (HA) subtending that arch, perpendicularly into two equal segments.

2. The diameter of a circle (AG) drawn from any angle (A) of a pentagone, does divide equally in two both the arch (CD) which the side of the pentagone opposite to that angle subtends, and also the opposite side it self (CD) and that perpendicularly.

Schol.

Here, according to our Promise, we shall lay down a ready praxis of the II. Prop. of the 4. Book.

Problem.



To find out the side of a pentagone to be inscribed in a circle ADB.

Draw the diameter AB, to which erect a perpendicular CD at the center C divide CB equally in E. and make $EF = ED$. then DF shall be the side of the pentagone.

a 6. 2.


For $BF \times FC + ECq = EFq = EDq = DCq - ECq$. d therefore $BF \times FC = DCq$ or c 47. 1. BCq . e wherefore $BF : BC :: BC : FC$. therefore d 3. ax. since BC is the side of an hexagone, f FC shall e 17. 6. be the side of a decagone. Consequently DF f 9. 13. $b = \sqrt{DCq - FCq}$ g is the side of a pentagone. g 10. 13. Which was to be done.

h 47. 1.

X 2

PROP.

PROP. XI.




If in a circle $ABCD$, whose diameter is rational AG , an equilateral pentagone be inscribed $ABCDE$; the side of the pentagone AB is an irrational line of that kind which is called a minor line.

Draw the diameter

- * 10. 6. BFH , and the right lines AC, AH ; and * make $FL = \frac{1}{4}$ of the radius FH ; and $CM = \frac{1}{4} CA$.
 a cor. 10. Because the angles AKF, AIC , are a right angles, and CAI common, the triangles AKF, AIC , are b equiangular: c therefore $CI.FK :: CA.FA$ (FB) $d :: CM.FL$. therefore by permutation $FK.FL :: CI.CM$ $d :: CD.CK$ ($2 CM$) and so by e composition $CD + CK.CK :: KL.FL$. f consequently $Q: CD + CK$ (g $5 CK$) $CKq :: KLq.FLq$. therefore $KLq = 5 FLq$. wherefore if BH (ρ) be taken 8, FH shall be 4, FL 1, and FLq 1, BL 5, and BLq 25, KLq 5, by which it appears that BL and KL are ρ b $\sqrt{5}$, k and so BK is a residual, and KL its congruent or adjoining line, but being $BLq - KLq = 20$, m cor. 8.6. l thence $BL = \sqrt{BLq - KLq}$. m whence BK and 17. 6. shall be a fourth residual line. Therefore because n 95. 10. ABq m is $= HB \times BK$, n shall AB be a minor line. Which was to be dem.

PROP.

PROP. XII.




If in a circle *ABEC* an equilateral triangle *ABC* be inscribed, the side of that triangle *AB* is in power triple to the line *AD* drawn from *D* the center of the circle to the circumference.

The diameter being extended to *E*, draw *BE*. Because the arch *BE* $a = EC$, the arch *BE* is the a cor. 10. sixth part of the circumference, b therefore *BE* 13. = *DE*. hence $Aeq\ c = 4\ DEq\ (4\ BEq)\ d = b$ cor. 15. 4. $ABq + BEq (+ ADq.)\ e$ consequently $ABq = c\ 4.\ 2.$ $3\ ADq.$ Which was to be dem. $d\ 47.\ 1,$ $e\ 3.\ ax.\ 1.$
Coroll.

1. $Aeq.\ ABq :: 4.\ 3.$
2. $ABq.\ AFq :: 4.\ 3.$ f For $ABq.\ AFq :: Aeq.\ f$ cor. 8. 6. and 22. 6.
3. $DF = FE$. For the triangle *EBD* g is equilateral, b and *BF* perpendicular to *ED*. h cor. 3. 3. therefore $EF = FD$.
4. Hence, $AF = DE + DF = 3\ DF$.

PROP. XIII.



To describe a pyramid *EGFI*, and comprehend it in a sphere given: and to demonstrate that the diameter of the sphere *AB* is in power sesquialter of the side *EF* of the pyramid *EGFI*.

X 3

About

a 10. 6. About AB describe the semicircle ADB; *a* and let AC be = 2 CB. from the point C erect the perpendicular line CD; and join AD, DB, then at the interval of the radius HE = CD describe the circle HEFG, wherein inscribe the equilateral triangle EFG from H erect IH = CA perpendicular to the plane EFG. produce IH to K, *d* so that IK = AB; and join the right lines IE, IF, IG. Then EFGI shall be the pyramide required.

For because the angles ACD, IHE, IHF, IHG, *e* are right angles; and CD, HE, HF, HG *e* equal. *e* and IH = AC; *f* therefore AD, IE, IF, IG shall be equal among themselves. But being AC (2 CB) CB *g* :: ACq. CDq. then shall ACq be = 2 CDq. therefore ADq *f* = ACq + CDq *b* = 3 CDq = 3 HEq *k* = EFq. *l* therefore AD, IE, IF, IG are equal, and so the pyramide EFGI is equilateral. But if the point C be placed upon H, and AC upon HI, the right lines AB, IH, *m* shall agree, as being equal. Wherefore the semicircle ADB being drawn about the axis AB or IK *n* shall pass by the point E, F, G, * and so the pyramide EFGI shall be inscribed in a sphere. Which was to be done.

o cor. 8. 6. Also it is manifest that BAq. ADq o :: BA. p constr. AC p :: 3. 2. Which was to be dem.

Coroll.

1. ABq. HEq :: 9. 2. For if ABq be put 9, then ACq (EFq) shall be 9, consequently HEq shall be 2.

2. If L be the center, then shall AB.LC :: 6. 1. For if AB be put 6, then AL shall be 3. *r* and thence AC 4. wherefore LC shall be 1. Hence

3. AB. HI :: 6. 4 :: 3. 2. whence


4. ABq. HIq :: 9. 4.

PROP.



Ab
from
draw
suar
the c
lar to
and j
shall
F
equ
the
tria
cor
LI
an
be
A
an
is
L

PROP. XIV.



To describe an Octaedron KEFGDL, and comprehend it in the given sphere, wherein a Pyramide is: and to demonstrate that AH the diameter of the sphere is in power double of AC the side of that Octaedron.

About H describe the semicircle ACH. and from the center B erect the perpendicular BC. draw AC, HC, then upon ED=AC. a make the a 46. 1. square EFGD, whose diameters DF, EG, cut in the center I. from I draw IL=AB perpendicular to the plane EFGD. produce IL, c till IK=IL. c 3. 1. and join KE, KF, KG, KD, LE, LF, LG, LD; then shall KEFGDL be the Octaedron required.

For AB, BH, FI, IE, &c. being semidiameters of equal squares are equal one to the other. d whence d 4. 1. the bases LF, LE, FE, &c. of the right-angled triangles LIE, LIF, FIE, &c. are equal, and consequently the eight triangles LFE, LEG, LGD, LDE, KEF, KFG, KGD, KDE, are equilateral, e e 27. def. and make an Octaedron, which may be inscri- 11. bed in a sphere, whose center is I, and IL or AB the radius. (because AB, IL, IF, IK, &c. ff constr. are equal.) Which was to be done. Moreover, it is evident that AHq (LKq) g = 2 ACq (2 g 47. 1. LDq.) Which was to be dem.

Coroll.

1. Hence it is manifest, that in the octaedron the three diameters, EG, FD, LK do cut one the other perpendicularly in the center of the sphere.

2. Also that the three planes EFGD, LEKG, LFKD are squares, cutting one another perpendicularly.

X 4


3. The

3. The Octaedron is divided into two like and equal pyramides EFGDL, and EFGDK, whose common base is the square EFGD.

75. II.

4. Lastly, It follows that the opposite bases of the Octaedron are parallel one to the other.

PROP. XV.



To describe a cube
EFGHIKLM, and
comprehend it in the
same sphere wherein
the former figures
were; and to demon-
strate that AB the
diameter of the

sphere is in power triple to EF the side of that cube.

210.6.

Upon AB describe a semicircle ACB; a and make $AB = 3 DA$, from D raise the perpendicular DC, and join BC and AC. Then upon $EF = AC$ b make the square EFGH, upon whose plane let the right lines EI, FK, HM, GL, stand perpendicular, being equal to EF, and connect them with the right lines IK, KL, LM, IM. The solid EFGHIKLM is a cube, as is sufficiently apparent from the construction.

b 46, II.

In the opposite squares EFKI, HGLM, draw the diameters EK, FI, HL, MG, by which let the planes EKLH, FI, MG be drawn, cutting one another in the line NO. which c shall divide equally in two parts the diameters of the cube EL, FM, GL, HK, in P the center of the cube, d therefore P shall be the center of a sphere passing by the angular points of the cube. Moreover, $ELq e = EKq + KLq, e = 3 KLq, f$ or $3 ACq$, but $ABq. ACq :: BA. DA f :: 3. i. b$ therefore $AB = EL$. wherefore we have made a cube, $\&c$. Which was to be done.

£ cor. 39.

II.

d 15. def.

II. and 14.

def. II.

€ 47. I.

‡ constr.

g cor. 8. 6

h 14. 5.

Coroll.

1. Hence it is manifest, that all the diameters of the cube are equal one to another, and do equally bisect one another in the center of the sphere. And by the same means the right lines which

which are equal.

2. The
the side
 $k = 1/BO$

To
ZGH
compa
were
lids;
FG
that
called


U
sph
AD
the
dic
BD
BE

which conjoin the centers of the opposite squares are equally bisected in the same center.

2. The diameter of a sphere contains in power the side of a tetraedron and of a eube, viz. $ABq = l BCq + m ACq$.

PROP. XVI.

k 47. r.
l 13. 15.
m 15. 13.



To describe an Icosaedron $ZGHKIFYVXRST$, and encompass it in the sphere, wherein were contained the foresaid solids; and to demonstrate that FG the side of the Icosaedron is that irrational line which is called a minor line.

Upon AB the diameter of a sphere describe the semicircle ADB ; and make $AB = 5 BC$. then from C erect CD perpendicular, and draw AD and BD . At the distance $EF = BD$ describe the circle EFK -

NG ; A



a 10. 6.

- b 11. 4. NG; *b* wherein inscribe the equilateral pentagone FKIHG. Divide equally in two parts the arches FG, GH, &c. and join the right lines FL, LG, &c. being the sides of a decagone. Then *c* erect EQ, LR, MS, NT, OV, PX equal to EF, and perpendicular to the plane FKNG; and connect RS, ST, TV, VX, XR; as also FX, FR, GR, GS, HS, ST, HT, IT, IV, KV, KX. Lastly, produce FQ, and take QY = FL, and EZ = FL: and conceive the right lines ZG, ZH, ZI, ZK, ZF to be drawn; as also YV, YX, YR, YS, YT. Then I say the Icosaedron required is made.
- d *constr.* For because EQ, LR, MS, NT, OV, PX, are *d*-equal and *e* parallel, also those lines that join them
- e 6. 11. EL, QR, EM, QS, EN, QT, EO, QV, EP, QX, *f* are
- f 33. 1. equal and parallel. And thence likewise LM (or FG) RS, MN, ST, &c. are equal one to the other.
- g 15. 11. *g* therefore the plane drawn by EL, EM, &c. is equidistant from the plane passing by QR, QS, &c. *h* and the circle QXRSTV drawn from the center Q is equal to the circle EPLMNO; and, RSTVX is an equilateral pentagone. But EF, EG, EH, &c. and QX, QR, QS, &c. being conceived to
- k 47. 1. be drawn; then because FRq *k* = FEq + LRq, *l* or
- l *constr.* EFq *m* = FGq, *n* therefore FR, FG, and so all
- m 10. 13. RS, FG, FR, RG, GS, GH, &c. shall be equal one
- n *sch.* 48. 1. to the other, and consequently the ten triangles
- and 1. *ax.* RFX, RFG, RGS, &c. are equilateral and equal.
- o *cor* 14. Moreover, because XQY is a *o* right angle, there-
11. fore XYq *p* = QXq + QYq *q* = VXq or FGq.
- p 47. 1. wherefore XY, VX. and likewise YV, YT, YS, YR,
- q 10. 13. ZG, ZH, &c. are equal. Therefore other ten triangles are made equilateral and equal both one to another, and to the ten former; and so an Icosaedron is made.

Moreover, divide equally EQ in *a*, draw the

r 15. *def.* 1. right lines *a*F, *a*X, *a*V; and because QX *r* = QV, and *a*Q the common side, and FQX, EQV are

f 4. 1. right angles, *f* therefore shall *a*X be = *a*V; and by the same reason all the lines *a*X, *a*R, *a*S, *a*T, *a*V,

*a*V, *a*F
ZQ, QV
x = *E*
Za =
fore the
ray, the
Icosaedron
Last

Za, q. *a*
QEq, *c*
1. *d* the
The
shall be
the per
is a mi

1. F
of the
semidi
sides *c*
2. *a*
the sp
that i
the de
sides
3. *l*
of an
For F

aV, aF, aG, aH, aI, aK are equal. But because t 9. 13.
 $ZQ.QE :: QE.ZE$. therefore $Zaq u = 5 Eaqu$ 3. 13.
 $x = EQq (EFq) + Eaq y = aFq$. therefore x 4. 2.
 $Za = aF$. z in like manner $aF = Ya$. there-
 fore the sphere, whose center is a , and aF the z 15. 5.
 ray, shall pass by the 12 angular points of the
 Icosaedron.

Lastly, Because $Za. aE :: ZY. QE$; a and foa 22. 6.
 $Zaq. aEq :: ZYq. QEq$, b therefore $ZYq = 5b$ 14. 5.
 QEq , or $5 BDq =$; but $ABq. BDq c :: AB.BC :: 5. c$ cor. 8. 6.
 1. d therefore $ZY = AB$. Which was to be done. d 1. ax. 1.

Therefore if AB be put ρ , c then $EF = \sqrt{ABqe}$ sch. 12.
 shall be also ρ . and consequently FG the side of re .
 the pentagone, and likewise of the Icosaedron, ff 11. 13.
 is a minor line. Which was to be dem.

Coroll.

1. From hence is inferred, that the diameter
 of the sphere is in power quintuple of the
 semidiameter of the circle encompassing the five
 sides of the Icosaedron.


2. Also it is manifest that the diameter of
 the sphere is composed of the side of a hexagone,
 that is, of the semidiameter, and two sides of
 the decagone of a circle encompassing the five
 sides of the Icosaedron.

3. It appears likewise that the opposite sides
 of an Icosaedron, such as RX, HI , are parallels.
 For RX a is parall. to LP . b parall. to HI .

a 33. 1.
 b sch. 26. 3.

PROP.

PROP. XVII.



To describe a Dodecaedron, and comprehend it in the sphere wherein the former figures were comprehended: and to demonstrate that the side RS of the Dodecaedron is an irrational line of that sort which is called an Apotome or residual line.

2 30. 6.

Let AB be a cube inscribed in the given sphere, and let all the sides thereof be divided equally in the points E, H, F, G, K, L, &c. and join the right lines KL, MH, HG, EF. & make $HI:IQ::IQ:QH$; and take $NO, NP, = IQ$. then erect OR, PS, perpendicular to the plane DB, and QT to the plane AC, and let OR, PS, QT, be equal to IQ, NO, NP, whence DR, RS, SC, CT, DT, being connected, DRSC T shall be a pentagone of the dodecaedron required. For draw NV parallel to OR, and having drawn NV out as far as the center of the cube X, join the right lines DS, DO, DP, CR, CP,

CP, HV, HT, RX. Because $DOq\ a = DKq\ (bKNq)\ a\ 47.\ 1.$
 $+KOq\ c = 3\ ONq\ (3\ ORq)\ d$ thence $DRq = b\ 7.\ ax.\ 1.$
 $ORq\ e = OPq$, or RSq . therefore $DR = RS.\ c\ 4.\ 13.$
 By the same reason DR, RS, SC, CT, TP are equal. $d\ 47.\ 1.$
 But because $OR\ f$ is $=$ and g parallel to PS , $e\ 4.\ 2.$
 therefore RS, OP , and consequently RS, DC shall *constr.* $96.$
 be also parallels. h therefore these with them $11.$
 that conjoin them DK, CS, VH , are in one and the $g\ 33.\ 1.$
 same plane. Moreover, because $HI, IQ\ b :: IQ.\ h\ 9.\ 1.$
 $(TQ.)\ QH\ k :: HN.\ NV.$ and both TQ, HN , and $k\ 7.\ 11.$
 $QH, NV\ k$ are perpendicular to the same plane, $l\ 6.\ 11.$
 and so likewise parallels, $m\ THV$ shall be a right $m\ 32.\ 6.$
 line. n therefore the Trapezium $DRSC$, and the $n\ 1.\ and\ 2.$
 triangle DTS are in one plane extended by the $11.$
 right lines $DC, TV.$ o therefore $DCTSR$ is a pen- $o\ 5.\ 13.$
 tagone, and that also equilateral, by what is
 shewn already. Furthermore, because $PK.\ KN ::$
 $KN.NP$; and $DSq\ p = DPq + PSq\ (PNq) = p\ p\ 47.\ 1.$
 $DKq + PKq + NPq,\ q$ thence $DSq = DKq + 3\ KNq\ q\ 1.\ ax.\ 1.$
 $= 4DKq\ (4\ DHq)\ r = DCq$. therefore $DS = DC.$ and $4.\ 13.$
 whence the triangles DRS, DCT , are equilateral $r\ 4.\ 2.$
 one to another, s therefore the angle $DRS = DTC,$ $f\ 8.\ 1.$
 therefore the pentagone $DTCSR$ is also equian-
 gular. Moreover, because $AX, DX, CX, \&c.$ are
 semidiameters of the cube. t thence is $XN = IH\ t\ 15.\ 13.$
 or KN , u and so $XV = KP$; wherefore because $u\ 1.\ ax.\ 1.$
 RVX , is a x right angle, z thence $RXq = XVq + x\ 29.\ 1.$
 $RVq\ (NPq) = KPq + NPq\ a = 3\ KNq\ b = AXq\ z\ 47.\ 1.$
 or $DXq, \&c.$ therefore RX, AX, DX , and by the $a\ 4.\ 13.$
 same reason XS, XT, AX , are equal one to another. $b\ 15.\ 13.$
 And if by the same method, whereby the penta-
 gone $DTCSR$ was made, twelve like penta-
 gones, touching the twelve sides of the cube, be
 made, they shall compose a Dodecaedron; and a
 sphere passing by their angular points, whose
 radius is AX or RX , shall comprehend that Do-
 decaedron. *Which was to be done.*

Lastly, because $KN.NO\ c :: NO.OK$, d thence $c\ constr.$
 $KL.OP :: OP.OK + PL$. Therefore if AB the dia- $d\ 15.\ 5.$
 meter of the sphere be supposed $\bar{p}$, then shall

KL

- e 15. 13. KL $e = \sqrt{\frac{AB}{3}}$ f be also p. g whence OP or
 f sch. 12. RS the side of the dodecaedron shall be a residual line. Which was to be dem.
 10.
 g 6. 13.

Coroll.

From this demonstration it follows, 1. that if the side of a cube be cut in extreme and mean proportion, the greater segment shall be the side of the dodecaedron inscribed in the same sphere.

2. If the lesser segment of a right line, cut in extreme and mean proportion, be the side of the dodecaedron, the greater segment shall be the side of the cube inscribed in the same sphere.


3. It is manifest also, that the side of the cube is equal to the right line which subtends the angle of a pentagone of the dodecaedron, inscribed in the same sphere.

P R O P. XVIII.

To find out the sides of the precedent five figures, and compare them together.

Let AB be the diameter of the sphere given, and AEB the semicircle, and let AC $a = \frac{1}{2} AB$, and AD $b = \frac{1}{3} AB$ then erect the perpendiculars CE, DF, and BG = AB.

join AF, AE, BE, BF, CG; and let fall the perpendicular HI from H; and CK being taken equal to CI, from K erect the perpendicular KL, and join AL. Lastly, c make AF. AO :: AO, OF.



a 10. 1.
 b 10. 6.

c 30. 6.

There-

Therefore 3. 2 $d :: AB. BD$ $e :: ABq. BFq$ the d *constr.*
 side of a Tetraedron and 2. 1 $:: a AB. AC :: ABq. e$ *cor.* 8. 6.
 $BEq. f$ the side of an Octaedron. f 14. 13.

Also 3. 1 $d :: AB. AD$ $e :: ABq. AFq. g$ the g 15. 13.
 side of an Hexaedron. h *constr.*

Moreover, because $AF. AO$ $b :: AO. OF.$ kk *cor.* 17.
 thence shall AO be the side of a Dodecaedron. 13.

Lastly, $BG, (2 BC.) BC$ $l :: HI. IC.$ m therefore l 4. 6.

$HI = 2 CI$ $n = KI.$ therefore HIq $o = 4 CIq.$ m 14. 5.

q consequently CHq $p = 5 CIq.$ q therefore n *constr.*

$ABq = 5 KIq.$ r therefore KI or HI is a ra- o 4. 2.

dius of a circle enclosing the pentagone of an p 47. 1.

Icosaedron; and AK or IB r is the side of a de- q 15. 5.

cagone inscribed in the same circle, f whence r *cor.* 16.

AL shall be the side of a pentagone, t and also 13.

the side of an Icosaedron. Whereby it appears f 10. 13.

that BF, BE, AE are δ $\square$. and AL, AO δ $\square$, t 16. 13.

and $BF \sqsubset BE$, and BE, AF , and $AF \sqsubset AO.$ And u 1. *ax.* 1.

because 3 $AFq = ABq$ $u = 5 KLq$, and $AF \times x$ 4. *ax.* 1.

$AO \sqsubset AF \times OF.$ x and so $AF \times AO + AF \times y$ 1. 2.

$OF \sqsubset 2 AF \times OF,$ r that is, $AFq \sqsubset z 2 AOq.$ z 17. 6.

thence shall 3 AFq ($5 KLq$) be $\sqsubset 6 AOq,$ a 47. 1.

consequently $KL \sqsubset AO$, and much rather AL

$\sqsubset AO.$

That we may express the sides in numbers; If
 AB be supposed $\sqrt{60}$, then, reducing what is
 already shewn to supputation, $BF = \sqrt{40}$, and
 $BE = \sqrt{30}$, and $AF = \sqrt{20}$. Also $AL = \sqrt{30}$
 $-\sqrt{180}$ (for $AK = \sqrt{15} - \sqrt{3}$, and $KL (HI)$
 $= \sqrt{12}$;) Lastly $AO = \sqrt{30} - \sqrt{500}$ ($\sqrt{25}$
 $-\sqrt{5}$.)

Schol.

It is very apparent that besides the five aforesaid figures, there cannot be described any other regular solid figure (viz. such as may be contained under ordinate and equal plane figures.)

For three plane angles at least are required to the constituting of a solid angle; *a* all which must be less than four right angles, *b* but 6 angles of an equilateral triangle, 4 of a square, and 6 of a hexagon, do severally equal 4 right angles; and 4 of a pentagon, 3 of a heptagon, 3 of an octagon, &c. do exceed 4 right angles: Therefore only of 3, 4, or 5 equilateral triangles, of 3 squares or 3 pentagones, it is possible to make a solid angle. Wherefore besides the five above-mentioned, there cannot be any other regular bodies.

a 21. II.

b See schol.

32. I.

Out of P. Herigon.

The Proportions of the sphere and the five regular figures inscribed in the same.

Let the diameter of the sphere be *z*. then shall

The Periphery or circumference of the greater circle, be 6, 28318.

The superficies of the greater circle, 3, 14159.

The superficies of the sphere, 12, 56637.

The solidity of the sphere, 4, 1879.

The side of the Tetraedron, 1, 62299.

The

The superficies of the tetraedron, 4, 6188.

The solidity of the tetraedron, 0, 15132.

The side of the hexaedron, 1, 1547.

The superficies of the Hexaedron, 8.

The solidity of the hexaedron, 1, 5396.

The side of the Octaedron, 1, 41421.

The superficies of the octaedron, 6, 9282.

The solidity of the octaedron, 1, 1333.

The side of the Dodecaedron, 0, 71364.

The superficies of the dodecaedron, 10, 51462.

The solidity of the dodecaedron, 2, 78516.

The side of the Icosaedron, 1, 05146.

The superficies of the Icosaedron, 9, 57454.

The solidity of the Icosaedron, 2, 53615.

Schol.

It is very apparent that besides the five aforesaid figures, there cannot be described any other regular solid figure (viz. such as may be contained under ordinate and equal plane figures.)

For three plane angles at least are required to the constituting of a solid angle; *a* all which must be less than four right angles, *b* but 6 angles of an equilateral triangle, 4 of a square, and 6 of a hexagon, do severally equal 4 right angles; and 4 of a pentagon, 3 of a heptagon, 3 of an octagon, &c. do exceed 4 right angles: Therefore only of 3, 4, or 5 equilateral triangles, of 3 squares or 3 pentagones, it is possible to make a solid angle. Wherefore besides the five above-mentioned, there cannot be any other regular bodies.

a 21. 11.
b See schol.
32. 1.

Out of P. Herigon.

The Proportions of the sphere and the five regular figures inscribed in the same.

Let the diameter of the sphere be *z*. then shall

The Periphery or circumference of the greater circle, be 6, 28318.

The superficies of the greater circle, 3, 14159.

The superficies of the sphere, 12, 56637.

The solidity of the sphere, 4, 1879.

The side of the Tetraedron, 1, 62299.

The

EUCLIDE's Elements.

337

The superficies of the tetraedron, 4, 6188.

The solidity of the tetraedron, 0, 15132.

The side of the hexaedron, 1, 1547.

The superficies of the Hexaedron, 8.

The solidity of the hexaedron, 1, 5396.

The side of the Octaedron, 1, 41421.

The superficies of the octaedron, 6, 9282.

The solidity of the octaedron, 1, 33333.

The side of the Dodecaedron, 0, 71364.

The superficies of the dodecaedron, 10, 51462.

The solidity of the dodecaedron, 2, 78516.





The side of the Icosaedron, 1, 05146.

The superficies of the Icosaedron, 9, 57454.

The solidity of the Icosaedron, 2, 53615.

The thirteenth Book of, &c.

If five equilateral and equiangular figures, like these in the schemes beneath, be made of Paper, and rightly folded, they will represent the five regular bodies.




The End of the thirteenth Book.

THE

THE FOURTEENTH BOOK OF EUCLIDE'S ELEMENTS

PROP. I.




A Perpendicular line DF drawn from D the center of a circle ABC to BC the side of a pentagone inscribed in the said circle, is the half of these two lines taken together, viz. of the side of the hexagone DE , and the side of the decagone EC

inscribed in the same circle ABC .

Take $FG = FE$, and draw CG : a Then CE a 4. 1.
is $= CG$, therefore the angle CGE b $= CEG$ b 5. 1.
b $= ECD$. therefore the angle ECG c $= EDC$ c 32. 1.
d $= \frac{1}{2} ADC$ e $= \frac{1}{2} CED$ ($\frac{1}{2} ECD$) f consequent- d hyp. and
ly the angle $GCD = ECG = EDC$. g wherefore 33. 6.
 $DG = GC$ (CE.) therefore $DF = CE$ (DG) + $EF =$ e 20. 3.
 $DE + CE$. f 7. ax.
2 Which was to be dem. g 6. 1.

PROP. II.



If two right lines AB, DE , be cut according to extreme and mean proportion ($AB. AG :: AG. GB.$ and $DE. DH :: DH. HE.$) they shall


be cut after the same manner, viz. into the same proportions ($AG. GB :: DH. HE.$)

Y 2

Take

- a 17. 6. Take $BC=BG$; and $EF=EH$. Then $AB \times BG$
 b 8. 2. is $a=5 AGq$. wherefore $ACq^b=4 ABG+AGq$
 c 2. ax. 1. $c=BGq$. In like manner shall DFq be $=5 DHq$.
 d 22. 5. & d therefore $AC. AG :: DF. DH$. whence by ad-
 22. 6. dition $AC+AG. AG :: DF+DH. DH$. that
 e 22. 5. is, $2 AB. AG :: 2 DE. DH$. e consequently $AB.$
 f 17. 5. $AG :: DE. DH$; f whence by division $AG.$
 $GB :: DH. HE$. Which was to be dem.

PROP. III.



a sch. 47. 1.

b 30. 6.

c 47. 1.

d 4. 2.

e 10. 13.

f 1. and

3. ax. The same circle ABD comprehends both $ABCDE$
 g 8. 13. the pentagone of a Dodecaedron, and LMN the trian-
 h 2. 13. & gle of an Icosaedron inscribed in the same sphere.

16. 5. Draw the diameter AG . and the right lines
 k 22. 6. & AC, CG . and let IK be the diameter of the
 sphere, a and $IKq = 5 OPq$. b and make OP .


4. 4. $OQ :: OQ. QP$. Because $ACq + CGq^c = AGq$
 l 15. 13. $d = 4 FGq$; and $ABq^e = FGq$. f thence ACq
 m constr. $+ABq = 5 FGq$. moreover, because $CA. ABg ::$
 n cor. 16. $AB. CA - AB$; and $OP. OQ :: OQ. QP$. h and
 13. so $CA.OP :: AB.OQ$. k therefore $3 ACq (/IKq)$

o 12. 13. $5 OPq (m IKq) :: 3 ABq. 5 OQq$. therefore 3
 p 10. 13. $ABq = 5 OQq$. But because ML^n is the side
 q 15. 5. of a pentagone inscribed in a circle, whose ra-
 * before dius is OP , thence $15 RMq. o = 5 MLq p = 5$

r 1. ax. 1. $OPq + 5 OQq = * 3 ACq + 3 ABq q = 15$
 and sch. FGq . r therefore $RM = FG$. f and consequently
 48. 1. the circle ABD is $=$ to the circle LMN . Which
 f 1. def. 3. was to be dem.

PROP.

PROP. IV.



If from F the center of a circle encompassing the pentagone of a dodecaedron $ABCDE$, a perpendicular line FG be drawn to one side of the pentagone CD ; the rectangle contained under the said side CD and the perpendicular FG , being thirty times taken, is equal to the superficies of a Dodecaedron. Also,

If from the center L of a circle inclosing the triangle of an Icosaedron HIK , a perpendicular line LM be drawn to one side of the triangle HK , the rectangle contained under the said HK , and the perpendicular LM , being thirty times taken, shall be equal to the superficies of an Icosaedron.

a 8. 1.

Draw FA, FB, FC, FD, FE . a then shall the triangles CFD, DFE, EFA, AFB, BFC be equal. but b 4. 1.
 $CD \times FG$ b = 2 triangles CFD . therefore 30 c 15. 5.
 $CD \times GF$ c = 60 CFD d = 12 pentagones d 6. ax.
 $ABCDE$ e = to the superficies of a dodecaedron. e 17. 3.
 Which was to be demr. f 41. 1.

Draw LI, LH, LK ; then $HK \times LM$ f is = g 15. 5.
 2 triang. LHK . therefore 30 $HK \times LM$ g = 60 h 16. 13.
 HLK = 20 HIK h = to the superficies of an Icosaedron. Which was to be dem.

Coroll.


k 15. 5.

$CD \times FG. HK \times LM$ k :: the superficies of a dodecaedron to the superficies of an Icosaedron.

Y 3

PROP.

PROP. V.



The superficies of a Dodecaedron has to the superficies of an Icosaedron inscribed in the same sphere, the same proportion that H the side of a cube has to AD the side of an Icosaedron.


Let the circle $ABCD$

a 3. 13. a inclose both the pentagone of a dodecaedron, and the triangle of an Icosaedron; whose sides are BD , AD . upon which from the center E let fall the perpendiculars EF , EGC , and draw CD .

b 9. 13. Because $EC \perp CD$. $EC \ b :: EC. CD$. thence
 c 1. 14. EG ($c \frac{1}{2} EC + CD$) EF ($d \frac{1}{2} EC$) $e :: EF. EG$
 d cor. 12. $- EF. (\frac{1}{2} CD)$ but $H. BD \ f :: BD. H - BD, g$
 $13.$ therefore $H. BD :: EG. EF$. consequently $H \times$
 e 15. 5. $EF = BD \times EG$. wherefore since $H. AD \ b ::$
 f cor. 17. $H \times EF. AD \times EF. k$ thence shall be $H. AD ::$
 $13.$ $BD \times EG. AD \times EF \ l ::$ the superficies of a dodecaedron to the superficies of an Icosaedron.
 g 2. 14. Which was to be dem.
 h 1. 6.
 k 7. 5.
 l cor. 4. 14.

PROP.

PROP. VI.



If a right line AB be cut in extreme and mean proportion, then as the right line BF , containing in power that which is made of the whole line AB , and that which is made of the greater segment AC , is to the right line E containing in power

that which is made of the whole line AB , and that which is made of the lesser segment BC ; so is the side of the cube BG to the side of an Icosaedron BK inscribed in the same sphere with the cube.

In the circle, whose semidiameter is AB , inscribe $BFGHI$ the pentagone of a dodecaedron, and BKL the triangle of an Icosaedron, a where- a cor. 17. fore BG shall be the side of a cube inscribed in 13. the same sphere. therefore $BKq\ b = 3\ ABq$; and $b\ 12.\ 13.$ Eq $c = 3\ ACq$. therefore $BKq.$ Eq $d :: ABq. c\ 4.\ 13.$ $ACqe :: BGq. BFq$. wherefore by inversion $BGq. d\ 15.\ 5.$ $BKq :: BFq.$ Eq. f whence $BG. BK :: BF.$ E. e 2. 14. Which was to be dem. f 22. 6.

PROP. VII.


A Dodecaedron is to an Icosaedron, as the side of a Cube is to the side of an Icosaedron, inscribed in one and the same sphere.

Because a the same circle comprehends both a 3. 14. the pentagone of a dodecaedron, and the triangle of an Icosaedron, b the perpendiculars drawn b 47. 1. from the center of the sphere to the planes of the pentagone and triangle, shall be equal one to another. Therefore if the Dodecaedron and Icosaedron be conceived divided into pyramids, right lines being drawn from the center of the

isphere to all the angles, the altitudes of all the pyramides shall be equal one to the other.
 c 5. and 6. Wherefore since the pyramides of equal height with the bases, and the superficies of the dodecaedron is equal to twelve pentagones, and the superficies of the Icosaedron to twenty triangles, the dodecaedron shall be to the Icosaedron, as the superficies of the dodecaedron is to the superficies of the Icosaedron, *d* that is, as the side of the cube is to the side of the Icosaedron.

d 5. 14.

P R O P. VIII.



The same circle *BCDE* comprehends both the square of the cube *BCDE*; and the triangle of the octaedron *FGH* inscribed in one and the same sphere.


Let *A* be the diameter of the sphere. Because
 a 15. 13. $Aq a = 3 BCq b = 6 BIq$; and also $Aq c = 2$
 b 47. 1. $GFq d = 6 KFq$; thence shall *BI* be $= KF$. *e*
 c 14. 13. therefore the circle *CBDE* $= GFH$. Which
 d 22. 13. was to be demonstrated.
 e 2. def. 1.

The End of the fourteenth Book.

THE

THE FIFTEENTH BOOK OF EUCLIDE'S ELEMENTS.

PROP. I.




IN a cube given $ABGHDCFE$ to describe a pyramide $AGEC$.

From the angle C draw the diameters CA, CG, CE ; and connect them with the diameters AG, GE, EA . All which are a equal among themselves, as being the diameters of equal squares: therefore the triangles CAG, CGE, CEA, EAG are equilateral and equal; and consequently $AGEC$ is a pyramide, which insits upon the angles of the cube, and therefore b is inscribed in it. Which was to
 be done. a 47. I.
b 31. def.
II.

PROP.

PROP. II.

In a pyramid given $ABDE$ to describe an octaedron $EFGKIFH$.



a 10. I.

a Bisect the sides of the pyramid in the points E, I, F, K, G, H , which join with the right lines $EF, FG, GE, \&c.$ All these are

b 4. II. b equal one to the other; consequently the 8 triangles $EHI, IHK, \&c.$ are equilateral and equal, and so make c an octaedron described in the given pyramid. Which was to be done.


c 27. def.

15.

d 31. def.

II.

PROP. III.



In a cube given $CHGBDEFA$ to describe an octaedron $NPQSOR$.

* 8. 4.

Connect * the centers of the squares N, P, Q, S, O, R , with the twelve right lines $NP, PQ, QS, \&c.$

a 4. I.

which are a equal among themselves; and so

b 31. and make eight equilateral and equal triangles: 27. def. II. wherefore b the Octaedron $NPQSOR$ b is inscribed in the cube. Which was to be done.

PROP.

PROP. IV.




In Octaedron given
 $ABCDEF$, to inscribe a
 cube.

Let the sides of the
 pyramide $EABCD$,
 whose base is the square
 $ABCD$, be equally bi-
 sected by the right lines
 LM, MN, NO, OL ,
 which are a equal and a 4. 1.
 b parallel to the sides b 2. 6.
 of the square $ABCD$. c 29. def. 1.
 then the quadrilateral

$LMNO$ is a square. In like manner, if the
 sides of the square $LMNO$ be equally bisected
 in the points G, H, K, I , and GH, HK, KI, IG
 connected, $GHKI$ shall be a square. And if in
 the other 5 pyramids of the octaedron, the cen-
 ters of the triangles be in the same sort conjoin-
 ed with right lines, then other squares will
 be described like and equal to the square
 $GHKI$. wherefore six such squares shall make a
 cube, which shall be described within an octa-
 edron, d being its eight angles touch the d 31. def.
 eight bases of the octaedron in their centers. II .
 Which was to be done.

PROP.



In an Icosaedron given to inscribe a Dodecaedron.

Let ABCDEF be a pyramide of the Icosaedron, whose base is the pentagone ABCDE; and the centers of the triangles G, H, I, K, L; which connect with the right lines GH, HI, IK, KL, LG. Then GHIKL shall be a pentagone of the dodecaedron to be inscribed.

For the right lines, FM, FN, FO, FP, FQ, passing by the centers of the triangles, *a* do equally divide their bases into two parts. *b* therefore the right lines MN, NO, OP, PQ, QM *c* are equal one to the other; *d* whence also the angles MFN, NFO, OFP, PFQ, QFM are equal. therefore the pentagone GHIKL is equiangular. *e* and consequently equilateral, being *e* 4. 1. FG, FH, FI, FK, FL *f* are equal. And if in *f* 12. 13. the other eleven pyramids of the Icosaedron, the centers of the triangles be in like sort conjoined with right lines, then will pentagones, equal and like to the pentagone GHIKL, be described. Wherefore 12 of such pentagones shall constitute a dodecaedron; which also shall be described in the Icosaedron, seeing the twenty angles of the dodecaedron consist upon the centers of the twenty bases of the Icosaedron. Whereby it appears that we have described a dodecaedron in an Icosaedron given. *Which was to be done.*

EUCLIDE'S



EUCLIDE'S DATA.

A

*Commentary or Preface written by
the Philosopher MARINUS, on
EUCLIDE'S DATA.*

IN the first place we ought to set down (as a Foundation) what that is, which we call *DATUM* or GIVEN; then to consider the Profit and Utility thereof; and in the third place, to what Art or Science this Tract doth appertain.

The Word *DATUM* therefore is diversly defined, for the Antients have defined it after one manner, and later Writers after another, whence it follows that it seemeth a difficult thing to give a true Explication thereof; for some of them have not delivered the Definition of the Word; but have with much Labour and Trouble sought certain Proprieties thereof, and some others collecting and mingling what hath been delivered

delivered by others before, have endeavored to define the Word *DATUM*; but not so exquisitely but that they have contradicted themselves; altho' what hath been said by all of them, seems to be grounded on one and the same notion and supposition; for they all take the Word *DATUM* to be a thing comprised; and therefore among such as have endeavored to describe it most simply, and with some simple difference, some of them have taken the Word *DATUM* to be the same with *ORDINATUM*, and so *Apollonius* understands it in his Tract of Inclinations, and in his universal Tract; and so others, as *Diodorus* takes it to be *COGNITUM KNOWN*; for in this Signification he takes the right line and the angles to be given, and all that may arrive to our Knowledge, altho' we may not be able well to express it. But others have believed that it hath the same signification as the Word [*Effabile*] that may be declared, and so *Ptolomy* would have it, who calls those things *GIVEN*, whose measure is known whether precisely, or near the matter. Others also have thought the Word *DATUM* to be what is granted us by the Proposer in the Hypothesis; being that in the first Elements, a point given, and a right line given, is diversly taken (that is to say, that who so would give and determine the quantity of a right line) all which things signify some *COMPREHENSION*; and therefore of all these Definitions, those are most agreeable, which do most openly declare the *COMPREHENSION*, as we shall make evident by what follows.

Let us now unfold the diverse Opinions of those, who writing the nature of *DATUM GIVEN*, have not taken one simple Mark, or only Character for its Definition; and let us reduce it as in a Summary or Epitomy, to the

end

end we may with the more ease know or number all their Differences. Some of them then have defined *DATUM* to be *Ordinatum* and *Porimon* together, and others *Ordinatum* and *Cognitum* together, and others *Porimon* and *Cognitum* together. Wherefore all seem to have so defined it as to have had regard to the *Comprehension*, or *Assuming* and *Invention* of the thing given; and to the end that we may the better conceive their Opinions, and that from the saying of many we may be able to draw a true Definition of what is proposed, we will take notice in the first place of the Signification of all the simple Terms which they make use of, as also of the Terms opposed to them, to wit, *Inordinatum* and *Incognitum*, *Aporon* and *Irrational*; for those things appertain to this Geometrical Business, to natural things, and to Mathematical Discipline.

Now we may call that *Ordinatum* (or Regulated) which doth always keep and observe that for which it is said to be ordered, whether you regard its Magnitude or Species, or touching some other such like thing: It is also thus defined, *Ordinatum* is that which cannot be done in divers manners, but in one only manner, and in some determined place: As for example, A right line drawn by two given points, is said to be ordered, by reason it cannot be otherwise done, nor in divers manners. But an angle passing by two points is said to be *Inordinatum* (or disordinate and irregular) for that it is made in infinite and divers manners by a great or small circle described by two points *ad infinitum*. Contrariwise, an angle constituted by three points, is said to be *Ordinatum*, as also those things which follow are said to be *Ordinatum*, as to constitute an equilateral triangle on a right line, for it cannot be diversly made, but unchangeably, on both the given

extremities of the line. Again, To divide a given right line according to a given proportion, for that cannot be done but in one certain point. The things *Inordinatum* are such as are done contrary to those last mentioned, as to constitute a Scalene triangle, and to divide a right line indefinitely. Wherefore the Problem is ordered, is proposed in the determination, considering that a certain thing may be in one manner said to be *Ordinatum*, and *Inordinatum* in another, as an equilateral triangle, considering the equality of the sides, it is *Ordinatum*, but considering its Magnitude it is *Inordinatum*, being in no wise determined.

But we call that *Cognitum* which is notorious, as clear and comprehended of us, and *Incognitum* that which is not known, or comprehended of us, as the length of a way is said to be known, when we know how many Miles it contains; also that the three lines of a Rectiline triangle are equal to two right angles; and in like manner that the Binomial is Irrational, such things are known, as also that it is only one right line that can touch a spiral line from a given point without it, from both parts; for if there were yet another line, two right lines would enclose a space, which is impossible. Again, irrational things are not said to be unknown, but such of them only which are neither known nor comprehended of us.

Porimon is that which we may make and constitute, that is to say, bring to our Understanding. Again, it is defined thus, *Porimon* is that which may be exhibited by Demonstration, or which is apparent without Demonstration, as to describe a circle from a center and with a space, as also to constitute not only an equilateral triangle, but also a Scalene; or to find a *Binomium*, or to find two right lines rational, commensurable in power only, and other things

things w
as to def
Aporon
example
it hath
certainly
the man
this pre
of that
called
been as
ble, is
Constr
hath
and is
ded, n
Effa
and ex
cies,
but th
prope
that v
acco
of a
TH
may
that
ther
all
ther
the
of
oth
ext
ma
po
by
is
m
th

things which are known infinitely, are *Porimon*; as to describe a circle by two points.

Aporon is wholly opposite to *Porimon*, as for example, the Quadrature of a circle, for that it hath not as yet been found; altho' it be certainly known that it may be: Nevertheless the manner of finding it out hath not been to this present comprehended. But we speak here of that which is already known, which is called *Porimon principale*; for what hath not been as yet made, and yet nevertheless is possible, is called *Poriston*, (or feasible) altho' the Construction be yet unknown. But *Aporon*, as hath been afore said, is opposite to *Porimon*, and is that whose Nature is not as yet decided, nor well determined.

Effabile, that is to say, rational (or speakable and explicable) is that whose Magnitude, Species, and Position, we may be able to declare; but this Definition is a little too general, for properly, and according to it self, *Effabile* is that which is known by certain things, and according to a Measure given by Position, as of a span, or a finger's breadth, &c.

These things then being thus unfolded, we may easily perceive in what all those things that we have afore spoken of do agree together, and wherein they do differ; and first of all how *Ordinatum* and *Cognitum* do agree together, and likewise their opposites the one to the other, for it cannot be said that any one of those things by counterchanging is the other, nor yet that the one hath not more extent than the other, altho' they agree in many things, as to describe a right line by two points, and to constitute an equilateral triangle by three circles. But to square a circle, that is indeed *Ordinatum*, yet nevertheless, *Incognitum*. Also that at a point of a spiral line there is but one touch line, that is, of the

kind we call *Ordinatum*, and cannot be otherwise done, yet nevertheless the Demonstration and Construction thereof is not yet known. Again, the indefinite section, and the Construction of the Scalenum is *Cognitum*; but is not *Ordinatum*; inasmuch as it is manifest, that amongst those things which are *Ordinatum*, there are some that are *Cognitum*, and others that are *Incognitum*; and contrariwise, that amongst the things that are *Cognitum*, there are some that are *Ordinatum*, and others that are *Inordinatum*; and therefore those things answer one another, as among Living Creatures, that which hath reason, with that which hath Feet, for there is no equality amongst them, neither doth the one extend more than the other.

In like manner, *Ordinatum* and *Inordinatum* agree together, respecting *Porimon* and *Aporon*; seeing that between them there is a very great Resemblance, and because that they do differ only in the manner before expressed; for in truth the spiral line is *Ordinatum*; but it was not *Porimon* before *Archimedes*; and by the same reason those things that are inordinate and known by an infinity of ways and means are *Porimon*, if any one shall undertake to invent their Constitution and Construction. Yet nevertheless they are not ordinate, as to constitute a Scalenum triangle, it being no difficult thing to make known the Constitution thereof by an equilateral triangle, yea it is most easy, altho' it be inordinate and known by an infinity of ways.

And in the same manner do agree *Ordinatum* and *Inordinatum*, together with *Effabile* and *Irrationale*; for they agree together in many things, differing nevertheless by the foregoing reason, seeing those things there mentioned bear no equality to each other, neither doth one thing

thing con-
such as
ordinate
pressible.
the squ
touching
because
minatel
sured b
explica

Now
is betw
difficul
rence;
come
seems
verthe
him
let it
line
point
the P
yet c
is Co
that
there
exter

N
agre
othe
for
nals
are
eve
nun
wa
for
th
ou
B

thing contain the other; for all Binomiums and such as are taken as Irrationals, are indeed ordinate, but yet they are not *Effabile*, or expressible, or to be unfolded, as the diameter of the square is in respect of its side. Now touching *Effabile*, there are divers inordinate; because they are diversly known, and indeterminately, for a Scalenum triangle may be measured by a defined and proposed measure, as explicable, altho' it be inordinate.

Now it is easy to see the agreement that there is between *Cognitum* and *Porimon*, but it is a difficult thing to express or unfold their difference; forasmuch as in their Natures they come so near to one another, as that there seems to be an equality between them: Nevertheless, there will some difference appear to him that shall consider it more strictly; for let it be consented to that there can be only one line that can touch a spiral line in a certain point, that is *Cognitum*, yet notwithstanding the Problem is not *Porimon*, it being not as yet comprehended; so as that all that which is *Cognitum*, is not therefore *Porimon*. But all that which is *Porimon* is also *Cognitum*, and therefore *Cognitum* appears to be of a greater extent than *Porimon*.

Now *Cognitum*, *Porimon*, and *Effabile*, do agree in some certain things, and do differ in other things by the same reason before alledged; for those lines which are there called Irrationals are in truth known; and yet nevertheless are not *Effabile* or explicable. Contrariwise, every number is indeed *Effabile*, and yet every number is not *Cognitum*. But *Effabile* is always of its own nature expressible, altho' that some lengths may be now *Effabile*, and at another time not, if it be examined with some other according to one and the same measure. But also that same length is sometimes known,

and other times not, tho' they wholly agree with one another. Now it is a difficult thing to find something that shall be *Effabile* and *Incognitum*, for *Cognitum* seems to be of a greater extent than *Effabile*, and by those things it is manifest that *Porimon* and *Aporon* do differ from RATIONAL or *Effabile*, and from IRRATIONAL, for of IRRATIONAL some of them may be *Porimon*; but of RATIONAL none of them can be Irrationals; and therefore it is very easy to perceive in what the before expressed things agree. Notwithstanding they seem to agree together, in such sort as that *Porimon* seems to be of a greater extent than *Effabile*.

Now by these things we may come to know the difference of those things that have been before spoken of, for in truth *Effabile* and *Irrationale* are so termed in respect of measure, which notwithstanding is not as yet arrived to our Understanding, seeing that something that is rational, may be as yet unknown to us, and in like manner may be rational, and yet may never be comprehended so to be. But *Ordinatum* and *Inordinatum* is so termed according to it self, and according to the proper nature of the thing on which we contemplate, altho' it be not comprehended by us. As *Archimedes* had perceived some things to be ordinate from the nature of the things, the which *Serenus* had before contemplated. But *Cognitum* and *Incognitum* is spoken in respect of us, so as the things before mentioned do differ among themselves; for these have respect to us, the others, some of them to their proper nature, and the rest to measure.

Having then explained the agreements and differences of the things that have been proposed, it remains now that we consider what is meant by the Word *DATUM*, for of all those

those th
that wh
the Hyp
because
GIVEN
GIVEN
sis, as r
been ma
fore wa
Definit
Ther
ed in t
consequ
such a
Word,
of the
in the
to ha
to wit
Word
and a
define
other
to o
thing
right
Defin
it o
thin
goo
Defin
the
ned
wh
tog
tha
cie
D
na
in

those that believe the Word *DATUM* to be that which is consented to by the Proposer in the Hypothesis, are wide from what is sought; because that all the Elements of the things GIVEN are not composed of this sort of GIVEN, which is according to the Hypothesis, as may be seen in those Tracts which have been made on this Subject GIVEN. Wherefore waving this Opinion, let us judge of the Definitions of others.

Then, that which is consented to, or granted in the Hypothesis, is something which is consequently known by the Principles; but such as make use of Definitions of one only Word, do define it and remark it by some one of the before mentioned, as hath been spoken in the beginning, so as that almost all seem to have had this common notion of GIVEN, to wit, that it is comprehended even as the Word *DATUM*, doth also manifest it to be; and amongst those, these are the chief that do define it by the Hypothesis or Supposition; and others have had regard to what is consented to or granted. But we making use of the said things as of a Rule and Direction to judge rightly, we may be able to find out a perfect Definition of *DATUM*; for it is certain that it ought to equal and be convertible with the thing defined, which is one thing proper to good Definitions. Now such seems to be the Definition of the thing proposed, which among the most simple and plain Expositors is defined *Porimon*, and amongst the more acute, that which defineth it to be *Porimon* and *Cognitum* together; but all the rest are imperfect; for that which defineth it *Ordinatum* is not sufficient for the Comprehension and Knowledge of *DATUM*; because that neither wholly ordinate nor alone ordinate, is not comprised, seeing that there are things inordinate that have

the same Condition, as hath been shewn. Again, that reason gives not Satisfaction neither, which describes it to be *Cognitum*; for all that is known is not comprehended, altho' that alone *Cognitum* be comprehended. Moreover, that also is not perfect which defineth it to be *Effabile*, for *Effabile* is not alone comprehended; seeing that some of the Irrationals are also comprehended. In like manner, all *Effabile* is not comprehended, as hath been before declared. Now amongst the Definitions which expound it by the only Word, there remains that which defineth it to be *Porimon*, which seemeth greatly to manifest the Comprehension; for whole *Porimon* and alone *Porimon* is comprehended. Wherefore EUCLIDE himself useth such a Definition in a Description of all the kinds of GIVEN by him conceived and regarded. But amongst such Definitions as are compounded, that is a perfect Definition which defineth *DATUM* to be *Cognitum* and *Porimon* together, having *Cognitum* for analogical kind, and *Porimon* for difference; but that is imperfect which hath *Ordinatum* and *Porimon* together; for those things which are such, are not alone GIVEN, and that which defineth it *Ordinatum* and *Effabile* together, comprehendeth likewise the GIVEN, with the defect or want. But that of *Cognitum* and *Ordinatum* together, is not to be received or admitted, because it doth exceed what is defined, for such is not given alone: Therefore those only which have declared that *DATUM* is *Cognitum* and *Porimon* together, seem to have attained the notion of GIVEN, for that which is such is all, and alone comprehended, which two things ought to be in those Definitions that are well given. But the former comes near to those which have thus defined it: *DATUM* is that to which we may find an equal, according to those things

we

we have
Hypoth
making
which
leaves
some o
that in
TUM i
kinds
GEOM
before

Wha

ar

or

A
prefer
It fo
Now
orda
for s
to a
have
Stre
ther
Can
wel
De
it
De
cel
to

we have proposed in the first Principles and Hypothesis, of which number *EUCLIDE* is one, making use throughout of the Word *περιόχου*, which signifies to exhibit or invent, altho' he leaves *Cognitum* as a Consequent of *Porimon*; some one nevertheless might reprove him, for that in the first place he hath not defined *DATUM* in general; but immediately some of the kinds of GIVEN, altho' in his Elements of GEOMETRY, he hath defined the line simple before the Species or Kinds.

What is the Utility and Profit that ariseth from this Tract of DATA, or things GIVEN.

AFTER having explained universally, and according to what seemed necessary for our present Use, what this Word *DATUM* signifieth: It follows to shew the Utility of this Tract. Now this Tract is such, as that it is not only ordained and instituted for its own respect, but for some other thing; for it is very necessary to a place which is called Resolved, and we have already declared elsewhere how much Strength a resolved place doth obtain in Mathematical Disciplines, as also in Opticks and Cannons, which come very near to them, as well for that Resolve is an Invention of the Demonstration, as for that in such like things it serves us much for the Invention of the Demonstration, or for that it is much more excellent to meet with a Resolutive Power, than to enjoy divers particular Demonstrations.

To what Art or Science this Tract is referred.

NOW seeing the Consideration of GIVEN is useful and profitable in all these kinds of

of Arts, for that it serveth much to RESOLUTION, it may well be said to be recalled, not only to one only Science, but to the Mathematicks universally which treat of Numbers, Time, Swiftneſs, and ſuch like things, which treateth likewise of Reaſons, as alſo of Proportions, and in a word of all *Medietez*: Wherefore for the perfect and demonſtrative Knowledge of things GIVEN, being of ſo great Utility, *EUCLIDE* hath taken pains to frame this Book of things GIVEN, which Author amongſt all ſuch as have compoſed the Elements of Geometry, hath juſtly deſerved the firſt place and rank, and who having invented the Elements, or rather the Introductions almoſt of all Mathematical Diſciplines, to wit, of all Geometry in 13 Books, of Astronomy, of Muſick, and Opticks, he hath left in Writing the Elements RESOLUTIVE, in this Treatiſe of things GIVEN; but as he was a Geometrician, he hath particularly accommodated to Magnitudes, that was of the GIVEN; yet nevertheless common in other things, which Method hath alſo been obſerved by him, when treating univerſally of Reaſons and Proportions, he appropriates them to the Magnitudes mentioned in his Fifth Book of Planes.

Now it hath been declared in general what is the meaning of *DATUM*, to what Science it appertaineth, and how profitable the Contemplation thereof is. We will add to what hath been ſaid, the Deſcription of this Science which treats of things GIVEN; ſeeing that it is (as appears by what hath been ſaid) a Comprehension in all manners of things GIVEN; and of their Accidents and Proprieties. But having reſpect to the propoſed Book, we ſhall declare it to be an Elementary Doctrine of the whole Knowledge of things GIVEN, whence it follows that it will be
very

very profi
tained, ſe
Now t
Species
in the f
which a
as are g
are giv
is given
larly co
in thoſe
he hath
Reason
are giv
EUCLID
Book, (M
Magnit
circular
obſerve
this Bo
of inſt
poſitio
ampl
Book.

The

very profitable, as also the things therein contained, seeing they refer to things GIVEN.


Now this Book is divided according to the Species or Kinds of the things GIVEN, and in the first Section are contained those things which are given by Reason. Secondly, such as are given by Position: and lastly, such as are given by Species or Kind; for that which is given by Magnitude is simple, and particularly contained in the others, and principally in those things given by Species or Kind. Now he hath begun with those things given by Reason and Position, forasmuch as those that are given by Species are constituted of them. *EUCLIDE* gives yet another Division to this Book, for that he divides it into universal Magnitudes, Lines, and Superficies, and into circular Theoremes, which Order he hath also observed in the Definitions and Suppositions of this Book. Moreover, he useth a certain way of instructing, which proceeds not by Composition, but by Resolution, as *Pappus* hath amply set down in his Commentaries on this Book.

*The End of the PREFACE of
MARINUS.*

DEFL.

DEFINITIONS.

- I. **P** Lanes or Spaces, Lines, and Angles, to which we may find others equal, are said to be given by Magnitude.
- II. A Reason is said to be given, when we may find one of the same or equal thereto.
- III. Rectiline figures, whose angles are given, and also the reason of the sides to one another, are said to be given by Species or Kind.
- IV. Points, Lines, and Angles, which have and keep always one and the same place and situation, are said to be given by Position or Situation.
- V. A Circle is said to be given by Magnitude, when the semidiameter thereof is given by Magnitude.
- VI. A Circle is said to be given by Position, and by Magnitude when the center thereof is given by Position, and the semidiameter by Magnitude.
- VII. Segments of Circles, whose angles and bases are given by Magnitude, are said to be given by Magnitude.
- VIII. Segments of a Circle, whose angles are given by Magnitude, and the bases of the segments by Position and Magnitude, are said to be given by Position and by Magnitude.
- IX. A Magnitude AB, is greater than another Magnitude C, by a given Magnitude BD, when having taken away the given



given
the ot
X. A M

A —
C —

XI. A M

A —
king
nitud
nitud

XII. A

D —

give
who
give
XIII.


a g
tion
ang

XIV.
giv
rig

XV.
Po
a

given Magnitude DB, the rest AD, is equal to the other Magnitude C.

- X. A Magnitude AB, is less than another Magnitude C, by a given Magnitude BD when having added thereto the given Magnitude BD, the whole AD is equal to the other magnitude C.



- XI. A Magnitude AB, is said to be greater than another magnitude CB, by a given magnitude AD, and in reason, when taking from the same magnitude the given magnitude AD, the rest DB, hath to the other magnitude CB, a given reason.



- XII. A magnitude AB is said to be less than another magnitude BC, by a given magnitude AD, and in reason, when the given magnitude AD being added thereto, the whole DB hath to the other magnitude BC, a given reason.




- XIII. A right line is said to be drawn down from a given point, unto a right line given in Position, the right line being drawn in a given angle.

- XIV. A right line is said to be drawn up from a given point, to a right line given in Position, the right line being drawn in a given angle.

- XV. A right line is against another right line in Position, when it is drawn parallel thereto through a given point.

PROPOSITION I.



a 1. def.

b 7. 5.

c 16. 5.


d 2. def.

TWO magnitudes *A* and *B*, being given, the reason they have to one another *A* to *B*, is also given.

Demonstration. For seeing that the magnitude *A* is given, *a* we may find one equal thereto, which let be *C*. Again, forasmuch as the magnitude *B* is given, we may also find one equal to that, and let that be *D*. Therefore seeing that *A* is equal to *C*, and *B* to *D*, as *A* is to *C*, *b* so is *B* to *D*, and by permutation, *c* as *A* shall be to *B*, so *C* shall be to *D*. Therefore *d* the reason of *A* to *B* is given, for it is the same reason as of *C* to *D*, as we have found, and which ought to be demonstrated.

PROP. II.

If a given magnitude *A*, hath to some other magnitude *B*, a given reason, that other magnitude *B*, is also given by magnitude.



a 2. def.

b 14. 5.


c 1. def.

Demonstr. For seeing that *A* is given, we may find one equal thereto, which let be *C*: And forasmuch as the reason of *A* to *B*, is also given, we may find *a* one of the same. Let it be found, and let the reason be of *C* to *D*. Now seeing that as *A* is to *B*, so *C* is to *D*; and by permutation, as *A* is to *C*, so *B* is to *D*: But *A* is equal to *C*, therefore *b* *B* shall be also equal to *D*. Therefore *c* the magnitude *B* is given, seeing that thereto there hath been found one equal, to wit, *D*.

PROP.



one equal
fore see
is equal
the wh
that D



also fi
ing t
the n
the n
equal
for t
wit,



PROP. III.


If given magnitudes AB and BC , are compounded, that magnitude AC , that is compounded of them shall be also given.



Demonstr. For seeing that AB is given, we may find one equal to it, which let be DE . Again, seeing that BC is given we may also find one equal to that, which let be EF . Wherefore seeing that DE is equal to AB , and EF is equal to BC , the whole AC ^a is equal to ^{a 2. ax. 1.} the whole DF . Therefore AC is given, seeing that DF is proposed equal thereto.

PROP. IV.


If from a given magnitude AB , there be taken away a given magnitude AC , the remaining magnitude CB is also given.



Demonstr. Forasmuch as AB is given, we may find one equal thereto, which let be DE . Again, seeing that AC is given, we may also find one equal to it, which let be DF . Seeing then that the magnitude AB is equal to the magnitude DE , and the magnitude AC to the magnitude DF ; the rest CB ^a shall be ^{a 3. ax. 1.} equal to the rest FE . Wherefore CB is given, for to it there hath been found an equal, to wit, FE .

PROP. V.

If a magnitude AB , hath a given Reason to some part thereof AC , it will have also a given reason to the part remaining CB .



Demonstr. Let DE be exposed as a given magnitude, and seeing that the reason of the magnitude AB ,

to

- a 2. def. to the magnitude AC, is given, *a* we may find one of the same, which let be DE to DE; therefore the reason of the same DE to DF is given. But DE being given, so is *b* also its part DF; and consequently, *c* the rest FE: Therefore *d* seeing that DE and FE are given, the reason of the same DE to FE, is also given. And forasmuch as DE is to DF, as AB is to AC, and by conversion, as DE to FE, so AB is to CB. But the reason of DE to FE is given, as hath been demonstrated; therefore the reason of AB to CB is also given.
- 19.5.

Scholium.

From this it is evident that if a magnitude hath to some part thereof a given reason, by division, the reason that one part hath to the other, shall be also given. For seeing that as DE is to FE, so is AB to CB; by division, as DF to FE, so AC to CB. But it hath been demonstrated that the parts DF and FE are given, and consequently their reason is also given: In like manner, therefore the reason of AC to CB is given.

PROP. VI.

A	D
B	E
C	F

If two magnitudes AB and BC, having to one another a given reason, are compounded, the magnitude AC compounded of them, shall also have a given reason to each of them AB and BC.

Demonstr. Let the given magnitude DE be exposed, and seeing that the reason of AB to BC is given; let there be made one and the same of the said DE to EF; therefore the reason of the same DE to EF is given; and therefore *a* the magnitude DE being given, both the one and the other of them DE and FE, is given. Wherefore *b* the whole DF shall be also given. There-

a 2. prop.

b 3. prop.

Therefore
of them I
asmuch th
in compo
to EF;
to is DE
is to ea
EF, so th
tudes AB
the same
and BC

A —
ment AC
Demon
CB is g
them (A
given:
and CB

A
B
C
given
the re
done
D is
that
be do
fore

Therefore *c* the reason of the same DF to each of them DE and EF, shall be given. And forasmuch then as AB is to BC, so is DE to EF; in compounding, *d* as AC is to BC, so is DF to EF; Therefore by conversion, as AC to AB, so is DF to DE. Therefore as the whole DF is to each of the other magnitudes DE and EF, so the whole AC is to each of the magnitudes AB and BC: Therefore *e* the reason of the same AC to each of the magnitudes AB and BC is given. 1. prop. 18. 2. def.

PROP. VII.

C
A ——— B

If a given magnitude AB, be divided according to a given reason AC to CB, each segment AC and CB is given.

Demonstr. For seeing the reason of AC to CB is given, the reason of *a* AB to each of them (AC and CB) is also given. But AB is given: Therefore *b* each of the segments AC and CB is also given. a 6. prop. b 2. prop.

PROP. VIII.

A	D	Magnitudes A and C, which have to one and the same a given reason B, shall be to one another in a given reason, A to C.
B	F	
C	E	
—	—	

Demonstr. For let the given magnitude D be exposed, and seeing that the reason of A to B is given, let the same be done of the said D to E. Now seeing that D is given, *a* E is also given. Again, seeing that the reason of B to C is given, let the same be done of E to F. But E is given, and therefore F is also given. But seeing that D is given,

a 2. prop.

- b 1. *prop.* given, b the reason of the same D to F is given; and seeing that as A to B, so D to E, and as B to C, so is E to F; in reason of equality, c as A is to C, so is D to F; but the reason of D to F is given. Therefore the reason of A to C is also given.

PROP. IX.




If two or more magnitudes A, B, and C, are to one another in a given reason, and that the same magnitudes A, B, and C, have to other

magnitudes D, E, and F, given reasons, altho' they be not the same, those other magnitudes D, E, and F, shall be also to one another in given reasons.

- Demonstr.* Forasmuch as the reason of A to B is given, as also that of A to D, the reason of D to B shall be given: But the reason of B to E is also given; therefore the reason of the same D to E shall be in like manner given. Again, seeing that the reason of B to C is given, and also that of B to E, the reason of E to C shall be given. But the reason of C to F is also given. Therefore a the reason of E to F shall be given. But it hath been demonstrated that the reason of D to E is also given; and therefore b the reason of D to F shall be given. Therefore the magnitudes D, E, and F, are to one another in given reasons.

PROP. X.



If a magnitude AB, be greater than another magnitude BC, by a given magnitude, and in reason, the magnitude

nitide
than the
and in
be gre
magnitu
shall be
magnitu
mainder
other i

Dem
BC by
the g
There
to BC
reason
magni
greate
tude,
Ag
than

A—
than
in re
whic
reaso
F
than
and
tude
less
it i
The
CB
son
AD
AB
giv

magnitude AC compounded of both, shall be also greater than that same magnitude, by a given magnitude, and in reason: But if that compounded magnitude be greater than the same magnitude, by a given magnitude, and in reason; either the remainder shall be also greater than that same by a given magnitude, and in reason; or else the same remainder is given with the following, to which the other magnitude hath a given reason.

Demonstr. For seeing that AB is greater than BC by a given magnitude, and in reason, let the given magnitude AD be taken away. Therefore a the reason of the remainder DB a 11. def. to BC is given; and in compounding, b the b 6. prop. reason of DC to BC is also given. But the magnitude AD is also given; therefore AC is greater than the same BC by a given magnitude, and in reason.

Again, Let the magnitude AC be greater than the magnitude BC, by a given magnitude, and in reason:

I say that the rest AB, is either greater

than the same BC by a given magnitude and in reason, or that the same AB, with that which followeth, to which BC hath a given reason, is given.

Forasmuch as the magnitude AC is greater than the magnitude BC, by a given magnitude, and in reason, cut off from it the given magnitude: Now the same given magnitude is either less than the magnitude AB, or greater: Let it in the first place be less, and let it be AD. Therefore the reason of the remainder DC to CB is given. Wherefore by division, the reason of DB to BC is given. But the magnitude AD is also given; therefore the magnitude AB is greater c than the magnitude BC by a c 11. def. given magnitude, and in reason. Now let the

A a given

given magnitude be greater than the magnitude AB, and let AE be put equal thereto; therefore *d* the reason of the remainder EC to CB is given; and by conversion, *e* the reason of the same BC to BE, is also given. But the same EB with BA is given, for that the whole AE is given: Therefore there is given AB, with that which follows BE, to which BC hath a given reason.

PROP. XI.

E D B

A — 1 — 1 — 1 — C

If a magnitude AB, be greater than a magnitude BC, by a given magnitude, and in reason, the same magnitude AB, shall be also greater than the magnitude compounded of them by a given magnitude, and in reason, and if the same magnitude be greater than the two others together by a given magnitude, and in reason, that same magnitude shall be also greater than the rest by a given magnitude, and in reason.

Demonstr. For seeing that the magnitude AB is greater than BC by a given magnitude, and in reason; let there be taken from it a given magnitude AD: Therefore *a* the reason of the rest DB to DC, is given, and therefore *b* the reason of DC to BD shall be also given: Let the same be done of AD to DE, therefore the reason of the same AD to DE is given. But *c* 2. prop. AD is given, therefore DE is also given, and consequently, *d* 4. prop. the rest AE, is also given. But seeing that as AD is to DE, so is DC to BD; by permutation, *e* as AD is to DC, so is DE to DB: Therefore by compounding, *f* as AC is to CD, so is EB to DB; and by permutation, *g* as AC is to EB, so is DC to DB. But the reason of DC to DB is given: Therefore also is AC to EB, and consequently, that of EB to AC. But it hath been demonstrated that AE is given, therefore *h* AB is greater than AC by a given magnitude, and in reason.

Eut

But no
given m
same AB
a given
For
given m
magnitu
fore i th
given,
that of
AD to
is given
AD to
ly that
therefo
and fe
whole
part c
der D
of AC
be gi
divisi
and c
DB
that
than
in re

A —
AB
But
be
the
dih
tu

But now let AB be greater than AC by a given magnitude, and in reason: I say that the same AB is also greater than the rest BC, by a given magnitude, and in reason.

For seeing that AB is greater than AC by a given magnitude, and in reason. Let the given magnitude AB be cut off there-from: Therefore i the reason of the remainder EB to AC is i *ii. def.* given, and consequently, also shall be given that of AC to EB. Let the same be done of AD to DE, therefore the reason of AD to DE is given; and by conversion, k the reason of k *s. prop.* AD to AE shall be also given, and consequently that of AE to AD. Now AE is given, therefore the whole AD l shall be also given; l *2. prop.* and seeing that as the whole AC is to the whole EB, so the part cut off AD, is to the part cut off ED, so also shall be m the remain- m *19. s.* der DC to the remainder DB. But the reason of AC to EB is given: Therefore also shall be given that of DC to DB. Wherefore by division, n the reason of BC to DB is given; n *schol. s.* and consequently also shall be given that of *prop.* DB to BC. But it hath been demonstrated that AD is given: Therefore o AB is greater o *ii. def.* than the same BC by a given magnitude, and in reason.

PROP. XII.

B C

A — 1 ——— 1 — D


If there be three magnitudes AB, BC, and CD, and that the first AB, with the second BC, to wit AC, be given. But the second BC, with the third CD, to wit, BD, be also given: Either the first AB shall be equal to the third CD, or the one shall be greater than the other by a given magnitude.

Demonstr. Forasmuch as each of the magnitudes AC and BD are given, the given magnitudes

A a 2

tudes


tudes are either equal to one another, or unequal. Let them be first equal: Therefore AC is equal to BD, take away the common magnitude BC, and there will remain a AB, equal to CD. But suppose them to be unequal, as in this second figure, and let BD be greater



than AC: Let then BE be put equal to AC. Now seeing that AC is given, BE is also given. But the whole BD is also given, the rest ED shall be so also; and forasmuch as BE is equal to AC, taking away the common magnitude c BC, there will remain AB equal to CE. But ED is given: Therefore CD is greater than AB by the given magnitude ED.

Scholium.

And if the first with the second, to wit, AC, were greater than the second with the third, to wit, BD,



as in the other figure, CE would be made equal to the same BD, and by the same reasons as was above demonstrated, that AE is given and equal to CD; and therefore AB greater than CD by a given magnitude.

PROP. XIII.



If there be three magnitudes AB, CD, and E, and that the first of them AB, hath a given reason to the second CD; but the second CD is greater than the third E, by a given magnitude, and in reason, also the first AB, shall be greater than

than the third E, by a given magnitude, and in reason.

Demonstr. For seeing that CD is greater than E by a given magnitude, and in reason, let the given magnitude CF be taken there-from: Therefore the reason of the rest FD to E is given. And forasmuch as the reason of AB to CD is given, let the same be done of AH to CF. Therefore the reason of the same AH to CF is given. But CF is given: Therefore a AH is also given. And seeing that as the a 2. prop. whole AB is to the whole CD, so the part cut off AH is to the part cut off CF, and so b also the rest HB is to the rest FD, the rea- b 19. 5. son of the same HB to FD is also given. But the reason of FD to E is also given: There- fore c the reason of HB to E is given. But it c 8. prop. hath been demonstrated that AH is given: Therefore d AB is greater than the said E by d 11. def. a given magnitude, and in reason.

P R O P. XIV.

$$\begin{array}{ccccccc}
 & B & & G & & & \\
 A & \text{---} 1 & \text{---} 1 & \text{---} & E & & \\
 & D & & & & & \\
 C & \text{---} 1 & \text{---} & F & & &
 \end{array}$$

If two magnitudes AB and CD, have to one another a given reason, and that to each of them there be added a given magnitude, to wit, BE and DF; either the whole AE and CF shall have to one another a given reason, or the one shall be greater than the other by a given magnitude, and in reason.

Demonstr. For seeing that each of those magnitudes BE and DF, is given, a the reason of a 1. prop. the said BE and DF is also given; and if that reason be the same with that of AB to CD, that of the whole AE to the whole CF, b b 12. 5. shall be the same; and therefore the reason of the said AE to CF is given.

A a 3

Now

- Now let the reason of BE to DF be not the same, with that of AB to CD, and let it be as AB to CD, so BG to DF. Therefore the reason of the said BG to DF is given. But
- c 2. prop. the magnitude DF is given, therefore c BG is also given; and seeing that the whole BE
- d 4. prop. is given, d the rest GE shall be also given. But forasmuch as AB is to CD, as BG is to
- e 12. 5. DF, e so also is the whole AG to the whole CF; and therefore the reason of the said AG to CF is given: But the magnitude GE is
- f 11. def. given: Therefore f the magnitude AE is greater than the magnitude CF by a given magnitude, and in reason.

PROP. XV.

E G

A ——— 1 ——— 1 ——— B

F

C ——— 1 ——— — D

If two magnitudes AB and CD, have to one another a given reason, and that from each of them be taken away a given magnitude (to wit, from the magnitude AB the magnitude AE, and from the magnitude CD the magnitude CF) the remaining magnitudes EB and FD, either shall have to one another, a given reason, or the one of them shall be greater than the other by a given magnitude, and in reason.

Demonstr. For seeing that each magnitude AE and CF is given, the reason of AE to CF is given; and if it be the same with that of AB to CD, that of the remainder EB to the remainder FD, a shall be also the same; and therefore the reason of the said EB to FD shall be also given.

E G

A ——— 1 ——— 1 ——— B

F

C ——— 1 ——— — D

But if it be not the same, let it be as AB to CD, so AC to CF. Now the reason of AB to CD is given, there-

therefore also that of AG to CF shall be given.
 But CF is given, therefore *b* AG is given. *b* 2. *prop.*
 But AE is also given, therefore *c* the rest EG *c* 4. *prop.*
 is given; and seeing that as AB is to CD, so
 the part cut off AG is to the part cut off CF,
 and so also is *d* the rest GB to the rest FD; the *d* 4. *prop.*
 reason of the said GB to FD is also given.
 Therefore seeing that EG is given, EB is
 greater than FD *e* by a given magnitude, and *e* 11. *def.*
 in reason.

PROP. XVI.

G E
 A ——— l ——— l ——— F
 E
 C ——— ——— l ——— D

If two magnitudes AB and CD, have to one another a given reason and that from one of them, to wit, CD, there be taken away a given magnitude DE, and to the other AB there be added a given magnitude BF, the whole AF shall be greater than the rest CE, by a given magnitude, and in reason.

Demonstr. For seeing that the reason of AB to CD is given, let the same be made of BG to DE: Therefore *a* the reason of the said BG *a* 2. *def.* to DE is given. But DE is given, therefore *b* BG is also given. But BF is also given, *b* 2. *prop.* therefore *c* the whole GF is given. And see- *c* 3. *prop.* ing that as AB is to CD, so the part cut off BG, is to the part cut off DE; and *d* so also *d* 19. 5. is the remainder AG to the remainder CE; the reason of the said AG to CE is given: But GF is given, therefore the magnitude AF is greater than the magnitude CE by a given magnitude, and in reason.

P R O P. XVII.



If there be three magnitudes AB , E , and CD , and that the first AB be greater than the second E , by a given magnitude, and in reason. But the third CD be also greater than the same second E , by a given magnitude, and in reason; the first AB shall have to the third CD , either a given reason, or else the one shall be greater than the other by a given magnitude, and in reason.

Demonstr. For seeing that AB is greater than E by a given magnitude, and in reason, let the magnitude AF be taken away: Therefore the reason of the remainder FB to E is given. Again, seeing that CD is greater than the said E by a given magnitude, and in reason, let the given magnitude CG be cut off therefrom; and the reason of the remainder GD to E shall be given: Therefore *a* the reason of FB to GD shall be also given. But to the said FB and GD are added the given magnitudes AF and CG : Therefore the whole AB and CD shall either have to one another a given reason, or the one shall be greater than the other by a given magnitude, and in reason.

P R O P. XVIII.



If there be three magnitudes AB , CD , and EF , and that the one of them, to wit, CD , be greater than either of the other AB or EF , by a given magnitude, and in reason; either the two others AB and EF , shall have

have to one
be greater
and in rea

Demonstr.
is greater
magnitud
tude DG
reason of
Let the
fore the
But DG
And seei
 BH , b fo
 AH , the
be also g

Again
than EF
let the r
Therefo
 EF is g
 FK . T
 FK sha
therefor
as CI i
whole
the sai
reason
Therefo
shall b
said A
and F
 EF e a
ther,
by a g

have to one another a given reason, or the one shall be greater than the other by a given magnitude, and in reason.

Demonstr. Forasmuch as the magnitude CD is greater than the magnitude AB by a given magnitude, and in reason, let the given magnitude DG be taken there-from: Therefore the reason of the remainder CG to AB is given. Let the same be made of GD to BH, therefore the reason of the said DG to BH is given. But DG is given, therefore *a* BH is also given. *a* 2. *prop.* And seeing that as CG is to AB, so is GD to BH, *b* so also is the whole CD to the whole *b* 12. 5. AH, the reason of the said CD to AH shall be also given.

Again, seeing that the same CD is greater than EF by a given magnitude, and in reason; let the magnitude DI be cut off there-from: Therefore the reason of the remainder CI to EF is given: Let the same be made of DI to FK. Therefore the reason of the said DI to FK shall be also given. But DI is given, therefore FK is also given. And seeing that as CI is to EF, so is ID to FK; so also is the whole *c* CD to the whole EK; the reason of *c* 12. 5. the said CD to EK shall be given. But the reason of the same CD to AH is also given: Therefore *d* the reason of the said AH to EK *d* 8. *prop.* shall be given. And seeing that from the said AH and EK, the given magnitudes BH and FK are cut off, the magnitudes AB and EF *e* are either in a given reason to one another, or the one is greater than the other by a given magnitude, and in reason. *e* 15. *prop.*

PROP.


PROP. XIX.



If there be three magnitudes AB , CD , and E , and that the first AB , be greater than the second CD , by a given magnitude, and in reason; and that the second CD be greater than the third E , by a given magnitude, and in reason; also the first magnitude AB shall be greater than the third E , by a given magnitude, and in reason.

Demonstr. For seeing that CD is greater than E by a given magnitude, and in reason; let the given magnitude CF be taken there-from: Therefore the reason of the remainder FD to E is given. Again, seeing that AB is greater than the same CD by a given magnitude, and in reason: Let the magnitude AG be taken there-from: Therefore the reason of the remainder GB to CD is given: Let the same be made of GH to CF : Therefore the reason of the said GH to CF is given. But CF is given: Therefore also GH is given, and then AG is

a 3. prop.




b 19. 5.

also given, the whole AH shall be also given. But as GB is to CD , so is GH to CF , and so also b the remainder HB to the remainder FD : Therefore the reason of the said HB to FD is given. But the reason of the same FD to E is also given: Therefore the reason of HB to E is in like manner given, and so is also the magnitude AE : Wherefore the magnitude AB *c* is greater than E by a given magnitude, and in reason.

c 11. def.

OTHER

OTHERWISE.



Construction. Let there be three magnitudes AB, C, and D, and let AB be greater than C by a given magnitude, and in reason; but

let C be also greater than D, by a given magnitude, and in reason: I say that AB is greater than D by a given magnitude, and in reason.

Demonstr. Forasmuch as AB is greater than C by a given magnitude, and in reason, let the given magnitude AE be cut off there-from: Therefore the reason of the remainder EB to C is given. But the magnitude C is greater than the magnitude D by a given magnitude, and in reason; therefore *d* EB is greater than *d* 13. *prop.* D by a given magnitude, and in reason: Wherefore let the given magnitude EF be cut off there-from; and the reason of the remainder FB to D shall be given. But AF is *e* given. Therefore *f* AB is greater than D by a given *e* 3. *prop.* magnitude, and in reason. *f* 11. *def.*

PROP. XX.



If there be two given magnitudes, AB and CD, and that from them there be taken magnitudes AE

and CF, having to one another a given reason; either the remaining magnitudes EB and FD, shall have to one another given reasons; or else the one shall be greater than the other by a given magnitude, and in reason.

Demonstr.

- Demonstr.* For seeing that both the magnitudes AB and CD, are given, the reason of the said AB to CD is *a* also given; and if it be the same as of AE to CF, that of the remainder EB to the remainder FD shall be *b* also the same; and therefore the reason of the said EB to FD shall be also given. But if it be not the same, let it be so as that AE be to CF, as AG to CD. Now the reason of the said AE to CF is given: Therefore the reason of the said AG to CD is given. But CD is given, therefore *c* AG is also given. But the whole AB is likewise given, therefore *d* the remainder BG is given. And seeing that as AE is to CF, so is AG to CD, and also the remainder EG to the remainder FD, the reason of the said EG to FD is given. But GB is also given: Therefore the magnitude EB is greater *e* than the magnitude BD by a given magnitude, and in reason.
- a 1. prop. b 19. 5. c 3. prop. d 4. prop. e 11. def.

P R O P. XXI.

$$\begin{array}{ccccccc} & & G & & B & & \\ A & - & 1 & - & 1 & - & E \\ & & & & D & & \\ C & - & & - & 1 & - & F \end{array}$$

If there be two magnitudes given AB and CD, and to them be added other magnitudes BE and DF, having to one another a given reason: Either the whole AE and CF shall have to one another a given reason, or else the one shall be greater than the other by a given magnitude, and in reason.

- Demonstr.* For seeing that both the magnitudes AB and CD are given, their reason *a* is also given; and if it be the same reason as of BE to DF, the reason of the whole AE to the whole CF shall be also given; for it shall be *b* the same. But if it be not the same, let it be as BE is to DF, so BG to CD: Therefore the reason of the said BG to CD is given.
- a 1. prop. b 12. 5.
- But

But CD is given, therefore *c* also BG shall be *c* 2. prop.
 given. But the whole AB is given, therefore
 also the *d* remainder AG shall be given. And *d* 4. prop.
 seeing that as BE is to DF, so is BG to CD,
 and also *e* the whole GE to the whole CF, *e* 12. 5.
 the reason of the said GE to CF shall be like-
 wise given. But AG is given, therefore the
 magnitude AE is greater than the magnitude
 CF by a given magnitude, and in reason.

P R O P. XXII.

$$\begin{array}{c} \text{B} \\ \text{A} \text{---} \text{---} \text{---} \text{---} \text{C} \\ \text{D} \end{array}$$
 If two magnitudes AB
 and BC, have to some
 other magnitude D, a
 given reason, also their
 compound magnitude AC,
 shall have to the same magnitude D, a given
 reason.


Demonstr. For seeing that each magnitude AB
 and BC, hath a given reason to D, the reason
a of AB to BC is given; and by compounding, *a* 8. prop.
b the reason of AC to BC is given. But that *b* 6. prop.
 of BC to D is also given, therefore *c* the rea- *c* 8. prop.
 son of the said AC to D shall be likewise
 given.

P R O P. XXIII.

$$\begin{array}{c} \text{E} \\ \text{A} \text{---} \text{---} \text{---} \text{---} \text{B} \\ \text{F} \quad \text{G} \\ \text{C} \text{---} \text{---} \text{---} \text{---} \text{D} \end{array}$$
 If the whole AB be to
 the whole CD in a given
 reason, and that the parts
 AE and EB, be to the
 parts CF and FD in gi-
 ven reasons, altho' they be not the same, the whole
 (to wit, AB, AE, and BE,) shall be to the whole
 (to wit, CD, CF, and FD,) in given reasons.

Demonstr. For seeing that AE is to CF in a
 given reason, let the same be made of AB to
 CG; therefore the reason of the said AB
 to

- a 19. 5. to CG is given ; and consequently, also that a of the rest EB to the rest FG. But the reason of FD to the same EB is also given : Therefore
- b 8. prop. fore the reason of FD to FG b is likewise given ;
- c 5. prop. and therefore c that of FD to the remainder GD is also given. But the reason of AB to each of the magnitudes CD and CG is given :
- d 8. prop. Therefore d also the reason of CD to CG is given, and again e that of CD to the remainder GD. But the reason of FD to DG is
- e 5. prop. given, therefore also f that of the same CD to FD, and consequently
- f 8. prop. that of g CD to the remainder FC ; and therefore also the reason of CF to FD shall be given. But the reason of EB to FD is proposed to be given ; therefore the reason of CF to EB shall be given. Again, for that the reason of AB to CD is given ; and also that of CD to each of those FC and FD, the reason of the same AB to each of the said FC and b FD, shall be likewise given. But the reason of the said FD to EB is given : Therefore the reason of AB to BE shall be also given, and consequently
- g 5. prop. AB to the remainder i AE. Wherefore by
- h 8. prop. division k the reason of AE to EB shall be likewise given. But the reason of EB to FD is given. Therefore also that of AE to FD. In like manner, seeing that the reason of CD to AB is given ; and that of AB to each of his parts AE and EB ; also the reason of the said CD to each of the said AE and EB, l shall be given : Wherefore each of the magnitudes AB, CD, AE, EB, CF, and FD, is to each of the others in a given reason.
- i 5. prop.
- k sch. 5. pr.
- l 8. prop.



A


B

C

Demonstr
right line
C is given
therefore
D is given
the two
a a mean
angle ma
square c
and F is
rectangle
reasons
also given
and con
given (see
But D
E is given
But as
the rect
F, so
D and
the re
is to the
gle of
(seeing
and the
fore
B, so

PROP.

P R O P. XXIV.



*If of three right lines
A, B, and C, proportion-
al, A to B, as B to C,
the first A hath to the
third C a given reason,
it will also have to the
second B a given reason.*

Demonstr. For, let there be exposed another right line D, and seeing that the reason of A to C is given: Let the same be made of D to F; therefore the reason of D to F is given. But D is given, therefore F is also given; betwixt the two right lines D and F, let there be taken a mean proportional E. Therefore the rect-
angle made under D and F is equal ^a to the ^b square of E. But the same rectangle of D and F is ^c given: (for all the angles of that rectangle are given, being right angles, and the reasons that the sides have to one another are also given;) therefore the square of E is given, and consequently the same right line E is also given (for one equal thereto may be found, ^d seeing that the rectangle of D and F is given.) But D is given, therefore ^e the reason of D to E is given, and as A is to C, so D is to F. But as A is to C, ^f so the square of A is to the rectangle of A and C, and also as D is to F, so the square of D is to the rectangle of D and F. Therefore as the square of A is to the rectangle of A and C, so the square of D is to the rectangle of D and F. But the rectangle of A and C is equal to the square of B, (seeing that A, B, and C, are proportional) and that of D and F to the square of E, therefore as the square of A is to the square of B, so the square of D is to the square of E:

Where-

a 13. 6.

b 17. 6.

c 3. def.

d 14. 2.

e 1. prop.

f 1. 6.

g 12. 6.
h 2. def.

Wherefore g as A is to B , so D is to E . But the reason of D to E is given, therefore also the reason of A to B is given.

OTHERWISE.

Demonstr. Forasmuch as the reason of A to C is given, and that as A is to C , so the square of A and C , the reason of the said square of A to the rectangle made of A and C , is also given. But to that rectangle made of A and C

the square of B is equal (seeing that A, B , and C , are proportional;) therefore the reason of the square of A to the square of B is given; and by consequence, the reason of the line A to the line B is given; for to each of them A and B , we have exhibited an equal to the proper square of each one.

PROP. XXV.



If two lines AB and CD , given by position do intersect the point E in which they intersect one another, is given by position.

a 4. def.

Demonstr. For if it change its place, the one or the other of the lines AB and CD , would change its position: But so it is that by Supposition it changeth not: Therefore a the point E is given by position.

PROP. XXVI.

A ——— B

If the extremities A and B , of a right line AB , be given by position, that same right line AB is given by position and by magnitude.

Demonstr.


Demonstr. For if the point A remaining in its place, the position, or the magnitude of the right line AB shall change, the point B will fall elsewhere. But so it is, that by Supposition it doth not fall elsewhere. Therefore the right line AB is given by position, and by magnitude.

PROP. XXVII.

A — B *If one of the extremities A of a right line AB, given by position and magnitude be given, the other extremity B shall be also given.*

Demonstr. For if the point A remaining in its place, the point B shall change and fall in some other place, either the position of the right line AB, or its magnitude would change: But so it is that according to the Supposition, neither the one nor the other doth change. Therefore the point B is given.

OTHERWISE.



Constr. On the center A, with the distance AB, describe the circumference BC.

Demonstr. Therefore *a* 6. def. that circumference BC is given by position. But the right line AB is also given by position; therefore the point *b* B is *b* 25. prop. given.

g 12. 6.
h 2. def.

Wherefore g as A is to B , so D is to E . But the reason of D to E is given, therefore also the reason of A to B is given.

OTHERWISE.

Demonstr. Forasmuch as the reason of A to C is given, and that as A is to C , so the square of A and C , the reason of the said square of A to the rectangle made of A and C , is also given. But to that rectangle made of A and C the square of B is equal (seeing that

A —————
 B —————
 C —————

A , B , and C , are proportional) therefore the reason of the square of A to the square of B is given, and by consequence, the reason of the line A to the line B is given; for to each of them A and B , we have exhibited an equal to the proper square of each one.

PROP. XXV.



If two lines AB and CD , given by position do manifest the point E in which they intersect one another, is given by position.

a 4. def.

Demonstr. For if it change its place, the one or the other of the lines AB and CD , would change its position: But so it is that by Supposition it changeth not: Therefore a the point E is given by position.

PROP. XXVI.

A ————— B If the extremities A and B , of a right line AB , be given by position, that same right line AB is given by position and by magnitude.

Demonstr.

Demonstr.
 its place
 right line
 fall else
 tion it d
 right line
 magnitu

A —————
 and mag
 shall be
Demonstr.
 its place
 some
 tight l
 But so
 neither
 Theref

A —————

given
 given


Demonstr. For if the point A remaining in its place, the position, or the magnitude of the right line AB shall change, the point B will fall elsewhere. But so it is, that by Supposition it doth not fall elsewhere. Therefore the right line AB is given by position, and by magnitude.

PROP. XXVII.

A ——— B *If one of the extremities A of a right line AB, given by position and magnitude be given, the other extremity B shall be also given.*

Demonstr. For if the point A remaining in its place, the point B shall change and fall in some other place, either the position of the right line AB, or its magnitude would change: But so it is that according to the Supposition, neither the one nor the other doth change. Therefore the point B is given.


OTHERWISE.



Constr. On the center A, with the distance AB, describe the circumference BC.

Demonstr. Therefore *a* 6. def. that circumference BC is given by position. But the right line AB is also given by position; therefore the point *b* B is *b* 25. prop. given.

PROP. XXVIII.




If by a given point *A*, there be drawn a right line *DAE*, against another right line *BC*, given by position, the right line *DAE* so drawn, is given by position.

a 13. def.
b 30. 1.

Demonstr. For if it be not given, the point *A* remaining in its place, the position of the right line *DAE* may change: Let it then change if it be possible, and fall elsewhere, remaining parallel to *BC*, and let it be the line *FAG*: Therefore *BC* is parallel to the said line *FAG*. But a the same *BC* is also parallel to *DAE*: Therefore b *DAE* is parallel to the said line *FAG*, which is absurd; seeing they joyn together, and meet in *A*: Therefore the position of the right line *DAE* falls not elsewhere. Wherefore the said line *DAE* is given by position.

PROP. XXIX.



If to a right line *AB*, given by position, and to a point *C* given therein, there be drawn a right line *CD*, which shall make

a given angle *ACD*, the line drawn *CD*, is given by position.

Demonstr. For if it be not given by position, the point *C* remaining in its place, the position of the line *CD* observing the magnitude of the angle *ACD*, will fall elsewhere. Let it fall elsewhere then if it be possible, and let it

it be CE. Therefore the angle ACD is equal to the angle ACE, the greater to the lesser, which is absurd. Therefore the position of the right line CD, shall not fall elsewhere; and therefore the said line CD is given by position.

PROP. XXX.



If from a given point *A*, be drawn to a right line *BC*, given by position, a right line *AD*, making a given angle *ADB*, that line drawn *AD* is given by position.

Demonstr. For if it be not given, the point *A* remaining in its place, the position of the right line *AD* keeping, magnitude of the angle *ADB*, will change. Let it change then, and let it be the right line *AE*: Therefore the angle *ADB* is equal to the angle *AEB*, the greater to the lesser, which is absurd. Therefore the position of the right line *AD* doth not change; and therefore the said line *AD* is given by position.

OTHERWISE.


Constr. By the point *A* let there be drawn the line *EAF*, parallel to the right line *BC*.

Demonstr. Then seeing that by the given point *A*, and against the right line *BC*, given by position, there is drawn the right line *EF*, those lines *EF* and *BC* are parallels. But on

B b z

the

b 29. 1.



the same lines doth also fall the right line AD. Therefore the angle FAD is equal to the given angle ADB; and

therefore it is also given. Wherefore to the right line EF given by position, and to the given point A therein, there is drawn the right line AD, making the given angle FAD.


c 29. prop. Therefore the said line AD is given by position.

OTHERWISE.

Constr. In the line BCE, let there be taken the given point C, and by the same let there be drawn the line CF, parallel to the said DA.

d 29. 1. *Demonstr.* Forasmuch as AD and FC are parallels, and that on them there doth fall the right line BCE, the angle FCB is equal to the given angle ADB; and therefore it is also given. And seeing that the right line BC

e 29. prop.




is given by position, and that to a given point C therein, there is drawn the right line FC, making the given angle FCB, that same line FC is given by position. But by the given point A, opposite to the line FC given by position, there is drawn the line

f 28. prop. AD. Therefore the said line AD is given by position.

OTHER-

OTHERWISE.



Constr. In the right line BC assume some point at F, and draw AF.


Demonstr. Forasmuch as each point A and F is given, the right line AF is given by position. But the line BC is also given by position. Therefore * the angle AFD is given. But by supposition, the angle ADF is given: Therefore DAF (which is the residue of two right angles) is given; and seeing that to the right line AF given by position, and to the given point therein A there is drawn the right line DA, making the given angle DAF, that same line DA is given by position. 26. prop. h 32. 1. i 25. prop.

Scholium.

* EUCLIDE supposeth here that two right lines being given by position, and inclining to one another do make a given angle, which some do demonstrate after this manner.

Demonstr. Forasmuch as the two right lines given by position, do incline to one another, the inclination of those lines is given. But the angle is the inclination of the lines: Therefore the angle which makes the right lines given by position, and inclining to one another, is given.

Another thus demonstrateth it.



Constr. Let there be two right lines inclining to one another, as AB and CB, given by position, and in the line AB let there be taken a given point A, and in BC also some point, as C; and let the right line AC be drawn.


Demonstr. Seeing that as well the point B, as each of the points A and C is

given, *k* the three right lines AB, BC, and AC, are given by magnitude. Wherefore of three direct lines equal unto them, a triangle may be constituted: Let there then be made the triangle FDE, having the side FD equal to the side AB, the side FE equal to the side AC, and the base DE equal to the base BC.

Seeing then the angles comprised of equal right lines are equal, we have found the angle FDE equal to the angle ABC; and therefore the same $\angle$ angle ABC is given.

l i. def.

P R O P. XXXI.




If from a given point A there be drawn to a right line given by position BC, a right line AD, given by magnitude, that line AD shall be also given by position.

Constr. From the center A, with the distance AD, let the circle DEF be described.

Demonstr.

Demonstr. Forasmuch as the center A is given by position, and the semidiameter AD by magnitude, the circle DEF is given by position. *a* 6. def. But the right line BC is also given by position: Therefore the point of intersection D *b* 25. prop. is given, and seeing that the point A is also given: *c* the right line AD is given by position. *c* 26. prop.

PROP. XXXII.



If unto parallel right lines AB and CD, given by position, there be drawn a right line EF, making the given angles BEF and EFD, the line drawn EF shall be given by magnitude.

Constr. For let there be taken in the line CD a given point G, and from that point let be drawn GH parallel to FE.

Demonstr. Forasmuch as the lines EF and HG are parallels, and that on them doth fall the line CD; *a* the angle EFD is equal to the *a* 29. 1. angle FGH. But the angle EFD is given, therefore the angle FGH is also given. And forasmuch as to the right line CD given by position, and to the point G given in the same, there is drawn the right line GH, making the given angle FGH, *b* the said line GH is given *b* 29. prop. by position. But AB is also given by position, therefore *c* the point H is given. But the *c* 25. prop. point G is also given: Therefore *d* the line *d* 26. prop. GH is given by magnitude, and is *e* equal to *e* 34. 1. EF. Wherefore *f* the said line EF is given *f* 1. def. by magnitude.

PROP. XXXIII.



If unto parallel right lines AB and CD, given by position, there be drawn a right line EF, given by magnitude, that line EF shall make the given angles BEF and DFE.

Constr. For let there be taken in the right line AB the point G, and by that point let there be drawn the line GH parallel to EF.

- Demonstr.* Therefore EF is equal to the said a GH. But EF is given by magnitude, therefore GH is also given by magnitude. But the point G is given, and therefore if on that point, with the distance GH, there be described a circle, b that circle shall be given by position: Let it be then described, and let it be HKL, the said circle HKL is therefore given by position. But the line CD which doth cut the circumference KHL in H, is also given by position. Therefore the said point of intersection H c is given. But the point G is given: Therefore d the right line GH is given by position. But the right line CD is also given by position: Therefore e the angle GHF is given. But to that angle f the angle EFD is equal: Therefore the angle EFD is given; and therefore also the angle BEF; for that it is the residue of the sum of two g right angles.

OTHERWISE.

Constr. Let there be taken in the right line CD the point G, and let GD be put equal to EF,

EF, then from the center G, with the distance GD, let there be described the circle DBH, and draw GB.

Demonstr. Forasmuch as the center G is given by position, and the semidiameter GD by magnitude, the circle BDH *h* is given by position. *h* 6. def. But the line AB is also given by position: Therefore *i* the point B is given. But the point *i* 25. prop.




G is also given, therefore *k* the right line GB *k* 26. prop. is given by position. But the right line CD is also given by position: Therefore *l* the angle BGD is *l* sch. 30. given. Wherefore if EF *prop.* be parallel to BG, the angle EFD *m* shall be *m* 29. 1. given, and consequently also the other angle

BEF. But the right lines BG and EF being not parallels, let them meet in the point H. Forasmuch as EB is parallel to FG, and EF is equal to GD, that is to say, to BG; also FH *n* shall be equal to GH (for EH and BH *n* 14. 5. being cut proportionally *o* by the parallel FG, *o* 2. 6. as EF is to FH, so is BG to GH; and by permutation, as EF is to BG, so is FH to GH :) Therefore *p* the angle HFG is equal to *p* 5. 1. the angle HGF, but the said angle HGF is given (for that it is equal *q* to the given angle *q* 15. 1. BGD :) Therefore the angle HFG is also given. But to that angle the angle BEF is equal; and therefore is given, as also the remaining angle EFG.

PROP.


PROP. XXXIV.



Constr For from the point E let there be drawn the line EH, perpendicular to the line CD.

Demonstr. Forasmuch as from the given point E there is drawn to the line CD the right line EH, making the given angle EHG *a* the said line EH is given by position, but both the one and the other lines AB and CD is also given by position. Therefore *b* the points of intersection K and H, are given. But the point E is also given: Therefore *c* each line EK and KH is given. Wherefore *d* the reason of the said EK to KH is given. But as EK is to KH, so is EF to FG; (for in the triangle GEH the line KF being parallel to HG, the sides EH and EG are cut proportionally :) Therefore the reason of the said EF to FG is given.

OTHERWISE.



say that the reason of GE to EF is given.

Demonstr.

If from a given point E, there be drawn unto parallel right lines AB and CD, given by position, a right line EFG, that right line EFG shall be divided in a given reason (to wit) as EF to FG.

Demonstr.
drawn
duced to
the p
position
the gi
given
is also
point
the po
the li
and th
EK is
to EF
being
the tr
and th
and b
EF.)
EG to

If
given
which
wit)
E, t
right
be g
C

draw
BC
I
poi
the
AH
Bu
the

Demonstr. For from the point E let there be drawn to CD the perpendicular EH, and produced to the point K; seeing therefore that from the point E to the right line CD, given by position, there is drawn the line EH, making the given angle EHG, *a* the said line EH is given by position. But each line AB and CD *a* 30. *prop.* is also given by position: Therefore *b* each point of intersection H and K is given. But *b* 25. *prop.* the point E is also given, therefore *c* each of the lines EH and EK is given by magnitude; *c* 26. *prop.* and therefore *d* the reason of the said EH to EK is given. But *e* as EH is to EK, so is EG *d* 1. *prop.* to EF (for the opposite angles at the point E *e* 4. 6. being equal, and the lines AB and CD parallels, the triangles EHG and EKF are equiangled; and therefore as EH is to EG, so is EK to EF; and by permutation as EH to EK, so is EG to EF.) Therefore the reason of the said lines EG to EF is given.

P R O P. XXXV.


If from a given point A, to a right line BC, given by position, there be drawn a right line AD, which let be divided in E, in a given reason (to wit) as AE to ED, and that by the point of section E, there be drawn a right line FEG, opposite to the right BC, given by position, the line FG drawn shall be given by position.

Constr. For from the point A, let there be drawn the line AH, perpendicular to the line BC.

Demonstr. For seeing that from the given point A there is drawn to BC given by position, the right line AH making the given angle AHD, *a* the said line AH is given by position, *a* 30. *prop.* But BC is also given by position: Therefore *b* 25. *prop.* the point H is given. But the point A is also given:

e 26. prop. given : Therefore c the line AH is given by magnitude and by position.

d 2. 6.



e 6. prop.


f 2. prop.

g 27. prop.

h 28. prop.

given by magnitude. But AK is also given by position, and the point A is given: Therefore g the point K is also given, and seeing that by the said given point K there is drawn the line FG, opposite to the right line BC given by position; the said line FG h is given by position.

PROP. XXXVI.




If from a given point A, there be drawn to a right line BC given by position, a right line AD, and to it be added a right line AE, having to the same AD a given reason, and that by the extremity E of the added line AE, there be drawn a right line FEK, opposite to the line BC, given by position, that same line FEK shall be given by position.

Constr. For from the point A let there be drawn to the line BC, the perpendicular AL, and let it be prolonged to the point G.

Demonstr. Forasmuch as from the given point A, there is drawn to the right line BC, given

given by position, the right GL, which makes
 the given angle GLD, *a* that line GL is a 30. prop.
 given by position. But BC is also given b 26. prop.
 by position, therefore *b* the point L is gi-
 ven; and seeing that the point A is also c 26. prop.
 given, the line *c* AL is given. But forasmuch
 as the reason of AE to AD is given, and that d 4. 6.
 as the said AE is to AD, so is AG to AL;
 (because the triangles ALD and AGE are
 equiangled) the reason of AG to AL is also
 given. But AL is given by magnitude: There- e 2. prop.
 fore *e* AG is given by magnitude. But it is
 also given by position, and the point A is f 27. prop.
 given: Therefore *f* the point G is also given.
 And seeing that by the same given point G,
 there is drawn the line FK, opposite to the g 28. prop.
 right line BC, given by position, *g* the said
 line FK is given by position.

PROP. XXXVII.



If unto parallel right
 lines AB and CD, gi-
 ven by position, there be
 drawn a right line EF,
 divided in the point G,
 in a given reason, (to
 wit, of EG to GF;) *but*
 if by the point of
 section G, there be
 drawn opposite to the

right lines AB or CD, given by position, a right
 line HGK, that line drawn shall be given by po-
 sition.

Constr. For let there be taken in the line AB
 the given point L, and from that point let
 there be drawn the line LN, perpendicular
 to CD.

Demonstr. Seeing that from the given point
 L, there is drawn to the right line CD, the
 line LN, making the given angle LND, the
 said

- a 30. *prop.* said LN *a* is given by position. But CD is also given by position: Therefore the point
 b 25. *prop.* N *b* is given. But the point L is also given:
 c 26. *prop.* Therefore *c* the line LN is given; and seeing that the reason of FG to GE is given, and that * as FG is to GE, so is NM to ML, the reason of the said MN to ML is given;
 d 6. *prop.* and in compounding, *d* the reason of LN to LM is also given. But LN is given by magnitude, therefore ML is *e* given by magnitude.
 e 2. *prop.* But it is also given by position, and the point
 f 27. *prop.* L is given: Therefore the point M *f* is also given. And considering that by the said point M there is drawn the right line KH, opposite to the right line CD, given by position, the said line KH is also given by position.

Scholium.

* EUCLIDE supposeth here, that as FG is to GE, so NM is to ML; but by another it is thus demonstrated.

The lines EF and LN are parallels or not parallels: Let them in the first place be parallels, and forasmuch as by Construction the lines EL, FN, EF, and LN, are parallels, EN shall be a parallelogram; and therefore the side EF is equal to the side LN. Again, seeing that MG is parallel to FN, and GF to MN, GN shall be also a parallelogram; and therefore the side GF is equal to the side MN. Wherefore the equal sides EF and LN, shall have to the equal sides FG and MN, *g* one and the same reason. Therefore as EF is to FG, so is LN to MN; and in dividing, *h* as GE to GF, so is LM to MN.

g 7. 5.

h 17. 5.

Now
parallels

A

H

CF

drawn
OG and
k as O
poundr
it bath
is NM
as FG

K


B

D

add
again
CD,
sition
C
AB,

Now

Now suppose that the lines EF and LN be not parallels, but that they meet in the point O. For-



asmuch as in the triangle OFN there is drawn HK, parallel to FN one of the sides; i the sides i 2. 6.

OF and ON are divided proportionably; and therefore as FG is to GO, so is NM to MO. Again, seeing that in the triangle OGM there is

drawn EL, parallel to the side GM, the sides OG and OM are divided proportionally: Wherefore k as OE is to EG, so is OL to LM; and by com- k 2. 6. pounding, l as OG is to EG, so is OM to LM; but it hath been demonstrated that as FG is to GO, so l 18. 5. is NM to MO; therefore in reason of equality, m m 22. 5. as FG is to GE, so is NM to ML.

PROP. XXXVIII.



If unto parallel right lines AB and CD, there be drawn a right line EF, and that to it there be added some other right line EG, which hath a given reason to the same EF, but if by the extremity G of the

added line EG, there be drawn a right line HK, against the parallels given by position AB and CD, the line drawn HK shall be also given by position.

Constr. For let there be taken in the line AB, the given point N, and from thence let there

Again, Let EF be put equal to BC ; and seeing that BC is given by magnitude, EF shall be so also. But the said EF is in like manner given by position, and the point E is given: Therefore the point F is given.

b 27. prop.

Furthermore, Let FG be taken equal to AC . Now forasmuch as the said AC is given by magnitude, FG is so also. But FG is also given by position, and the point F is given: Therefore the point G is also given. Now from the center E , with the distance ED , let there be described the circle DHK , c c 6. def. and that circle shall be given by position. Again, on the center F , and distance FG , let there be described the circle GLK . Therefore the said circle GLK is given by position; and therefore the point of intersection I is given. But each of the points E and F is given: Therefore each line EK , EF , and FK , is given by position, and magnitude. Therefore the triangle EKF is given * by kind; but it is equal and alike to the triangle ABC ; and therefore the triangle ABC is also given by kind.

Scholium.

EUCLIDE supposeth here that a triangle whose sides are given by magnitude and position, is given by kind; but the ancient Interpreters demonstrate it in a manner thus. Forasmuch as the right lines KE and EF are given, g g 1. prop. the reason which they have to one another is given. Also the right lines EF and FK being given, their reason is also given; and in like manner, the reason

C c


of



there be drawn to CD the perpendicular NM, and let it be prolonged to the point L.

- Demonstr.* Forasmuch as from the given point N there is drawn to the right line CD, given by position, the right line NM making a given angle NMF, the said angle NMF *a* is given by position. But the line CD is also given by position: Therefore *b* the point M is given. But the point N is also given: Therefore *c* the line NM is given, and for that the reason of EG to EF is given, and that *d* as EG is to EF, so is LN to NM, the reason of LN to NM is also given: But NM is given, therefore LN is *e* also given. But the point N is given: Therefore *f* the point L is also given. Seeing then that by the given point L there is drawn the right line HK, opposite to the line AB given by position, *g* the said line AK is also given by position.
- a* 30. *prop.*
b 25. *prop.*
c 26. *prop.*
d sch. 37. *prop.*
e 2. *prop.*
f 27. *prop.*
g 28. *prop.*

PROP. XXXIX.



If all the sides of a triangle ABC are given by magnitude, the triangle is given by Kind.

Constr. For, let there be exposed the right line DG given by position, ending in the point D; but being infinite towards the other part G, and therein let there be taken DE, equal to AB.

Demonstr. Now seeing the said AB is given by magnitude, DE is so also; but the same DE is also given by position, and the point

- a* 27. *prop.* D is given: Therefore *a* the point E is given.

Again;

Again, Let EF be put equal to BC; and seeing that BC is given by magnitude, EF shall be so also. But the said EF is in like manner given by position, and the point E is given: Therefore *b* the point F is given.

b 27. prop.

Furthermore, Let FG be taken equal to AC. Now forasmuch as the said AC is given by magnitude, FG is so also. But FG is also given by position, and the point F is given: Therefore the point G is also given.

Now from the center E, with the distance ED, let there be described the circle DHK, *c* *c* 6. def. and that circle shall be given by position.


Again, on the center F, and distance FG, let there be described the circle GLK. Therefore *d* the said circle GLK is given by position; and therefore *e* the point of intersection *e* 25. prop.

K is given. But each of the points E and F is given: Therefore each line *f* EK, EF, and FK, is given by position and magnitude. *f* 26. prop.

Therefore the triangle EKF is given * by kind; but it is equal and alike to the triangle ABC; and therefore the triangle ABC is also given by kind.


Scholium.

* EUCLIDE supposeth here that a triangle whose sides are given by magnitude and position, is given by kind; but the antient Interpreters demonstrate it in a manner thus. Forasmuch as the right lines KE and EF are given, *g* *g* 1. prop. the reason which they have to one another is given. Also the right lines EF and FK being given, their reason is also given; and in like manner, the reason of



of the said EK and FK is given. Again, seeing that the same lines KE and EF are given by position, *h* the angle KEF is given by magnitude: *prop.* Moreover, the right lines EF and FK being given by position, the angle EFK is given by magnitude, as is also the residue EKF , and so in the triangle EKF are all the angles given, and also the reasons of the sides: Therefore *i* the said triangle EKF is given by kind.

PROP. XL.



If the angles of a triangle ABC , are given by magnitude, the triangle is given by kind.

Constr. Let there be exposed the right line DE , given by position and by magnitude; and let there be constituted at the point D the angle FDE , equal to the angle CBA ; but in the point E the angle DEF , equal to the angle BCA ; therefore the third angle BAC is equal to the third angle DFE .

Demonstr. For each of the angles constituted in the points A , B , and C , is given: Therefore each of those which are posited in the points D , F , and E , is also given; and seeing that to the right line DE given by position, and to the point D given therein, there is drawn the right line DF , which makes the given angle EDF , *a* the line DF is given by position; and by the same reason, the line *b* EF is given by position: Therefore *b* the point F is given by position. But each of the *c* $26. prop.$ points D and E is given: Therefore *c* each of the

the line
tude.
by kind
Therefo

If a
and the
constitu
the tria
Const
line D
thereo
be con
angle
Dem
Theref
seeing
sition,
drawn
gle D

CA

E

F is
 DF ,
mag


the lines DF, DE, and EF, is given by magnitude. Wherefore the triangle DFE is given by kind; and is alike to the triangle ABC: Therefore the triangle ABC is given by kind.

P R O P. XLI.

If a triangle ABC, hath one angle BAC given, and that the two sides BA and AC, which do constitute it, have to one another a given reason, the triangle is given by kind.

Constr. For, let there be exposed the right line DF given by magnitude and position. But thereon, and at the given point F, let there be constituted the angle DFE equal to the angle BAC.

Demonstr. Now the angle BAC is given: Therefore also the angle DFE is given, and seeing that to the right line DF given by position, and from the given point F therein, is drawn a right line FE, making the given angle DFE, ^a the said line FE is given by position. But seeing that the ^{a 29. prop.} reason of AB to AC is given, let the same be made of DF to FE, then let DE be drawn. Therefore the reason of DF to FE is given. But DF is given: Therefore ^b FE is given by ^{b 2. prop.} magnitude. But the same FE is also given by position, and the point F is given. Therefore ^c the ^{c 27. prop.} point E is also given. But each of the points D and




F is given: Therefore ^d each of the right lines ^{d 26. prop.} DF, FE, and DE, is given by position and magnitude. Wherefore ^e the triangle DEF is ^{e 39. prop.} given

Cic 2

f 6. 6.

given by kind. And seeing that the two triangles ABC and DEF have an angle equal to an angle, that is to say, the angle BAC to the angle DFE , and the sides which constitute those equal angles, proportional; *f* the triangle ABC is alike to the triangle DEF . But the triangle DEF is given by kind: Therefore the triangle ABC is given by kind.

P R O P. XLII.



a 2. prop.

b 2. prop.

c 39. prop.

If the sides of a triangle ABC , be to one another in given reasons, the triangle ABC is given by kind.

Constr. For, Let there be exposed the right line D , given by magnitude, and seeing that the reason of BC to AC is given, let the same be made of D to E .

Demonstr. Now D is given, therefore a E is also given. Again, seeing

that the reason of AC to AB is given, let the same be made of E to F . Now E is given, therefore *b* F is also given. Now of three right lines, equal to the three given right lines D , E , and F , (and of which three lines, two of them, in what manner soever they be taken, are greater than the other :) Let there be constituted the triangle GKH , in such sort as D may be equal to HK ; but E is equal to KG , and GH equal to F ; therefore each of the said lines HK , KG , and GH , is given by magnitude: Wherefore *c* the triangle HKG is given by kind. And seeing that as BC is to CA , so is D to E , and that D is equal to HK , and E to KG , as BC is to CA ,

so HK
is to KG ,
that as
fore by
so is H
 ABC is



C
cle I
De
divid
said
midi
the
same
son
there
er th
than
EG
dift
Now
that

so HK is to KG. Again, seeing that as CA is to AB, so is E to F, and that E is equal to KG, and F to GH; as CA is to AB, so is KG to GH. But it hath been demonstrated that as BC is to CA, so is HK to KG: Therefore by reason of equality, as BC is to AB, so is HK to GH. Therefore *d* the triangle ABC is also given by kind. *d* 5. 6.

P R O P. XLIII.


If the sides BC and BA, about one of the acute angles of a rectangled triangle ABC, have to one another a given reason, that triangle is given by kind.

Constr. Let there be exposed the right line DE given by magnitude and position, and on it let there be described the semicircle DGE: Therefore *a* the semicircle DGE is given by position. *a* 6. def.

Demonstr. For the line DE being given, and divided in two equal parts, the center of the said circle is given by position, and the semidiameter by magnitude. And forasmuch as the reason of BC to BA is given, let the same be made of DE to F: Therefore the reason of DE to F is given. But DE is given, therefore F *b* is also given. Now BC is greater than *c* AB: Therefore ED is *d* also greater than F. Let DG be fitted equal to F, and let *d* EG be drawn; then on the center D, with the distance DG, let the circle GK be described. Now that circle *e* is given by position, seeing *e* that the center D is given, and the semidiameter


C c 3

meter



- meter DG also given by magnitude. But the semicircle DGE is also given by position:
 f 25. prop. Therefore f the point of intersection G is given.
 But the points D and E are also given, there-
 g 26. prop. fore g each of the right lines DE, DG, and EG, is given by position and magnitude.
 h 39. prop. Wherefore h the triangle DGE is given by kind. And seeing that the triangles ABC and DGE have an angle equal to an angle, to wit, the right angle BAC to the right angle DGE, and the sides about the angles CBA and EDG proportional. But each of the others ACB and DEG are less than a right angle: Those triangles ABC and DEG k are alike. But the triangle DGE is given by kind: Therefore the triangle ABC is also given by kind,

PROP. XLIV.



If a triangle ABC, hath one angle B given, and that the sides BA and AC, about another angle BAC, have to one another a given reason, the triangle ABC is given by kind.


Constr. Now the given angle B is either acute or obtuse, (for it was a right angle in the foregoing Proposition.) Let it be in the first place acute, and from the point A let AD be drawn perpendicular to BC.

- Demonstr.* Therefore the angle ADB is given: But the angle B is also given; and therefore the third angle BAD is given: Wherefore
 a 40. prop. a the triangle ABD is given by kind; and
 b 3. def. therefore b the reason of BA to AD is given. But the reason of the same BA to AC is also
 c 8. prop. given; Therefore c the reason of AD to AC is given, and the angle ADC is a right angle: Where-


Wherefore the triangle *d* ACD is given by *d* 43. prop. kind: Therefore *e* the angle C is given. But *e* 3. def. the angle B is also given; and therefore the other angle BAC is given: Therefore *f* the triangle ABC is given by kind.

Constr. Now let the angle ABC be obtuse, and on the side CB prolonged, let there be drawn the perpendicular AD.

Demonstr. Forasmuch as the angle ABC is given, the angle ABD which follows it, shall be given. But the angle ADB is also given: Therefore the third angle DAB is given. Wherefore *g* the angle ABD is given by kind; *g* 40. prop. and therefore *h* the reason of DA to AB is given. But the reason of AB to AC is also given: Therefore *i* the reason of DA to AC is given, and the angle D is a right angle: Therefore the triangle DAC is given by kind, and therefore the angle ACB is given. But the angle ABC is also given: Therefore the third angle BAC is given. Wherefore the triangle ABC is given by kind,



PROP. XLV.



If a triangle ABC , hath one angle BAC given, and that the line compounded of the two sides AB and AC , about the said given angle BAC , hath to the other side BC a given reason, the triangle ABC is given by kind.

Constr. For, let the angle BAC be divided into two equal parts by the line AD , therefore *a* the angle CAD is given.


Demonstr. Seeing that as AB is to AC , so *b* is BD to CD ; by compounding, *c* as the line compounded of CAB is to CA , so is BC to CD ; and by permutation, as the line compounded of CAB is to CB , so is CA to CD . But the reason of the line compounded of CAB to BC is given; therefore the reason of CA to CD is also given, and the angle CAD is given. Therefore *d* the triangle ACD is given by kind, and therefore the angle C is given. But the angle BAC is also given: Therefore *e* 40. *prop.* the third angle B is given: Wherefore *e* the triangle ABC is given by kind.

OTHERWISE.


Constr. Let BA be prolonged directly unto the point D , in such sort as that AD may be equal to AC , and let CD be joined.

Demonstr.

Demonstr. Forasmuch as the reason of the line compounded of CAB to CB is given, and that AD is equal to AC, the reason of the whole line BD to BC is given. But the angle ADC is also given, for it is the half of the given angle BAC (for that the said angle BAC *f* is *f* 32. 1. equal to the two internal angles ACD and ADC, which are *g* e-g 5. 1. equal to one another, be-^h 44. prop. ing the sides AC and AD are equal :) Wherefore the triangle BDC *h* is given by kind, and therefore the angle B is given. But the angle BAC is also given: Therefore the remaining angle ACB is given: Wherefore i the triangle i 40. prop. ABC is given by kind.



PROP. XLVI.



If a triangle ABC hath one angle B given, and that the line CAB compounded of the two sides, AC and AB, about another angle BAC, hath to the other side BC a given reason, the triangle ABC is given by kind.

Constr. For let the angle BAC be divided into two equal parts, by the line AD.


Demonstr. Therefore (as hath been shewn in the foregoing Proposition) the compound line CAB is to CB, as AB is to BD. But the reason of the said compound line CAB to CB is given: Therefore also the reason of AB to BD is

a 41. prop.
b 2. def.

is given. But the angle B is also given: Therefore the triangle ABD *a* is given by kind; and therefore *b* the angle BAD is given. But the angle BAC is double to that of BAD; and therefore it is also given. Therefore the third angle C is given. Wherefore the triangle ABC is given by kind.

OTHERWISE.

Constr. Let BA be prolonged directly, and let AD be put equal to AC, and let CD be joined.




c 41. prop.
d 3. def.

D is given: Therefore the angle BAC which is double to BDC, is also given: Wherefore the other angle ACB is given; and therefore the triangle ABC is given by kind.

Demonstr. Forasmuch as the reason of the line compounded of CAB to CB is given, and that AD is equal to AC, the reason of BD to BC is given; and the angle B is also given: Therefore the triangle CDB *c* is given by kind; and therefore *d* the angle

PROP.

PROP. XLVII.




Rectiline figures as *AB-CDE*, given by kind, are divided into triangles given by kind.

Constr. For let the right lines *EB* and *EC* be drawn.

Demonstr. Forasmuch as the rectiline figure *ABCDE* is given by kind, the angle *a* *BAE* *a* 3. def.

is given, and the reason of the side *AB* to *AE* is also given: Therefore *b* the triangle *BAE* *b* 41. prop. is given by kind. Wherefore the angle *ABE* is given. But the whole angle *ABC* is also given: Therefore *c* the remaining angle *EBC* *c* 4. prop. is given. But the reason of the side *AB* to the side *BE*, and also that of *AB* to *BC* is given: Therefore *d* the reason of *BC* to *BE* is *d* 8. prop. given, and the angle *CBE* is also given: Therefore *e* the triangle *BCE* is given by *e* 41. prop. kind. By the same discourse it may be demonstrated that the triangle *CDE* is given by kind. Therefore the rectiline figures given by kind divide themselves into triangles given by kind.

PROP. XLVIII.




If on one and the same right line *AB*, are described triangles as *ACB* and *ABD*, given by position, those triangles shall have to one another a given reason, as *ACB* to *ABD*,

Constr.

Constr. For from the points A and B, let there be drawn at right angles on the line AB, the lines AE and BG, and prolonged unto the points F and H; but by the points C and D, let there be drawn the lines ECG and FDH, parallel to AB.

- Demonstr.* Forasmuch then as the triangle ABC is given by kind, *a* the reason of CA to BA is given, and the angle CAB also given; but the angle BAE is given: Therefore the remaining angle CAE is also given; but the angle CAE is given; and therefore the other angle ACE is also given. Wherefore *b* the triangle AEC is given by kind. Now the reason of EA to AB *c* is given; (for *d* the reason of EA to AC, and that of AC to AB is given;) and in like manner, the reason of FA to AB is given. Therefore *e* the reason of EA to AF is given; but as AE is to AF, so *f* the parallelogram AH to the parallelogram AG; but ACB is *g* the half of AH, and ADB the half of AG; therefore the reason of the triangle ACB to the triangle ADB is given; for it is the same reason with that of AH to AG *h*; that is to say, of EA to AF, which is given.
- a 3. def.
b 40. prop.
c 8. prop.
d 3. def.
e 8. prop.
f 1. 6.
g 41. 1.
h 15. 5.

PROP. XLIX.



If on one and the same right line AB there be described any two rectilineal figures AECFB and ADB, given by kind, they shall have to one another a given reason (to wit) AECFB to ADB.

Constr. For let the lines FA and FE be drawn:

drawn
AFE,
Dem
right
ECF
ECF t
pound
given
FAB
given
right
CF t
pound
given
ABD
AEC

CI
TI
BU
fo
W
B
T
is


EUCLIDE: DATA.

411

drawn: Therefore each of the triangles a ABF, a 47. prop. AFE, and ECF is given by kind.

Demonstr. Seeing that on one and the same right line EF there are described the triangles ECF and EAF, given by kind; the reason of ECF to EAF b is given. Therefore by com- b 48. prop. pounding, c the reason of AECF to EAF is c 6. prop. given. But the reason of the said EAF to FAB is given, d being they are triangles d 48. prop. given by kind, described on one and the same right line AF: Therefore e the reason of AE- e 8. prop. CF to FAB is given. Wherefore by com- pounding, f the reason of AECFB to FAB is f 6. prop. given. But the reason of the same FAB to ABD g is given: Therefore h the reason of g 48. prop. AECFB to ABD is also given. h 8. prop.

PROP. L.




If two right lines AB and CD, have to one another a given reason, and that on those lines there be described rectiline figures AEB and CFD, alike, and alike posited, they will have to one another a given reason.

Demonstr. To the two lines AB and CD, let there be taken a third proportional: Therefore as AB is to CD, so is CD to G. But the reason of AB to CD is given: Therefore the reason of CD to G is also given: Wherefore a the reason of AB to G is given. a 8. prop. But b as AB is to G, so is AEB to CFD: b cor. 19. Therefore the reason of the same AEB to CFD $20.$ 6. is given.

PROP.

PROP. LI.



If two
right lines
 AB and
 CD , have
to one ano-
ther a gi-
ven reason,
and that
upon them
there be
described


any rectiline figures AEB and CFD , given by kind, they will have to one another a given reason, (to wit, that of AEB to CFD .)

Constr. For on AB let the rectangled figure AH be described alike and alike posited to DF .

Demonstr. Now DF is given by kind: Therefore also AH is given by kind. But AEB is also given by kind, and described on the same line AB : Therefore *a* the reason of AEB to AH is given: And seeing that the reason of AB to CD is given, and that on those lines are described the rectiline figures AH and DF alike, and alike posited, the reason *b* of the said line AH to DF is given. But the reason of AEB to AH is also given: Therefore the *c* 8. *prop.* reason *c* of AEB to DF is given.

PROP.

PROP. LII.



If on a right line *AB*, given by magnitude, there be described a figure *ACB*, given by kind, that figure *ACB* is given by magnitude.

Constr. For on the same line *AB*, let the square *AD* be described. Therefore *AD* is given by kind * and by magnitude.


Demonstr. Seeing that on the right line *AB*, are described the two rectiline figures *ACB* and *AD*, given by kind, a the reason of *ACB* to *AD* is given: Therefore b *ACB* is given by magnitude. a 29. prop.
b 2. prop.

Scholium.

* The antient Interpreter hath noted here that every square is given by kind; for that all the angles thereof are given; being all equal and right angles: But also the reasons of the sides are given; for those sides being all equal, their reasons are also equal. Moreover, whensoever a square is exposed, a square equal thereto may be exhibited; and therefore the square is given by magnitude, as also each side thereof.

PROP.

PROP. LIH.




If there be two figures AD and EH , given by kind, and that one side BD , of the one, hath to a side FH of the other, a given reason; the other sides shall have also to the other sides given reasons.

Demonstr. For seeing that the reason of BD to FH is given, and

a 3. def. also that a of BD to BA , b the reason of the
 b 8. prop. said AB to FH is given. But the reason of the
 c 3. def. same FH to FE is also given: Therefore d
 d 8. prop. the reason of AB to EF is given. In like manner also the reasons of the other sides to the other sides are given.

PROP. LIV.



If two figures A and B , given by kind, have to one another a given reason, also their sides shall be to one another in a given reason.

Constr. For either the figure A is alike and alike posited to B , or is not: Let it in the first place be alike, and alike posited; and let there be taken the line G , a third proportional to the lines CD and EF .

Demonstr.

Demonstr. As CD is to G, so is A to B. *a cor. 19,*
But the reason of A to B is given; therefore *20. 6.*
also the reason of CD to G is given. And
seeing that CD, EF, and G, are proportional,
also the reason of CD to EF is given. But *b 24. prop.*
A and B are given by kind: Therefore the other sides shall have given reasons to the other sides. *c 53. prop.*

Now let the figure A be not alike to the figure B, and let there be described on EF the figure EH, alike and alike posited to A: Therefore the figure EH is given by kind; but the figure B is also given by kind: Therefore the reason of B to EH is given; and therefore the reason of A to the same EH is also given: But A is alike to EH: Therefore (by what is abovesaid) the reason of CD to EF is given; and in like manner the reason of the other sides to the other sides is given. *d 49. prop. e 8. prop.*

OTHERWISE.



which figures M and N shall be consequently given by kind.


Demonstr. Therefore seeing that as CD is to EF, so is GH to LK, and that on those lines CD, EF, GH, and LK, are described the figures
D d A, B,

f 22. 6. A, B, M, and N, alike and alike posited; *f* as A is to B, so is M to N. But the reason of A to B is given: Therefore the reason of M to N is given. But *g* M is given, considering that it is a figure given by kind, described on a right line given by magnitude; therefore N is also given.

h sch. 52. *Constr.* 2. Now, on LK let the square O be described: Therefore *h* the figure O is given by kind.

Demonstr. 2. Wherefore the reason of O to N is given. But N is given: Therefore O is given; and consequently, *i* also KL. But *i* sch. 52. GH is given: Therefore *k* the reason of GH to KL is given. But as GH is to LK, so is CD to EF. Therefore the reason of CD to EF is given; and therefore the figures A and B being given by kind, *l* the other sides of the same figures shall also have to the other sides given reasons. But if the figures be not alike, the latter part of the demonstration here above must be observed.

PROP. LV.




If a space A be given by kind, and by magnitude, the sides thereof shall be given by magnitude.

Constr. For, let the right line BC given by position and by magnitude, be exposed; and thereon let there be described the space D, alike and alike posited to A; therefore the said space D is given by kind.

Demonstr. For that it is described on the line BC, given by magnitude, it is also *a* given by magnitude. But the figure A is also given: There-

Therefore *b* the reason of A to D is given. *b* 3. *prop.*
 But those figures A and D are given by kind:
 Therefore *c* the reason of the line EF to the *c* 54. *prop.*
 line BC is given. But BC is given: Therefore
d EF is also given. But the reason of the *d* 3. *def.*
 same EF to FG is given: Therefore *e* FG is *e* 2. *prop.*
 given. And by the same reasons it may be
 demonstrated that each of the other sides are
 given by magnitude.


OTHERWISE.



Constr. Let the
 space GHIKL be
 given by kind and
 by magnitude: I
 say that the sides
 thereof are given
 by magnitude. For
 on the right line
 GH let there be
 described the square
 GM: therefore *f* *f* *sch.* 52.
 GM is given by *prop.*
 kind.

Demonstr. But the
 space GHIKL is
 also given by kind:
 Therefore *g* the reason of the same space GK *g* 49. *prop.*
 to GM is given. But GK is given by magni-
 tude: Therefore *h* GM is also given by magni- *h* 2. *prop.*
 tude; and seeing that GM is the square of the
 line GH, *i* that line GH is given by magni *i* *sch.* 52.
 tude. Wherefore in like manner, each of the *prop.*
 other lines HI, IK, KL, and LG, is given.

PROP. LVI.



given reason that the parallelogram *A* hath to the parallelogram *B*.

Constr. For let *HD* be prolonged directly to *L*, so that as *CD* is to *FG*, so *HD* may be to *DL*; and finish the parallelogram *DK*.

- Demonstr.* Seeing that as *CD* is to *FG*, so *HD* is to *DL*, and *a* that *CD* is equal to *KL*; as *LK* is to *FG*, so is *GE* to *DL*; and thus the sides about the equal angles *DLK* and *EGF* are reciprocally proportional: Wherefore *b* *DK* is equal to *B*; and therefore seeing the reason of *A* to *B* is given, and that *B* is equal to *DK*, the reason of *A* to *DK* is given. But as *c* *A* is to *DK* (that is to *B*) so is *HD* to *DL*; therefore the reason of *HD* to *DL* is also given: and seeing that as *CD* is to *FG*, so *GE* is to *DL*, and that the right line *HD* hath to *DL* a given reason; to wit, that which the space *A* hath to the space *B*; as *CD* is to *FG*, so *GE* is to that to which *HD* hath the given reason that the space *A* hath to the space *B*, that is to say, the reason of *HD* to *DL*.


PROP.

PROP. LVII.

If a given space AD be applied to a given right line AB in a given angle CAB , the breadth CA of the application is given.

Constr. For on AB , let there be described the square AF ; therefore a the same AF is *a sch. 52.* given: Let the lines EA , FB , and CD , be *prop.* prolonged to the points G and H .


Demonstr. Seeing therefore that each space AD and AF is given, their reason is also given. But b AD is equal to AH : Therefore the reason of AF to AH is given: Wherefore the reason of EA to AG is given. (for c it is the *c 1. 6.*



same with that of AF to AH .) But EA is equal to AB ; therefore the reason of AB to AG is given. Now seeing that the angle CAB is given, and the angle GAB also given, the residue CAG is given. But the angle CGA is also gi-

ven, being a right angle: Therefore the remaining angle ACG is given. Wherefore the triangle d CAG is given by kind. Therefore *d 4c. prop.* the reason of CA to AG is given. But the reason of AB to the same AG is also given: Therefore the reason of CA to AB is given; and the said AB is given: Wherefore CA is also given.

PROP. LIX.




If a given space AB be applied according to a given right line AC , exceeding it by a fi-

gure CB given by kind, the breadths of the excesses CE and CF are given.

Constr. For DE being divided into two equal parts in G , let there be described on GE the rectiline figure GH , alike and alike posited to CB .

Demonstr. Now seeing that CB is alike to GH , those figures CB and GH * are about one and the same diameter, and GH is given by kind, as is CB . But it is described on the given line GE : Therefore *a* the same GH is also given by magnitude. But AB is given: Therefore AB and GH are given by magnitude. Now those figures AB and GH , are equal to LI , (for AG , LE , and EI , being equal, the Gnomon GFH is equal to AB ; and therefore adding GH common to both, LI shall be equal to AB and GH ;) therefore LI is given by magnitude; but it is also given by kind, being it is *b* alike to CB . Therefore *b* 24. 6. *c* the sides of the said LI are given, seeing *c* 55 prop. it is equal to GE : Therefore *d* the remainder *d* 4. prop. CF is given, and in a given reason *e* to CE . *e* 3. def. Wherefore *f* CE is given. *f* 2 prop.

Scholium.



* EUCLIDE sup-
poseth here that CB
and GH are about one
and the same diameter,
but we shall thus de-
monstrate it: Let CB
and GH be two alike
parallelograms disposed
as above, that is to
say, that the equal
angles join together in

E, the side CE meets directly with his homologal
side EH, and the side BE, his correspondent side
EG; and let the diameter FE be drawn, I say
that the said diameter FE prolonged, will pass by
the point K; that is to say, the parallelograms GH
and CB, consist about one and the same diameter.
For if it be denied, the diameter EF being pro-
duced, will pass above the point K, or below it.
Let it in the first place pass above it, and let it cut
GK, prolonged in the point M, and by the point M,
let there be drawn MN, parallel to KH, which
shall meet EH, prolonged in the point N, and FB
in O.

Demonstr. Forasmuch as the parallelograms
GN and CB are with the parallelogram LO
about one and the same diameter, they are
alike to one another. Wherefore as FC is to
CE, so is EG to GM. In like manner, seeing
the parallelograms CB and GH are alike, as
FC is to CE, so is EG to GK; Therefore
as EG is to GM, so is EG to GK. Where-
fore GM and GK are equal, a part to the
whole, which is absurd; By the same rea-
sons it may be demonstrated; that the diame-
ter prolonged will not fall below the point K:
There-

g 24. 6.

h 11. 5.

i 9. 5.

Therefor
sist abo

F

G

CE and
Dem
the G
rall
by kin
a the
theref
given
given
CE an


F

G

L
and
figu

Therefore the parallelograms CB and GE consist about one and the same diameter.


PROP. LX.



If a parallelogram AB, given by kind and by magnitude, be augmented or diminished by a Gnomon CFD, the breadths of the Gnomon (consisting of the lines

CE and DG) are given.

Demonstr. For seeing that AB is given, and the Gnomon CFD also given, the whole parallelogram BF is given: But it is also given by kind, seeing it is alike to BA: Therefore the sides of the same BF are given; and a 55. prop. therefore each of the lines BE and BG is given. But each of the lines BC and BD is given; therefore each of the remaining lines CE and DG is also given.




Constr. Now let the parallelogram BF, given by kind and by magnitude, be diminished by the given Gnomon CFD: I say that each of the lines CE and DG is given.

Demonstr. 2. For seeing that BF is given, and the Gnomon CFD given, the remaining figure AB is also given. But it is also given by

by kind, seeing it is alike to BF : Therefore
 b 55. *prop.* b the sides of the said AB are given, and there-
 fore each of the lines CB and BD is given.
 But each of the lines BE and BG is given :
 Therefore also each of the remaining lines CE
 and DG is given.

PROP. LXI.



If to one side of a figure ABCE, given by kind, there be applied a space parallelogram CD, in a given angle BCF, and that the given figure AC hath to the parallelogram CD a given reason, the parallelogram CD is given by kind.

Constr. For by the point B, let BH be drawn parallel to CE, and by the point E let EH be drawn parallel to CB, and let EC and HB be prolonged to the points K and G.

Demonstr. Forasmuch as the angle BCE is given, and the reason of EC to CB, a the parallelogram CH is given * by kind. But the figure ABCE is also given by kind, and is described on the same line BC, as the parallelogram CH given by kind is : Therefore b
 b 49. *prop.* the reason of the figure ABCE to the parallelogram CH is given. But by supposition, the reason of the said figure ABCE to the parallelogram CD is also given; and CD is c
 c 36. 1. equal to CG : Therefore d the reason of CH to CG is given. Wherefore the reason of
 d 8. *prop.* the line EC to the line CK is given; (for e
 e 1. 6. as CH is to CG, so is EC to CK.) But the reason of EC to CB is also given : Therefore
 f 8. *prop.* f the reason of the said CB to CK is given.
 And

And see
 the follo
 angle B
 the rem
 angle C
 to the
 CFK is
 is give
 of FC
 to the
 the res
 angle
 parallelog

* Al
 hath o
 about
 kind,
 withsta
 demon

See
 ECB
 the ri
 CB,
 same
 fore
 other
 of E
 to C
 one

And seeing that the angle ECB is given, also the following angle BCK *g* is given. But the *g* 13. r. angle BCF is proposed given; and therefore 4. prop. the remaining angle FCK is given. Also the angle CKF is given, for that *h* it is equal *h* 29. r. to the angle BCK: Therefore the other angle CFK is given: Wherefore *i* the triangle FCK *i* 40. prop. is given by kind; and therefore the reason of FC to CK is given. But the reason of CB to the same CK is also given: Therefore *k* 8. prop. the reason of FC to CB is given; and the angle BCF is also given. Wherefore the parallelogram CD is given by kind.


Scholium.

* *Altho' it be manifest that a parallelogram that hath one angle given, and the reason of the sides about the same angle also given, is given by kind, as Euclide doth here declare, so it is notwithstanding that the antient Interpreter doth thus demonstrate it.*

Seeing that in the parallelogram CH the angle ECB is given, the angle CEH is also given; for the right line EC falling on the parallels EH and CB, doth make the two internal angles on the same part equal to two right angles. And therefore seeing that the angle ECB is given, the other angles are given; and seeing that the reason of EC to CB is given, and that BH is equal to CE, and EH to BC, the reason of the sides to one another is also given.

PROP.

PROP. LXII.



If two
right lines
AB and
CD, have
to one
another a
given rea-
son, and
that on
one of
them AB,
there be


described a figure AEB, given by kind; but on the other CD, a space parallelogram DF, in a given angle DCF, and that the figure AEB hath to the parallelogram DF a given reason, the parallelogram DF is given by kind.

Constr. For on the line AB let there be described the parallelogram AH, alike and alike posited to DF.

Demonstr. Seeing then that the reason of AB to CD is given, and that on those lines are described the rectiline figures AH and FD, alike and alike posited, *a* the reason of AH to FD is given. But the reason of FD to AEB is also given: Therefore *b* the reason of AH to AEB is given. But the angle ABH is also given, being equal to the angle FCD, and so the figure AEB is given by kind; and to AB one of the sides thereof, the parallelogram AH is applied in a given angle ABH, and the reason of the said figure AEB to the said parallelogram AH is given: Therefore *c* the parallelogram AH is given by kind; and therefore FD which is alike thereto, is also given by kind.

PROP.

P R O P. LXIII.




If a triangle ABC be given by kind, the square BE , CD , and CF , which is described on each of the sides, shall have a given reason to the triangle ABC .

Demonstr. For seeing that on one and the same right line BC , there are

described the two rectiline figures ABC and CD , given by kind, *a* the reason of the same ABC to CD is given; and therefore the reason of the squares BE and CF , to the triangle ABC , is also given. *a 49. prop.*

P R O P. LXIV.



If a triangle ABC , hath an obtuse angle ABC given, that space of which the side AC subtending the obtuse angle ABC , is more in power than the sides AB and BC , that comprehend the said angle, shall have a given reason to the triangle ABC .

Constr. Let the line CB be prolonged directly, and from the point A let the perpendicular AD be drawn: I say

a 12. 2.

say that the space of which the square of the line AC doth exceed the squares of the lines AB and BC, that is to say, *a* the double of the rectangle contained under CB and BD, shall have a given reason to the triangle ABC.

b 40. prop.

c 3. def.

d 1. 6.

e 41. 1.

f 8. prop.

Demonstr. For seeing that the angle ABC is given, the angle ABD is also given. But the angle ADB is also given; therefore the other angle BAD is given: Wherefore *b* the triangle ABD is given by kind; therefore *c* the reason of AD to DB is given. But as AD to DB, so *d* the rectangle of AD and BC is to the rectangle of BC and BD. But the reason of AD to BD is given: Therefore also is the reason of the rectangle of AD and BC to the rectangle of BC and BD given: Wherefore the reason of the double of the said rectangle BC and BD to the rectangle of AD and BC is also given. But the said rectangle of AD and BC hath also a given reason to the triangle ABC (to wit, double reason; for the rectangle is *e* double to the triangle) therefore the reason of the double of the rectangle of BC and BD *f* to the triangle ABC is given. But the same double of the rectangle of CB and BD is that space of which the square of the line AC doth exceed the squares of the lines AB and BC: Therefore the same space hath a given reason to the triangle ABC.


PROP.



line AC
of wh
than t
that is
of BC
triangl

Dem
given,
other
triang
fore t
consec
BC a
There
angle
and
rectan
is gi
triang
ble c
trian
same
is th
less
that

PROP. LXV.




If a triangle ABC , hath one acute angle ACB given, that space, of which the side subtending the said acute angle is less in power than the sides comprehending the same acute angle, shall have a given reason to the triangle.

Constr. From the point A let there be drawn the line AD , perpendicular to BC : I say that space of which the square of the line AB is less than the squares of the lines AC and CB , that is to say, *a* the double of the rectangle of BC and CD , hath a given reason to the triangle ABC . 13. 2.

Demonstr. For seeing that the angle C is given, and the angle ADC also given, the other angle DAC is given: Wherefore the triangle b ADC is given by kind; and therefore the reason of AD to DC is given, and consequently also *c* that of the rectangle of BC and CD to the rectangle of BC and AD : 40. prop.
Therefore the reason of the double of the rectangle of BC and CD to the rectangle of BC and AD is given. 1. 6. But the reason of the same rectangle of BC and AD to the triangle ABC is given (for *d* the rectangle is double to the triangle:) Therefore *e* the reason of the double of the rectangle of BC and CD to the triangle ABC is given. 41. 1.
And seeing that the same double of the rectangle of BC and CD is that whereof the square of the line AB is less than the squares of the lines AC and BC , that space of which the square of the line AB 8. prop.

AB is less than the squares of the lines AC and BC, shall have a given reason to the triangle ABC.

PROP. LXVI.




If a triangle ACB, hath one angle B given, the rectangle made of the lines AB and BC, containing the same angle, shall have a given reason to the triangle.

Constr. For from the point A let AD be drawn perpendicular to CB.

- Demonstr.* Therefore seeing that the angle B is given, and also the angle ADB; the other angle BAD is likewise given. Wherefore the triangle ADB *a* is given by kind; and consequently the reason of AB to AD is given.
- a* 40. *prop.* But as AB is to AD, *b* so the rectangle of AC and CB is to the rectangle of CB and AD: Therefore the reason of the rectangle of AC and CB to the rectangle of CB and AD is given. But the reason of the said rectangle of CB and AD to the triangle ACB is also given; (for that it is double reason, the rectangle being double *c* to the triangle:)
- c* 41. *prop.* Therefore *d* the reason of the rectangle of AC and CB to the triangle ABD is given.
- d* 8. *prop.*

PROP

PROP. LXVII.



If a triangle ABC hath one angle BAC given, that space by which the square of the line compounded of the two sides BA and AC , that contain the same given angle BAC doth exceed the square of the other side, it shall have a given reason to the triangle ABC .

Constr. For let BA be prolonged in such sort as that AD may be equal to AC , then having drawn DCE infinitely, from the point B let BE be drawn parallel to AC , meeting the said DE in the point E .

Demonstr. Forasmuch as AD is equal to AC , DB is equal to BE ; (for the two triangles ADC and BDE are alike) and from the top B is drawn to the base DE , the right line BC : Therefore * the rectangle of DC and CE , with the square of BC , is equal to the square of BD ; but the same BD is compounded of BA and AC ; therefore the square of the compound of AB and AC is greater than the square of BC , of the rectangle of DC and CE .

Now I say that the rectangle of DC and CE hath a given reason to the triangle ABC : Forasmuch as the angle BAC is given, the angle DAC is also given. But each of the angles ADC and ACD is given, it being the half of the angle BAC which is given. Therefore b the triangle ADC is given by kind; and therefore the reason of DA to DC is given. Therefore c the reason of the square of the said DA to the square of DC is also given.

E e


given.

d 2. 6.

given. And seeing that as BA is to AD, d

e 1. 6.

f 1. 6.



so is EC to CD, and
also as BA is to AD,
e so is the rectangle of
BA and AD to the
square of AD; and as
EC is to CD, f so also
is the rectangle of EC
and CD to the square
of CD; by permutation,
as the rectangle of BA
and AD is to the rect-
angle of EC and CD,

so is the square of AD to the square of DC.
But the reason of the said square of AD to
the square of DC is given: Therefore the rea-
son of the rectangle of BA and AD to the
rectangle of EC and CD is also given. But
AD is equal to AC: Therefore the reason of
the rectangle of BA and AC to the rectangle
of EC and CD is given. But the reason of
the rectangle of BA and AC to the triangle
g 6. prop. ABC g is given, because the angle BAC is
h 8. prop. given: Therefore h the reason of the rectangle
EC and CD to the triangle ABC is given.
But the rectangle of EC and CD is that
whereof the square of the line compounded of
BA and AC is greater than the square of BC:
Therefore that space to which the square of
the line compounded of BA and AC is greater
than the square of BC, shall have a given
reason to the triangle ABC.

Scholium.

* EUCLIDE supposeth in this place that when
in an Isosceles triangle a right line is drawn
from the top to the base, the square of that line,
with the rectangle contain'd under the segments of
the

the base
the other
thus dem

Constr
whose
A let
that th
BD and
of the

Demo




angle
shall
But
squa
of A
DC,
N
but
BC
shal
poi
rect
DE
add
gle
and
and
th

the bases, is equal to the square of either of the other legs, which the antient interpreter doth thus demonstrate.

Constr. Let ABC be an Iſosceles triangle, whose legs are AB and AC; and from the top A let AD be drawn to the base BC: I ſay that the ſquare of AD with the rectangle of BD and DC, is equal to the ſquare of either of the legs AB or AC.

Demonſtr. Now the line AD is perpendicular to BD, or not: Let



it in the firſt place be perpendicular: Therefore it will cut the baſe BC into two equal parts in the point D; and therefore the rectangle contained under BD and DC is equal to the ſquare of the ſaid BD, and adding to them the common ſquare of AD, the rect-

angle of BD and DC with the ſquare of AD, ſhall be equal to the ſquares of DB and AD. But to thoſe ſquares of AD and DB i the ſquare of AB is equal: Therefore the ſquare of AB is equal to the rectangle of BD and DC, and the ſquare of AD together. 47. 1.

Now ſuppoſe AD not to be perpendicular, but that from the point A there doth fall on BC the perpendicular AE, that being ſo, BC ſhall be cut into two parts equally in the point E, and unequally in D. Wherefore the rectangle of BD and DC, with the ſquare of DE, k is equal to the ſquare of BE; and k 5. 2. adding the common ſquare of AE, the rectangle of BD and DC, with the ſquares of DE and AE, ſhall be equal to the ſquares of BE and AE. But l the ſquare of AD is equal to l 47. 1. the two ſquares of DE and AE: Therefore

E e 2


the

the rectangle of BD and DC, with the square of AD is equal to the squares of BE and AE. But to these squares of BE and AE the square of AB is equal: Therefore the square of AD, with the rectangle of BD and DC, is equal to the square of AB.

OTHERWISE.

Constr. Having done, as in the foregoing Demonstration, from the point A, let AF be drawn perpendicular to CD, and let AE be drawn.

Demonstr. Forasmuch as the angle BAC is given, the half thereof ACF shall be also given. But the angle AFC is given; and therefore the triangle AFC is given by kind: Therefore the reason of AF to FC is given. But the reason of CD to the same FC is also given, seeing that CD is double to FC: Therefore *m* the reason of CD to AF is given; and therefore also the reason of the rectangle of CD and EC, to the rectangle of AF and EC, is given; (for it is the same reason *n* as that of CD to AF.)



o 41. 1.

p 37. 1.

But the triangle ACE is equal to the triangle ABC *p*, they being both constituted on one and the same base AC, and between the same parallels


parallel
of the
gle A
CE a
of the
great
space
pound
square
angle



great
rectan
square
and
pound
square
gle o
reason
to the
Co
acute
draw
De
an C
AD

parallels AC and BE: Therefore *q* the reason *q* 8. *prop.* of the rectangle of CE and CD to the triangle ABC is given. But the said rectangle of CE and CD is the space by which the square of the line compounded of AB and AC, is greater than the square of BC: Therefore that space by which the square of the line compounded of AB and AC is greater than the square of BC, hath a given reason to the triangle ABC.

OTHERWISE.



For the given angle A is either a right, acute, or obtuse angle: Let it in the first place be supposed a right angle: Therefore the square of the line compounded of BAC, is

greater than the square of BC, by twice the rectangle of BA and AC; (seeing that *r* the *r* 47. 1. square of BC is equal to the squares of BA and AC; and the square of the line compounded of BAC *s* is equal to those two *s* 5. 2. squares of BA and AC, and twice the rectangle of the said BA and AC:) Wherefore the reason of double the rectangle of BA and AC to the triangle ABC is given.


Constr. Now let the angle C be supposed acute, and from the point A let there be drawn on CB the perpendicular AD.

Demonstr. Forasmuch as the triangle CAB is an Oxigonium triangle, and the perpendicular AD being drawn, the square of CA and CB

t 13. 5.

are equal *t* to the square of AB with twice the rectangle of CB and CD; adding therefore the common double rectangle of CA

u 4. 2.



x 1. 2.

and CB, the squares of CA and CB, with the double rectangle of the said CA and CB, that is to say, *u* the alone square of the line compounded of ACB, are equal to the square of AB, with the double of the rectangle of CD

y 40 prop.


z 6. prop.

a 1. 6.

b 66. prop.

and CB, and over and above the double of the rectangle of AC and CB, that is to say, the double of the rectangle contained under the compound line of ACD and CB (for the rectangle of ACD and CB is *x* equal to the rectangles of AC and CB, and of CD and CB :) Therefore the square of the line compounded of ACB is greater than the square of AC, by double the rectangle of ACD and CB. And seeing that the angle ACB is given, and the angle BDA also given, the other angle CAD is given: Therefore *y* the triangle CAD is given by kind, and therefore the reason of CD to CA is given, and by consequence the reason of the line compounded of ACD to CA *z* is also given. Wherefore the reason of the rectangle of those lines compounded of ACD and CB *a* to the rectangle of AC and CB is also given. But the reason of the said rectangle of AC and CB to the triangle CAB *b* is given, seeing the angle C is given; therefore the reason of double the rectangle of the line compounded of ACD and CB to the triangle CAB is given.

Lastly,



Lastly, Let the angle BAC be supposed to be obtuse, and having prolonged BA from the point C, let the perpendicular CE be drawn on the said line BA prolonged; and let AF be pro-

posed to be equal to AE.

Demonstr. Forasmuch as the angle BAC is obtuse, and the perpendicular CE being drawn, the squares of AB and AC, and the double of the rectangle under BA and AE, or AF, are all alike equal *c* to the square of BC, and *c* 12. 2. adding the common double rectangle of BA and AC, the squares of the said AB and AC, with the double of the rectangle of the same AB and AC, that is to say, *d* the square of *d* 4. 2. the line compounded of BAC, and the double of the rectangle of BA and AF are together equal to the square of BC, with the double of the rectangle of BA and AC. Let the common double of the rectangle of BA and AF be taken away, and there will remain the square of the line compounded of BAC, equal to the square of BC, with the rectangle of AB and CF; (for the rectangle of AB and AC is equal *e* to the two rectangles of *e* 1. 2. AB and AE, and of AB and CF :) Therefore the square of the line compounded of BAC is greater than the square of BC by the double of the rectangle of AB and CF. And forasmuch as the angle BAC is given, the angle CAE *f* is given. But the angle AEC is *f* 13. 11. also given; therefore the other angle ACE is given: Wherefore *g* the triangle ACE is given *g* 40. *prop.* by kind, and therefore the reason of CA to AE, that is to say, to AF is given. Therefore *h* the reason of the said CA to FG is *h* 5. *prop.*


E e 4 also

- also given. But the reason of the same CA to CE is given; therefore the reason of CE to CF is also given. Wherefore the reason of the rectangle of EC and AB to the rectangle of FC and AB is given; (for the rectangle is to the rectangle *k* as CE is to CF) and also that of the rectangle of AC and AB, to the rectangle of EC and AB. Therefore the reason of the rectangle of FC and AB to the rectangle of AC and AB is given. But the reason of the rectangle of AC and AB to the triangle ABC *m* is given: Therefore also the reason of the double of the rectangle of FC and AB, to the triangle ABC is given. But the same double of the rectangle of FC and AB is that, whereof the square of the line compounded of BAC is greater than the square of BC, whereof that space of which the square of the line compounded of BAC is greater than the square of BC, hath a given reason to the triangle ABC.

OTHERWISE.

Constr. Let the line BA be prolonged to the point D, in such sort as AD may be equal to AC, and let CD be drawn.


- Demonstr.* Forasmuch as the angle BAC is given, each of the angles ADC and ACD, which is the half thereof shall be also given; and therefore the other angle, DAC is also given: Therefore *n* the triangle ACD is given by kind. Wherefore the reason of AC to CD is given. And forasmuch as the an-



gle ADC is given: Let each of the angles DEC and AFC be made † equal to the

the said
gle BDC
angle DE
and DBC
the other
angle BD
Therefore
Wherefore
is to say
with th
of BD,
line con
to AC
and CE
say, th
BAC i
angle
the rea
CE to
much
gle BC
angle
to the
is also
the r
remain
gle A
and th
to C
CD,
to C
of A
let A
as mu
angl
is g

the said ADC : Therefore seeing that the angle BDC is equal to the angle DEC, and the angle DBE is common to the triangles DBE and DBC, the other angle BDE is equal to the other angle BCD ; and therefore the triangle BDE is equiangled to the triangle BDC. Therefore *o* as EB is to BD, so is BD to CB : *o* 4. 6. Wherefore the rectangle of EB and CB, that is to say, *p* the rectangle of EC and CB, *q* *p* 5. 2. with the square of CB is equal, *r* to the square *q* 5. 2. of BD, that is to say, to the square of the *r* 17. 6. line compounded of BAC ; for AD is equal to AC ; and therefore the rectangle of EC and CB, with the square of CB, that is to say, the square of the line compounded of BAC is greater than the square of the rectangle of BC and CE : I say therefore that the reason of the said rectangle of BC and CE to the triangle ABC is given. Forasmuch as the angle BDE is equal to the angle BCD, and the angle ADC equal to the angle ACD, the other angle CDE is equal to the other angle ACB : But the angle DEC is also equal to the angle AFC ; therefore the remaining angle CAF is equal to the remaining angle DCE. Wherefore the triangle AFC is equiangled to the triangle DCE ; and therefore *s* as CA is to AF, so is CD *s* 4. 6. to CE ; and by permutation, as AC is to CD, so is AF to CE. But the reason of AC to CD is given : Therefore also the reason of AF to CE is given. From the point A let AH be drawn perpendicular to BC : Forasmuch as the angle AFC is given, and the angle AHF also given, the third angle HAF is given : Wherefore *t* the triangle AHF is *t* 40. prop. given



u 8. prop.

x i. 6.

y 41. 1.

But the reason of AF to CE is also given; Therefore *u* the reason of AH to CE is given; and therefore the reason of the rectangle of AH and BC *x* to the rectangle of BC and CE is also given. But the reason of the rectangle of AH and BC to the triangle ABC is likewise given; (for the rectangle *y* is double to the triangle) and the rectangle of BC and CE is that whereof the square of the line compounded of BAC is greater than the square of BC. Therefore that space of which the square of the line compounded of BAC is greater than the square of BC by a given reason to the triangle ADC.

Scholium.

† The antient Interpreter pretending to shew the construction of the angle DEB equal to the angle ADC, saith that on the line BD and in the point D, the angle BDE ought to be made equal to the angle BCD, and that the right lines BC and DE be drawn until they intersect in E, in such sort as he supposeth the angle BCD, to be given, but it is not.

The same Interpreter afterward shews how there may universally from a given point, be drawn a right line given by position to a right line, making an angle equal to a given angle. But we will also reject this way, seeing we have elsewhere shewn

given by
kind;
and by
consequence
the reason
of AF to
AH is
given.

shewn and
if we w
given b
equal to
we have
in the
CKL eq
KL do
line re
the poi
rallel t
E, and
angle
DE,
the an
hath
and b
equal

If
anoth
a giv
likew

Co
ven
ven.
appl
para
fort
† T
right

shewn another more brief and easy. For example,
 if we would from the point D draw to the line BC
 given by position a right line, making an angle
 equal to a given angle ADC , as is here required,
 we have no more to do but to assume the point K
 in the said line BC , and there make the triangle
 CKL equal to the given angle ADC : If the line
 KL doth meet with the point D , it shall be the
 line required. But if it meet not with it, from
 the point D let there be drawn the line DE , pa-
 rallel to the said KL , cutting BC prolonged in
 E , and the angle DEC shall be equal to the given
 angle ADC , for on the two parallel lines LK and
 DE , there doth fall the line BE ; and therefore
 the angle DEC is equal to the angle LKC , which $\angle$ 29. 1.
 hath been made equal to the given angle ADC ;
 and by consequence the same angle DEC is also
 equal to ADC .


P R O P. LXVIII.

If two parallelograms AB and CD , have to one
 another a given reason, and that a side hath also
 a given reason to a side, the other side shall have
 likewise a given reason to the other side.

Constr. Let the reason of BE to FD be gi-
 ven: I say the reason of AE to FC is also gi-
 ven. For to the right line EB let there be
 applied the parallelogram FH , equal to the
 parallelogram CD , and constituted in such
 sort as AE and EG may make one right line:
 † Therefore KB and BH will also make one
 right line.

Demonstr.

a 1. 6.



b 14. 6.

therefore the reason of AE to EG is also given. Seeing therefore that EH is equal and equiangled to CD, as *b* EB is to FD, so is FC to EG. But the reason of EB to FD is given: Therefore also the reason of FC to EG is given. But the reason of AE to the same EG is also given: Therefore the reason of AE to FC is given.

Scholium.

c 13. 7.

d 29. 1.

† EUCLIDE having posited AE and EG directly in one right line, presently concludeth that KB and BH shall also make a right line; but we shall demonstrate it thus. Seeing the lines AB and EG are posited directly, the angles AEB and BEG *c* are equal to two right angles; and seeing that AB is a parallelogram, the lines AK and EH are parallels, on which the line AE doth fall; and therefore the two internal angles A and BEA *d* are also equal to two right angles, and taking away the common angle BEA, there will remain the angle A, equal to the angle BEG; and consequently their opposite angles EBK and H are also equal to one another: Again, seeing that BG is a parallelogram, the two lines BE and HG are parallels, on which BH doth fall; and therefore the two internal angles H and EBH *d* are equal to two right angles. But it hath been demonstrated that


Demonstr.

Forasmuch as the reason of AB to CD is given, and that EH is equal to the said CD; the reason of AB to EH is given; and

that H is
gles EBA
angles;
BH do

Constr.
fed, and
ven, let
fore th

Demo
also gi
CD to
K to M
given.



to E
of K
M to
CD
with
and
the
CD
fore
sam
M
of

that H is equal to EBK : Therefore the two angles EBK and EBH are also equal to two right angles; and therefore the two lines KB and BH do meet directly according to EUCLIDE.

OTHERWISE.

Constr. Let the given right line K be exposed, and seeing that the reason of A to B is given, let the same be made of K to L ; therefore the reason of K to L is also given.

Demonstr. But K is given; therefore L is *prop. 2.* also given. Again, seeing that the reason of CD to EF is given, let the same be made of K to M : Therefore the reason of K to M is given. But K is given, therefore M is also *prop. 1.*



given; and therefore the reason of L to M is given. Now seeing that A is equiangular to B , the reason of the said A to B is compounded of that of the sides, that is to say, of CD

to EF , and of CG to EH . But also the reason of K to L is compounded of K to M , and of M to L ; therefore the reason compounded of CD to EF , and of CG to EH , is the same with that which is compounded of K to M , and of M to L (the reason of K to L being the same as of A to B :) But the reason of CD to EF is the same as of K to M : Therefore the other reason of CG to EH is also the same as of M to L . But the said reason of M to L is given: Therefore also the reason of CG to EH is given.

PROP.

PROP. LXIX.



If two parallelograms, CB and EH, having the angles D and F given, and that a side hath also a given reason

to a side; in like manner the other side shall have a given reason to the other side.

Constr. Let the reason of BD to FH be also given: I say that the reason of AB to EF is given. For if CB be equiangled to HE, it is manifest by the precedent Proposition; but if it be not equiangled thereto, let the right line DB be constituted, and in the given point B therein, let the angle DBK be made equal to the angle EFH, and finish the parallelogram DK.

Demonstr. Forasmuch as each of the angles BKL and BAK is given, † the other angle KBA is given: Wherefore the triangle a ABK is given by kind; and therefore the reason of AB to BK is given. But the reason of CB to EH is supposed to be given, and b CB is equal to DK; therefore the reason of DK to EH is given; and seeing that DK is equiangled to EH, and the reason of the said DK to EH is c 68. prop. given, as also that of DB to FH, c the reason of BK to FE is given. But the reason of the said BK to BA is also given: Therefore d the reason of AB to FE is given.

Scholium.

† EUCL
gram havin
gles are als
preters as o
gle F bein
given, it b
for that o
doth fall i
ternal ang
to two rig
posite ang
are also g
From
F being
of the t
given: T
equal to
to the
angle B
BDC, i
the rem
ven; in
angles
doth de

If
about
gles
anoth
CB to
EH
Co
on t
be a

Scholium.

† EUCLIDE supposeth there that a parallelogram having one angle given, all the other angles are also given, and as well the antient Interpreters as others, do give the reasons why, the angle F being given, the other angle E shall be also given, it being the remainder of two right angles, for that on the parallel lines EG and FH there doth fall the line EF , which makes the two internal angles (of the same part) F and G , equal to two right angles. But to those angles f the opposite angles G and H are equal, and therefore they are also given.

From whence it follows that the angles BDC and F being given by supposition, all the other angles of the two parallelograms CB and EH , are also given: Therefore the angle DBK having been made equal to the angle F , the angle K shall be equal to the angle E , and given as that is: But the angle BAL which is opposite to the given angle BDC , is also given; and therefore BAK which is the remainder of two right angles, shall be also given; in such sort as in the triangle ABK , the two angles BAK and BKA are given, as EUCLIDE doth declare in this place.


PROP. LXX.

If of two parallelograms AB and EH , the sides about the equal angles, or about the unequal angles (yet nevertheless given angles have to one another a given reason, to wit (AC to EF , and CB to FH) also the same parallelograms AB and EH shall have to one another a given reason.

Constr. For let AB be prolonged to EH , and on the right line CB let the parallelogram CM be applied equal to the parallelogram EH , in such

such sort as AC may be direct to CN; that is to say, that AC and CN make one right line; and by consequence DB shall be a direct-ly with BM.

Demonstr. Forasmuch then as CM is equi-angled and equal to EH, the sides about the equal angles shall be reciprocally *b* proportion-
b 14. 6. al: Wherefore as BC is to HF, so is FE to



NC. But the reason of BC to HF is given: Therefore the reason of FE to NC is also given.
c 8. *prop.* But the reason of AC to the same EF is given: Therefore *c* the reason of AC to NC is also given. Wherefore the reason of AB to CM is given; (for it is the same *d* as of AC to CN.) But CM is equal to EH: Therefore the reason of AB to EH is given.

d 1. 6.

Constr. Now suppose AB not to be equi-angled to EH, and on the right line CB, and in the given point C therein: Let there be constituted the angle BCK, equal to the given angle F, and so finish the parallelogram CL.

Demonstr. Forasmuch as the angle ACB is given, and the angle BCK also given, the remaining angle ACK is given: Therefore the triangle ACK *e* is given by kind; and therefore the reason of AC to CK is given: But the reason of AC to EF is also given: Therefore the reason of CK to EF is given. But the reason of BC to HF is also given, and the angle BCK is equal to the angle F; there-fore

e 40. *prop.*


fore (b)
reason
CL, A
AB to

less g
son (c
triang
reason

Co
be fi
D
AG
ang
ang
son
lelo
ven
the
the
the
gle

fore (by the first part of this Proposition) the reason of CL to EH is given. But to the said CL, AB is equal: Therefore the reason of AB to EH is given.

P R O P. LXXI.




If of two triangles ABC and DEF, the sides about the equal angles A and D, or else about the unequal angles (yet neverthe-

less given angles) have to one another a given reason (to wit AB to DE and AC to DF) the same triangles shall have also to one another a given reason ABC to DEF.

Constr. Let the parallelogram AG and DH be finished.

Demonstr. Seeing that the two parallelograms AG and DH, have the sides about the equal angles A and D, or else about the unequal angles (nevertheless given) have a given reason to one another, the reason *a* of the parallelogram AG to the parallelogram DH is given. *a 70. prop.* But the triangle ABC is the half of the parallelogram AG *b* and the triangle DEF *b 34. prop.* is the half of the parallelogram DH. Therefore the reason of the triangle ABC to the triangle DEF is given.

PROP. LXXII.




If of two triangles ABC and DEF , the bases BC and EF , are in a given reason, BC to EF , and that from the angles A and D , there be drawn to those bases the right lines AG and DH , making the equal angles AGC and DHF , or else unequal (yet nevertheless given) which shall have to one another given reasons AG to DH , those triangles ABC and DEF shall have also a given reason to one another, to wit, ABC to DEF .

Constr. For let the parallelogram KC and LF be finished.

Demonstr. Forasmuch as the angles AGC and DHF are equal, or unequal (yet given) and that the angle AGC *a* is equal to the angle KBC . But the angle DHF equal to the angle LEF , the angles at the points B and E are equal, or else unequal (yet given) and for that the reason of AG to DH is given, and AG is equal to KB . But DH is equal to LE , also the reason of KB to LE is given. But the reason of BC to EF is also given, and the angles at the points B and E are equal, or
a 29. I.
b 7c. prop. else unequal (yet given :) Therefore *b* the reason of the parallelogram KC to the parallelogram LF is given; and therefore the reason of the triangle ABC to the triangle DEF is given, seeing those triangles *c* are the one half of the parallelograms.

PROP.

PROP. LXXIII.



If of two parallelograms A-B and EG, the sides about the equal angles C and F, or else about the unequal angles (but nevertheless given) are in such sort to one another, that as the side CB of the first, is to the side FG of the second; so the other side EF of the second, is to some other right line CN. But that the other side AC hath also to the same right line CN a given reason, those parallelograms will have also to one another a given reason AB to EG.

Constr. For in the first place, let the parallelogram AB be equiangled to EG, and having placed CN directly to AC: Let the parallelogram CM be finished.

Demonstr. Forasmuch then as CB or NM its equal, is to FG, so is EF to CN, and that the angles N and F are equal (for N is equal to the angle ACB, which is put equal to F) the parallelograms CM and EG are equal: a 14. 6.

But as AC to CN, so b the parallelogram AB b 1. 6: is to the parallelogram CM or EG: Therefore seeing that the reason of AC to CN is given, the reason of AB to EG is also given.

Constr. 2. Now suppose the parallelogram AB not to be equiangled to the parallelogram EG, and let there be constituted at the given point C in the line CB, the angle BCK, equal to the angle EFG, and so finish the parallelogram CL.

F f 2

Demonstr.

Demonstr. 2. Seeing that each of the angles ACB and KCB is given, the remaining angle ACK is also given. But *c* the angle CAK is given, as also the remaining angle AKC: *c* *sch.* 69. *prop.* Therefore *d* the triangle ACK is given by kind; and therefore the reason of AC to CK is given. But the reason of the same AC to CN is also given: Therefore *e* the reason of CK to CN is given. And seeing that as CB is to FG; so is EF to the right line CN, to which the other side KC hath a given reason, and that the angle BCK is equal to the angle F, the reason of the parallelogram CL to the parallelogram EG is given (by the first part of this Proposition) but the parallelogram CL is equal to the parallelogram AB: Therefore the reason of the parallelogram AB to the parallelogram EG is given.

P R O P. LXXIV.

If two parallelograms (as in the former figure) AB and EG, in equal angles C and F, or else in unequal angles (yet nevertheless given angles) have a given reason to one another, as one side CB of the first shall be to one side FG of the second, so the other side EF of the second, shall be to that to the which the other side AC of the first hath a given reason.

Constr. For either AB is equiangled or not; suppose it in the first place to be equiangled, and to the right line BC let there be applied the parallelogram CM, equal to the parallelogram EG, and so posited, as that *a* *sch.* 68. *prop.* AC and CN may be direct: Therefore a DB and BM shall be also direct (that is as one right line.)


Demonstr. Seeing that the reason of AB to EG is given, and that CM is equal to EG, the reason of AB to CM is also given; and therefore the reason of AC to CN is given (seeing

(seeing AB is to CM , b as AC is to CN ;) and b 1. 6.
for that CM is equal and equiangled to EG ,
the sides about the equal angles of the paral-
lelograms CM and EG , c are reciprocally pro- c 14. 6.
portional; and therefore as CB is to FG , so
is EF to CN . But the reason of AC to CN
is given: Therefore as CB is to FG , so is
 EF to that to which AC hath a given reason.

Constr. 2. Now suppose AB not to be equi-
angled to EG , and in the given point C of
the line CB , let there be constituted the an-
gle BCK equal to the angle EFG , and finish
the parallelogram CL .

Demonstr. 2. Seeing then that the reason of
 AB to EG is given, and d that AB is equal d 36. 1.
to CL , also the reason of CL to EG is given,
and the angle BCK is equal to the angle F ,
and therefore CL e is equiangled to EG : e sch. 69.
Therefore (by the first part of this Proposi. *prop.*
tion) as CB is to FG , so is EF to that to
the which CK hath a given reason. But the
reason of AC to CK is given; (as appears
by what hath been demonstrated in the latter
part of the precedent Proposition.) Therefore
as CB is to FG , so is EF to that to which
 AC hath a given reason.

P R O P. LXXV.



If two tri-
angles ABC
and DEF , in
equal angles
 A and D , or
else unequal
(yet neverthe-
less given)
have to one

another a given reason, as the side AB of the first,
shall be to the side DE of the second, so the
F i 3 other

other side DF of the second, shall be to that right line to the which the other side AC of the first hath a given reason.

Constr. For let the parallelograms AG and DH be finished.

Demonstr. Forasmuch as the reason of the triangle ABC to the triangle DEF is given, also the reason of the parallelogram AG to the parallelogram DH is given.

Seeing therefore that the two parallelograms AG and DH in equal angles, or unequal angles (nevertheless given) have to one another a given reason; as a AB is to DE , so is DF to that to which AC hath a given reason.

PROP. LXXVI.




If from the top A of a triangle ABC , given by kind, there be drawn to the base BC , a perpendicular line AD , that line AD shall have to the base BC a given reason.

Demonstr. For seeing that the triangle ABC is given by kind, the reason of AB to BC is given; and the angle B is also given. But the angle ADB is given; therefore the other angle BAD is given. Wherefore a the triangle ABD is given by kind; and therefore the reason of AB to AD is given. But the reason of AB to BC is given: Therefore b the reason of AD to BC is given.

PROP.

PROP. LXXVII.

If two figures ABC and DEF , given by kind, have to another a given reason, the reason also shall be given of which you please of the sides of one of the figures, to which you please of the sides of the other figure.




Constr. For on the right lines BC and EF , let there be described the squares BG and EH .

Demonstr. Forasmuch as on one and the same right line BC , are described two figures ABC and BG ,

given by kind, *a* the reason of the said ABC ^{a 49. prop.} to BG is given : In like manner, the reason of DEF to EH is given ; and seeing that the reason of ABC to DEF is given, and also that of the same figure ABC to BG ; and again the reason of DEF to EH : *b* the reason of BG ^{b 8. prop.} to EH is given ; and therefore the reason of BC to EF is also given.

PROP. LXXVIII.



If a given figure ABC, hath a given reason to some rect-angled figure DF, and that one side BC hath a given reason to one side DE, the rectan-

gled figure DF is given by kind.

Constr. For on the right line BC let the square BH be described, and to the right line DE, let the parallelogram DK be applied equal to BH, in such manner, as that GD and DI may be placed directly, *a* and by consequence FE and EK also directly.

a sch. 68. prop.

Demonstr. Therefore seeing that on one and the same right line BC are described the two rectilineal figures ABC and BH, given by kind, *b* the reason of ABC to BH is given. But the reason of the said ABC to DF is also given: Therefore *c* the reason of BH to DF is given. But BH is equal to DK: Therefore the reason of DK to DF is also given. And seeing that BH is equal and equiangled to DK, both the one and the other being rect-angles, *d* the sides of those figures are recipro-

d 14. 6.

cally proportional; and as BC is to DE, so is DI to CH. But by supposition, the reason of BC to DE is given; therefore also the reason of DI to CH is given; but the reason of

of

of DI to
DG e a
son of D
to BC, t
the reaso
reason o
therefore
and the
fore g I

If tw
gle B e
angles I
and FH
the bas
perpend
triangle
triangle


Co
be de
line
let
the
and

of DI to DG is also given : (for DI is to DG *e* as DK to DF :) Therefore *f* the reason of DG to CH is given. But CH is equal *f* 8. *prop.* to BC, seeing that BH is a square ; therefore the reason of BC to DG is given. But the reason of the same BC to DE is also given ; therefore the reason of DE to DG is given, and the angle at D is a right angle ; Therefore *g* DF is given by kind.

g sch. 6r.
prop.

PROP. LXXIX.

If two triangles ABC and EFG, have an angle B equal to an angle F. But from the equal angles B and F there be drawn perpendiculars BD and FH, to the bases AC and EG ; and that as the base AC of the first triangle ABC, is to the perpendicular BD, so also the base EG of the other triangle EFG, is to the perpendicular FH, those triangles ABC and EFG are equiangled.



Constr. For about the triangle EFG let there be described the circle FFLG, then on the right line EG, and in the point E given therein, let there be made the angle GEL, equal to the angle C, and let FL and LG be drawn, and the perpendicular LM.

Demonstr.

a 21. 3.

b 7. 5.

c 28. 1.

d 33. 1.

e 29. 1.

f 21. 3.

Demonstr. Seeing then that the angle GEL is equal to the angle C, and the angle ELG is equal to the angle EFG, *a* they being in one and the same segment of the circle; the third angle EGL is equal to the third angle A. Wherefore the triangle ABC is alike to the triangle ELG, and the perpendiculars BD and LM are drawn: Therefore † as AC is to BD, so is EG to LM; but by supposition as AC is to BD; so is EG to FH: Therefore *b* LM is equal to FH. But the said LM is *c* parallel to FH: Therefore *d* FL is also parallel to EG; and therefore the angle FLE *e* is equal to the angle LEG. But the angle C is also equal to the said angle LEG, and the angle FLE to the angle FGE *f*: Therefore also the angle C is equal to the angle FGE. But by supposition the angle ABC is equal to the angle EFG: Therefore the third angle BAC is equal to the third angle FEG: Wherefore the triangle ABC is equiangled to the triangle EFG.

Scholium.

g 4. 6.

h 4. 6.

† Now that as AC is to BD, so EG is to LM, it is by some thus demonstrated. Forasmuch as the angle C is equal to the angle GEL, and the angle BDC to the angle LME, each being a right angle, the other angle CBD is equal to the other angle ELM: Therefore *g* as EM is to ML, so is CD to DB. Again, seeing the angle ABC is equal to the angle ELG, and the angle CBD to the angle ELM, the remaining angle ABD is equal to the remaining angle MLG; but the angle ADB is also equal to the angle LMG; and therefore the third angle A is equal to the third angle LGM: Therefore *h* as AD is to DB, so is GM to ML. But it hath been demonstrated that as

CD

GD is to
AC is to

If a t
that the
AC, com
reason
triangle
Const
there
BE.

Dem
given,
ABE
reason
reason
rectan

also
AB
The
BC
and
to t
rect
let
ma
be

GD is to DB, so is EM to ML : Therefore *i* as *i* 14. 5.
AC is to BD, so is EG to LM.

PROP. LXXX.

If a triangle *ABC* hath one angle *A* given, and that the rectangle contained under the sides *AB* and *AC*, comprising the given angle *A*, hath a given reason to the square of the other side *BC*, the triangle *ABC* is given by kind.

Constr. For from the points *A* and *B*, let there be drawn the perpendiculars *AD* and *BE*.

Demonstr. Forasmuch as the angle *BAE* is given, and also the angle *AEB*, the triangle *ABE* is given by a kind; and therefore the *a* 40. *prop.* reason of *AB* to *BE* is given: Therefore the reason of the rectangle of *AB* and *AC* to the rectangle of *BE* and *AC* is also given (for

it is the same reason *b* 1. 6.

b as of *AB* to *BE*.) But

the rectangle of *AC* and

BE is equal to the rect-

angle of *BC* and *AD*;

for that each of those rect-

angles is *c* double to the

triangle *ABC*. Therefore

the reason of the rectangle

of *AB* and *AC* to the rect-

angle of *BC* and *AD* is

also given. But the reason of the rectangle of

AB and *AC* to the square of *BC* is given:

Therefore *d* also the reason of the rectangle of *d* 8. *prop.*

BC and *AD* to the square of *BC* is given;

and therefore the reason of the right line *EC*

to the right line *AD* is given. (For that *e* the *e* 1. 6.


rectangle is to the square as *AD* to *BC*.) Now

let the right line *FG* given by position and

magnitude, be exposed; and thereon let there


be described the segment of a circle *FIG*;

capable



f 8. def.

g 4. def.



h 2. prop.

i 27. prop.

k 28. prop.

l 25. prop.

m 26. prop.

n 39. prop.

o 34. prop.

p 79. prop.

capable of an angle equal to the angle A. And seeing the said angle A is given, also the angle in the segment FLG shall be given; and therefore f the same segment is given by position. From the point G let there be erected at right angles on the line FG, the line

GH, which g is given by position: Let it be so made, that as BC is to AD, so FG may be to GH; and seeing that the reason of BC to AD is given, also that of FG to GH is given. But FG is given: Therefore h GH is given by

magnitude. But it is also given by position, and the point G is given: Therefore the point

i H is i also given. Now by the point H let there be drawn HI, parallel to FG, and that

k line HI shall be given by k position. But the segment of the circle FIG is also given by

l position. Therefore l the point I is given. Let the right lines IF and IG be drawn, and the perpendicular IK: Therefore IK is given

by position. But the point I is given, as also

m each of the points F and G: Therefore m each of the lines FG, FI, and IG is given

n by position and magnitude: Wherefore n the triangle FIG is given by kind; and seeing

that as BC is to AE, so is FG to GH, and

o that to GH, IK is equal, as BC is to AE, so is FG to IK, and the angle A is equal to

p the angle FIG: Therefore p the triangle ABC is equiangular to the triangle FIG. But FIG

is given by kind: Therefore also the triangle ABC is given by kind.

OTHER-

Const.
is given
ained u
be given
ven by
Demon
that Ip
compou
of BC,
ABC.
the reaf
But the
rectang




to the
s the r
BC is
the re
BC ro
fore r
poun
ven;
BC i
poun
of th
is g

OTHERWISE.

Constr. Let the triangle ABC, whose angle A is given, and the reason of the rectangle contained under AB and AC, to the square of BC be given: I say that the triangle ABC is given by kind.

Demonstr. For seeing the angle A is given, that space of which the square of the line compounded of BAC is greater than the square of BC, *q* hath a given reason to the triangle *q* 67. *prop.* ABC. Now let that space be D: Therefore the reason of D to the triangle ABC is given. But the reason of the triangle ABC to the rectangle of AB and AC is given; *r* seeing *r* 66. *prop.*


the angle A is given: Therefore the reason of the space D to the rectangle of AB and AC is given. But the reason of the rectangle of AB and AC



to the square of BC is also given: Therefore the reason of the space D to the square of BC is given. Wherefore by compounding, the reason of the space D, with the square of BC to the said square of BC is given: Therefore the reason of the square of the line compounded of BAC, to the square of BC is given; (for that the space D with the square of BC is equal to the square of the line compounded of BAC;) and therefore *u* the reason of the said line compounded of BAC to BC is given. But the angle A is also given: There-

x 46. *prop.* Therefore x the triangle ABC is given by kind.

PROP. LXXXI.



If of three right lines A, B, and C, proportional to three other proportional right lines D, E, and F, the extremes A and D, C and F, are in a given reason (so

wit, as A to D, and C to F,) also the means B and E shall be in a given reason, and if one extreme hath a given reason to an extreme, and the mean to the mean, the other will have also a given reason to the other.

Demonstr. Forasmuch as the reason of A to D, and of C to F is given, the rectangle of A and D a shall have a given reason to the rectangle of C and F. But the rectangle of A and D is equal b to the square of B; and the rectangle of C and F to the square of E. Therefore the reason of the square of B to the square of E is given; and therefore c the reason of the line B to the line E is also given.

Again, Let the reason of A to D, and B to E, be given: I say that the reason of C to F is also given. For seeing that the reason of A to D, and of B to E is given, also the reason d 50. *prop.* of the square of B d to the square of E is given. But the square of B is equal to the rectangle of A and C, and the square of E to the rectangle of D and F: Therefore the reason of the rectangle of A and C to the rectangle of D and F is given. But the reason of a side A to a side D is given: Therefore e the reason of the other side C to the other side F is also given.

PROP.

A —
B —
C —
D —
E —
F —

which I
as that
Demon
therefor
And se
And ag
son of
But E i
reason,
ven rea
B hath
D hat

A ha
De
E, th
equa

PROP. LXXXII.

A ——— If there be four right lines A, B, C,
 B ——— and D, proportional, as the first A,
 C ——— shall be to that line to which the se-
 D ——— cond B hath a given reason, so the
 E ——— third C, shall be to that to which
 F ——— the fourth D hath a given reason.

Constr. Let E be the line to which B hath a given reason, and let it be so as that B may be to E, as D is to F.

Demonstr. Now the reason of B to E is given, therefore also the reason of D to F is given. And seeing that as A is to B, so is C to D. And again, as B is to E, so is D to F, by reason of equality as A is to E, so is C to F. But E is that line to which B hath a given reason, and F that to which D also hath a given reason: Therefore as A is to that to which B hath a given reason, so C is to that to which D hath a given reason.

PROP. LXXXIII.


A ——— If four right lines A, B, C,
 B ——— and D, are in such sort to one
 C ——— another, that of any three of them
 D ——— A, B, and C, and a fourth E,
 E ——— taken proportional, to which that
 ——— line D, which remains of the four
 ——— lines, hath a given reason, the
 ——— four lines A, B, C, and E, are
 ——— proportional; as the fourth D is
 ——— to the third C, so the second B
 ——— shall be to that to which the first

A hath a given reason.

Demonstr. Forasmuch as A is to B as C is to E, the rectangle contained under A and E is ^{a 16. 6.} equal to the rectangle contained under B and C;

- C; and seeing that the reason of D to E is given, also shall be given the reason of the rectangle of A and D to the rectangle of A and E (for *b* it is the same reason as of D to E.) But the rectangle of A and E is equal to the rectangle of B and C: Therefore the reason of the rectangle of A and D to the rectangle of B and C is given. Wherefore *c* as D is to C, so is B to that to which A hath a given reason.

PROP. LXXXIV.




If two right lines AB and AE comprehending a given space AF in a given angle BAE, and that the one AB be greater than the other AE by a given line CB, also each of the lines AB and AE is given.

- Demonstr.* For seeing that AB is greater than AE by the given line CB, the remainder AC is equal to AE: Finish the parallelogram AD. Therefore seeing that AE is equal to AC, the reason of AE to AC is given, and the angle A is also given: Therefore *a* AD is given by kind. Wherefore the given space AF is applied to the given right line CB, exceeding it by the given figure AD given by kind; and therefore *b* the breadth of the excess is given. Therefore AC is given. But CB is also given: Therefore the whole AB is given. But AE is also given: Therefore each of the right lines AB and AE is given.

PROP.

PROP. LXXXV.



If two right lines AC and CD, do comprehend a given space AD in a given angle ACD, the line compounded of those lines AC and CD is given, also each of those lines AC and CD is given.

Constr. For let AC be prolonged to the point B, and let CB be put equal to CD, then by the point B let BF be drawn parallel to CD, and so finish the parallelogram CF.


Demonstr. Seeing then that CB is equal to CD, and the angle DCB is given; for that angle that follows is the given angle; and therefore *a* the parallelogram DB is given by kind; and again, seeing that the line compounded of ACD is given, and CB is equal to CD, also AB is given. And thus to the right line AB there is applied the given space AD, deficient by the figure DB given by kind; and therefore *b* the breadths of the defects are also given: Therefore the right lines DC and CB are given. But the compounded line ACD is also given: Therefore *c* each of the lines AC and CD is given.

a sch. 61. prop.

b 58. prop.

c 4. prop.

PROP. LXXXVI.



If two right lines AB and BC , do comprehend a given space AC , in a given angle ABC , the square of the one BC , is greater than the square of the other AB , by a given space (yet in a given reason) also each of those

lines AB and BC shall be given.

- Demonstr.* For seeing that the square of BC is greater than the square of AB by a given space (yet in a certain reason.) Let the given space be taken away, that is to say, the rectangle contained under CB and BE : Therefore
- a 11. def. a the reason of the remainder, b which is the
 - b 2. 2. rectangle contained under BC and CE to the square of AB is given. And forasmuch as the rectangle $\dagger$ under AB and BC is given, and
 - c 1. prop. also that of CB and BE , their c reason is given.
 - d 1. 6. But as the rectangle under AB and BC is to the rectangle under CB and EB , d so AB is to BE ; and therefore the reason of AB to BE
 - e 50. prop. is given: Wherefore e the reason of the square of AB to the square of BE is also given. But the reason of the square of AB to the rectangle under BC and CE is given: Therefore f
 - f 8. prop. also the reason of the rectangle under BC and CE to the square of BE is given. Wherefore the reason of four times the rectangle under BC and CE to the square of BE is given; and by compounding, g the reason of four
 - g 8. prop. times the rectangle under BC and CE , with the square of BE to the square of BE is given. But four times the rectangle of BC and CE , with the square of BE , h is the square of the
 - h 8. 2. compound line BCE : Therefore the reason of the

the


square of the compound line BCE to the square of BE is given : Wherefore *i* the reason of the *i* 54. prop. line compounded of BC and CE to BE is given, and by compounding, *k* the reason of the *k* 6. prop. compound of the lines BC, CE, and BE, that is to say, the double of BC to BE is given ; and therefore the reason of the only line BC to BE is also given. But as BC is to BE, / so *l* 1. 6. the rectangle under BC and BE is to the square of BE : Therefore the reason of the rectangle under BC and BE to the square of BE is given. But the rectangle of BC and BE is given : Therefore *m* the square of BE is also gi- *m* 2. prop. ven, and consequently the line BE is given. Wherefore BC is also given, seeing that the reason of BE to BC is given. But the space AC is given, and also the angle B : Therefore *n* AB is given. Wherefore each of the lines *n* 57. prop. AB and BC is given.

Scholium.

† Instead of saying in this place [what is under, &c.] we have used this Word Rectangle, it being manifest by what follows that such was the intention of EUCLIDE, seeing he makes use in the said Demonstration of the second and eighth Proposition of the twelfth Element ; and also that the space or Parallelogram given being not rectangled, it may be reduced thereto, making on BC, and in the given point B, a right angle CBA, so as that there will be two Parallelograms constituted on one and the same base BC, and between the same parallels, as in the 69th Proposition by means whereof this Conclusion is drawn.

Note, This serves also for the next Prop.

PROP. LXXXVII.



If two right lines AB and BC , do comprehend a given space AC , in a given angle B , the square of the one BC , is greater than the square of the other AB , by a given space,

also each of those lines AB and BC shall be given.

- Demonstr.* For seeing that the square of BC is greater than the square of AB by a given space: Let the given space be taken away, and let the rectangle be contained under BC and BE : Therefore the remainder, *a* which is the rectangle of BC and CE , is equal to the square of AB . And seeing that the rectangle of BC and BE is given, and also the space or rectangle AC ; the reason of the said rectangle of BC and BE to AC is given. But as *b* the rectangle of BC and BE is to the rectangle of AB and EC , so is BE to AB : Therefore the reason of BE to AB is given, and therefore *c* the reason of the square of the said DE to the square of AB is also given. But to that square of AB the rectangle of BC and CE is equal: Therefore the reason of the said rectangle of BC and CE to the square of BE is given; and therefore the reason of the quadruple of the said rectangle of BC and CE to the square of BE is also given; and by compounding, *d* the reason of four times the rectangle of BC and CE , with the square of BE , to the said square of BE is given. But four times the rectangle of BC and CE , with the square of BE , *e* is the square of the compound line BCE : Therefore the reason of the square of that compound line BCE to the square of BE is


is also
compre
fore
compre
twice
reason
But
given
given
BE is
so th
and
rectan
the
being
was
give
But
there

A

C
and
and
giv
AE
ang

is also given; and therefore the reason *f* of the *f* 54. *prop.*
compound line BCE to BE is given. Where-
fore by compounding, *g* the reason of the said *g* 6. *prop.*
compound line BCE and EB, that is to say,
twice BC to BE is also given; therefore the
reason of the only line BC to BE is given.
But the reason of the same BE to AB is also
given: Therefore *h* the reason of AB to BC is *h* 8. *prop.*
given. And seeing that the reason of BC to
BE is given, and that as the said BC is to BE,
so the square of BC to the rectangle of BC *i* 1. 6.
and BE, the reason of the square of BC to the
rectangle of BC and BE is also given. But
the said rectangle of BC and BE is given, it
being that which was taken away, and which *k* 2. *prop.*
was given. Therefore the square of BC *k* is
given, and therefore the line BC is given.
But the reason of the same BC to BA is given,
therefore AB is also given.

P R O P. LXXXVIII.



If in a circle
ABC, given by mag-
nitude, there be
drawn a right line
AC, which shall take
away a segment
ABC, which doth
comprehend a given
angle AEC, that
line AC is given by
magnitude.

Constr. For let D be the center of the circle;
and let the diameter thereof ADE be drawn,
and let EC be joined.

Demonstr. Forasmuch as the angle ACE is
given, for *a* it is a right angle. But the angle *a* 31. 3.
AEC is also given, and therefore the other
angle CAE is given. Wherefore the triangle
G g 3 ACE

- b 40. prop. $\angle ACE$ is given by kind; and therefore the reason of EA to AC is given. But AE is given by magnitude, seeing that the circle ABC is given by magnitude. Therefore AC is also given by magnitude.

PROP. LXXXIX.

If in a circle ABC , given by magnitude, there be drawn a right line AC , given by magnitude, that line AC will take away a segment ABC , comprehending a given angle.

Constr. For having taken the point D for the center of the circle, let the diameter ADE be drawn, as also the right line EC .


Demonstr. Forasmuch as each of the right lines AE and AC are given, the reason of the line AE to AC is given; and the angle ACE is a right angle: Therefore the triangle ACE is given by kind, and therefore the angle AEC is given.

PROP. XC.

If in the circumference of a circle ABC given by position, and by magnitude, there be taken a given point B , and that from that point B , to the circumference of the circle ABC , there doth bend a right line BAC , making a given angle BAC the other extremity C of the bent line shall be given.

Constr. For let the center of the circle be D , and let the right lines BD and BC be drawn.


Demonstr.



Demonstr. Forasmuch as each point B and D is given, the right line BD, *a* 26. *prop.* is given by position; and seeing that the angle BAC is given, the angle BDC is also given. Wherefore to the right line BD given by position, and in the point D given therein, there is drawn the right line CD; which

makes the given angle BDC; and therefore *b* 29. *prop.* the line DC is given by position. But the circle ABC is given by position and magnitude: Therefore *c* the right line DC is given by position and by magnitude. But the point D is given: Therefore *d* the point C is also given. *d* 27. *prop.*

PROP. XCI.



If from a given point C, there be drawn a right line CA, which shall touch a circle AB, given by position; that line CA is given by position and by magnitude.

Constr. For having taken the point D

for the center of the circle, let the right lines DA and DC be drawn.


Demonstr. Forasmuch as each point C and D is given, the right line CD *a* is given by position and by magnitude. But the angle CAD *b* is a right angle; and therefore the semicircle *b* 18. 3. described on CD shall pass by the point A: Let it then pass by that point, and let the semicircle be DAC: Forasmuch as the same

G g 4

DAC

c 6. def. DAC c is given by position, and also the circle
 d 25. prop. ABE, d the point A is given. But the point
 e 26. prop. C is also given: Therefore e the right line
 AC is given by position and by magnitude.

P R O P. XCII.



If without a circle ABC, given by position, there be taken some point D, and from that given point there be drawn a right line DB, cutting the circle, the rect-
 angle comprised un-


der the whole line BD, and the part DC, between the point D, and the circumference convex AC shall be given.

Constr. For from the point D let the right line DA be drawn, which shall touch the circle in the point A.

a 91. prop. Demonstr. Therefore DA a is given by position and magnitude; and therefore the square
 b 52. prop. of the said DA is b given. But the said
 c 36. 3. square of DA is equal c to the rectangle of BD and DC: Therefore the said rectangle of BD and DC is also given.

OTHER-

OTHERWISE.




Constr. Let E be the center of the circle, and by the same center let there be drawn from the point D the right line DA.

Demonstr. Forasmuch as each point D and E

is given, the right line DE is *d* given by position and by magnitude. But the circle ABC is given by position and by magnitude: Therefore each point A and F *e* is given, and the point D is also given; and therefore *d* each *e* 25. *prop.* line AD and FD is given. Wherefore the rectangle of the lines AD and DF is also given. But the said rectangle of AD and DF is equal to the rectangle of DB and DC: Therefore the rectangle of DB and DC is given.

PROP. XCIII.




If in a circle given by position there be taken a given point A, and by that point A there be drawn a right line BC to the circle, the rectangle comprised under the segments of the same line BC shall be given.

Constr. For let D be taken for the center of the circle, and having

having drawn the right line AD prolong it to the points E and F.

Demonstr. Forasmuch as each point A and D is given, the right line AD *a* is given by position. But the circle BEC is also given by position: Therefore each point E and F is also given by position, and the point A is given. Wherefore each line *b* AE and AF is given: Therefore the rectangle of the same lines AE and AF is given; and is equal to the rectangle *b* of AB and AC: Therefore the said rectangle of AB and AC is given.

P R O P. XCIV.



If in a circle ABC, given by magnitude, there be drawn a right line BC, which doth take away a segment which doth comprehend a given angle ABC, and that the said angle being in the segment is cut into two equal parts, the line compounded of the right lines BA and AC, which comprehend the given angle BAC shall have a given reason to the line AD, which doth divide that angle into two equal parts; and the rectangle contained under the line compounded of those lines BA and AC, comprehending the given angle BAC, and that part ED of the intersecting line which is below the segment between the base BC and the circumference, shall be given.

Constr. Let BD be drawn.


Demonstr. Forasmuch as in the circle ABC given by magnitude, there is drawn the right line BC, which takes away the segment BAC comprehending the given angle BAC, that

line BC
given:
is cut i
BA
pound
and by
CA to
is equ
ACE
AEC
theref
to the
CE, f
so th
BC:
to E
as th
BC t
given
line
I say
line
the
gle
ang
BE
as
line
line
for
and
BE
gi
gi
po

line BC *a* is given; and therefore BD is also a *88. prop.*
 given: Therefore the reason of BC to BD *b* is *b 1. prop.*
 given. And seeing that the given angle BAC
 is cut in two equal parts by the right line AD, as
 BA is to CA, so is BE to CE; and by com- *c 3. 6.*
 pounding, as BAC is to CA, so is BC to CE;
 and by permutation, as BAC is to BC, so is
 CA to CE. And seeing that the angle BAE
 is equal to the angle CAE, and the angle
 ACE *d* to the angle BDE, the other angled *21. 3.*
 AEC is equal to the other angle ABD; and
 therefore the triangle ACE is equiangled
 to the triangle ABD: Therefore *e* as AC is to *e 4. 6.*
 CE, so is AD to BD. But as AC is to CE,
 so the line compounded of BA and AC is to
 BC: Therefore as the compound line BAC is
 to BC, so is AD to BD; and by permutation,
 as the compound line BAC is to AD, so is
 BC to BD. But the reason of BC to BD is
 given: Therefore the reason of the compound
 line BAC to AD is also given. Moreover,
 I say that the rectangle under the compound
 line BAC and ED is given. For seeing that
 the triangle AEC is equiangled to the trian-
 gle BDE, (for the angle ACE *d* is equal to the
 angle BDE, and the angle AEC *f* to the angle *f 15. 1.*
 BED) as BD is to DE, so is AC to CE. But
 as AC is to CE, so is also the compound
 line BAC to BC: Therefore as the compound
 line BAC is to BC, so is BD to DE. Where-
 fore the rectangle of the compound line BAC
 and DE *g* is equal to the rectangle of BC and *g 16. 6.*
 BD. But the rectangle of BC and BD is
 given, (for that those lines BC and BD are
 given:) Therefore the rectangle under the com-
 pound line BAC and ED is also given.

OTHER-

OTHERWISE.



Constr. Let CA be prolonged to the point E, and let AE be put equal to BA, and let BE and BD be joined.

Demonstr. Forasmuch as the angle BAC is double to each of the angles CAD and AEB (for the angle BAC is cut into two equal parts by the line AD, and equal b

h 32.1.

i 5.1.

k 21.3.


to the two angles ABE and AEB, which i are equal) the angle ABE is equal to the angle CAD, that is to say, k to the angle CBD; adding therefore the common angle ABC, the whole angle ABD shall be equal to the whole angle FBE. But the angle ACB is k equal to the angle ADB: Therefore the third angle AEB is equal to the third angle BAD; and therefore the triangle CEB is equiangular to the triangle ABD: Wherefore as CE is to CB, so is AD to BD. But the right line CE is compounded of the two lines CA and AB: Therefore as the compound line BAC is to CB, so is AD to BD; and by permutation, as the compound line BAC is to AD, so is CB to BD. But the reason of CB to BD is given, seeing that each of those lines is given: Therefore the reason of the compound line BAC to AD is also given. And seeing that the

the trian
FBD (fo
gle BFD
ADB) a
EC is
Therefo
CB, so
angle o
equal to
the rect
dering
given:
pound

C
CF
line
I
and
are
to
is

the triangle CEB is equiangled to the triangle
 FBD (for the angle AFC is equal l to the an- l 21. 3.
 gle BFD, and the angle ECB m to the angle m 16. 6.
 ADB) as EC is to CB, so is BD to DF. But
 EC is equal to the compound line BAC:
 Therefore as the compound line BAC is to
 CB, so is BD to DF. Wherefore n the rect- n 16. 6.
 angle of the compound line BAC and DF is
 equal to the rectangle of CB and BD. But
 the rectangle of CB and BD is given, consi-
 dering that each of the lines CB and BD is
 given: Therefore the rectangle of the com-
 pound line BAC and DF is given.

OTHERWISE.



Constr. Let AC be prolonged to F, and let
 CF be put equal to AB, and let the right
 lines BD and DF be drawn.


Demonstr. Forasmuch as BA is equal to CF,
 and o BD to DC, the two sides AB and BD o 26, 29,
 are equal to the two sides CD and DF, each 3 .
 to his corresponding side, and the angle ABD
 is equal to the angle DCF, p seeing that the p 22. 3.

four

q 4. r.

four sided figure ABDC is within the circle: Therefore the base AD is *q* equal to the base DF, and the angle DAB to the angle DFC. But the angle BAD is given, (being the half of the given angle BAC) Therefore the angle DFC is so also. But DAF is also given: Therefore the triangle ADF is given by kind. Wherefore the reason of FA to AD is given. But AF is the compound of BA and AC, for that CF is equal to AB: Therefore the reason of the compound line BAC to AD is given: The same Demonstration will serve to shew that the rectangle contained under the compound line BAC and ED is given also.

P R O P. XCV.



If in the diameter BC of a circle ABC given by position, there be taken a given point D, and that from that point D there be drawn a right line DA, to the circumference

of the circle. But from the section of the said line there be drawn a right line AE, perpendicular thereto, and by the point E where that perpendicular doth meet with the circumference, there be drawn a parallel EF, to the first line drawn AD, that point F in which the parallel meets with the diameter, is given; and the rectangle contained under the parallel lines AD and EF is also given.

Constr.

Constr. Let the right line EF be prolonged to the point G, and let the right line AG be drawn.

Demonstr. Forasmuch as the angle AEG is a right angle, the right line AG is the diameter of the circle. But BC is also the diameter: Therefore the point H is the center of the circle. Now the point D is given; and therefore *a* the line DH is given by mag. *a* 26. *prop.* nitude. But seeing that AD is parallel to EG, and AH equal to GH; *b* DH is equal to FH, *b* 26. *prop.* and AD to FG; (for the angles AHD and FHG *c* are equal, and DAH and FGH *d* are *c* 15. 1. also equal.) But the line DH is given: *d* 29. 1. Therefore FH is also given. But each of those lines DH and HF is also given by position, and the point H is given: Therefore *e* the point F is also given. And seeing that in the circle ABC given by position, is taken the given point F, and through the same is drawn the right line EFG; the rectangle under EF and FG *f* is given. But FG is equal *f* 93. *prop.* to AD. Therefore the rectangle comprehended under AD and EF is given. Which was to be demonstrated.

The End of EUCLIDE'S DATA.

Let the right line AB be produced
to the point G , and let the right line AD be
drawn. I say, that the angle BAC is
greater than the angle BCD . For, the angle
 BAC is the sum of the angles BAD and
 DAC . But the angle BCD is less than
the angle BAD . Therefore the angle BAC is
greater than the angle BCD . Q.E.D.

Let the right line AB be produced
to the point G , and let the right line AD be
drawn. I say, that the angle BAC is
greater than the angle BCD . For, the angle
 BAC is the sum of the angles BAD and
 DAC . But the angle BCD is less than
the angle BAD . Therefore the angle BAC is
greater than the angle BCD . Q.E.D.

Let the right line AB be produced
to the point G , and let the right line AD be
drawn. I say, that the angle BAC is
greater than the angle BCD . For, the angle
 BAC is the sum of the angles BAD and
 DAC . But the angle BCD is less than
the angle BAD . Therefore the angle BAC is
greater than the angle BCD . Q.E.D.

Let the right line AB be produced
to the point G , and let the right line AD be
drawn. I say, that the angle BAC is
greater than the angle BCD . For, the angle
 BAC is the sum of the angles BAD and
 DAC . But the angle BCD is less than
the angle BAD . Therefore the angle BAC is
greater than the angle BCD . Q.E.D.

Let the right line AB be produced
to the point G , and let the right line AD be
drawn. I say, that the angle BAC is
greater than the angle BCD . For, the angle
 BAC is the sum of the angles BAD and
 DAC . But the angle BCD is less than
the angle BAD . Therefore the angle BAC is
greater than the angle BCD . Q.E.D.



BR

R

F

base
ther
mix



A

BRIEF TREATISE,

Added by FLUSSAS,

O F

Regular Solids.

Regular Solids are said to be composed and mix'd when each of them is transformed into other Solids, keeping still the form, number and inclination of the bases, which they before had to one another; some of which yet are transformed into mix'd Solids, and other some into simple.

H h

Into



Into mixt, as a Dodecaedron and an Icosaedron, which are transformed or altered, if you divide their sides into two equal parts, and take away the solid angles subtended of plain superficial figures, made by the lines coupling those middle sections; for the Solid remaining after the taking away of those solid angles, is called an Icosidodecaedron. If you divide the sides of a Cube and of an Octoedron into two equal parts, and couple the sections, the solid angles subtended of the plain superficies made by the coupling lines, being taken away, there shall be left a solid, which is called an Exoctoedron. So that both of a Dodecaedron and also of an Icosaedron, the Solid which is made shall be called an Icosidodecaedron; and likewise the Solid made of a Cube, and also of an Octoedron, shall be called an Exoctoedron. But the other Solid, to wit, a Pyramis or Tetraedron, is transformed into a simple Solid; for if you divide into two equal parts each of the sides of the pyramis, triangles described of the lines which couple the sections, and subtending and taking away the solid angles of the Pyramis, are equal and like unto the equilateral triangles left in each of the bases, of all which triangles is produced an Octoedron, to wit, a simple, and not a composed Solid. For the Octoedron hath four bases, like in number, form, and mutual inclination with the bases of the pyramis, and hath the other four bases with like situation opposite and parallel to the former. Wherefore the application of the pyramis taken twice, maketh a simple Octoedron, as the other Solids make a mix'd compound Solid.

DEFL.

I. An
tain
equ

DEFINITIONS.

- I. *An Exoctaedron is a solid figure contained of six equal squares, and eight equilateral and equal triangles.*



II. An Icosidodecaedron is a solid figure contained under twelve equilateral, equal, and equiangled Pentagons, and twenty equal and equilateral triangles.

For the better Understanding of the two former Definitions, and also of the two Propositions following, I have here set two figures, whose figures if you first describe upon pasted Paper or such like matter, and then cut them and fold them accordingly, they will represent unto you the perfect forms of an Exoctoedron, and of an Icosidodecaedron.

PROBLEME I.

To describe an equilateral and equiangled Exoctoedron, and to contain it in a given Sphere, and to prove that the Diameter of the Sphere is double to the side of the said Exoctoedron.

Constr. Suppose a Sphere whose diameter let be AB, and about the diameter AB let there be described a square *a*, and upon the square let there be described a Cube *b*, which let be CDEFQTVR; and let the diameter thereof be QR, and the center S. Divide the sides of the Cube into two equal parts in the points G, H, I, K, L, M, N, O, P, &c. and couple the middle sections by the right lines IN, NO, OP, PI, and such like, which sub-tend the angles of the squares or bases of the Cube; and they are equal *c*, and contain right angles, as the angle NIP. For the angle NID, which is at the base of the Isosceles triangle

a 6. 4.

b 15. 13.


c 4. 1.

R
triangle
to likew



A
fore t
the r
by th
KGH
Cube
ber,
of th
angle
the C
solid
the
trian
ing
mits
the

triangle NDI , is the half of a right angle, and so likewise is the opposite angle RIP . Where-




fore the residue NIP is a right angle, and so the rest. Wherefore $NIPO$ is a square. And by the same reason shall the rest $NMLK$, $KGHI$, &c. inscribed in the bases of the Cube, be squares, and they shall be six in number, according to the number of the bases of the Cube. Again, forasmuch as the triangle KIN subtendeth the solid angle D of the Cube, and likewise the triangle KGL the solid angle C , and so the rest which subtend the right solid angles of the Cube, and these triangles are equal and equilateral (to wit) being made of equal sides, and they are the limits or borders of the squares, and the squares the limits or borders of them; as hath been

.H h 3 before

before proved. Wherefore LMNOPHGK is an Exoctaedron by the definition, and is equilateral; for it is contained of equal subtendant lines, it is also equiangled; for every solid angle thereof is contained under two superficial angles of two squares, and two superficial angles of two equilateral triangles.

Demonstr. Forasmuch as the opposite sides and diameters of the bases of the Cube are parallels, the plain extended by the right lines QT and VR, shall be a parallelogram. And for that also in that plain lyeth QR, the diameter of the Cube, and in the same plain also is the line MH, which divideth the said plain into two equal parts, and also coupleth the opposite angles of the Exoctaedron; this



line MH therefore divideth the diameter into
two

two equal parts d ; and also divideth it self ^{d cor. 34. r.} in the same point, which let be S , into two equal parts e . And by the same reason may ^{e 4. r.} we prove that the rest of the lines which couple the opposite angles of the Exoctaedron, do in S the center of the Cube, divide one another into two equal parts, for each of the angles of the Exoctaedron are set in each of the bases of the Cube. Wherefore making the center the point S , with the distance SH or SM describe a Sphere, and it shall touch every one of the angles equidistant from the point S .

And forasmuch as AB the diameter of the sphere given, is put equal to the diameter of the base of the cube, to wit, to the line RT , and the same line RT is equal to the line MH ^f, which line MH coupling the opposite ^{f 33. r.} angles of the Exoctaedron, is drawn by the center. Wherefore it is the diameter of the Sphere given which containeth the Exoctaedron.

Lastly, forasmuch as in the triangle RFT , the line PO doth cut the sides into two equal parts, it shall cut them proportionally with the bases, to wit, as FR is to FP , so shall RT be to PO ^g. But FR is double to FP by supposition: Wherefore RT , or the diameter HM , is also double to the line PO , the side of the Exoctaedron. Wherefore we have described, &c. Which was required to be done.

PROBLEME II.

*To describe an equilateral and equian-
gled Icosidodecaedron, and to comprehend
it in a sphere given, and to prove that the*

$H h$ 4 diame-

diameter being divided by an extreme and mean proportion, maketh the greater segment double to the side of the Icosidodecaedron.

- Constr.* Suppose that the diameter of the sphere given be NL, *a* divide the line NL, by an extreme and mean proportion in the point I, and the greater segment thereof let be NI; and upon the line NI describe a Cube *b*; and about this Cube let there be circumscribed a Dodecaedron *c*; and let the same be ABCDEFHKMO, and divide each of the sides into two equal parts in the points Q, R, S, T, V, X, Y, Z, P, ϵ , δ , G, &c. and couple the sections with right lines, which shall subtend the angles of the Pentagons, as the lines PG, GV, VQ, QY, YR, RQ, VT, TX, XV, and so the rest.

Demonstr. Forasmuch as these lines subtend equal angles of the Pentagons, and those equal angles are contained of equal sides, to wit, of the halves of the sides of the Pentagons; therefore those subtending lines are equal *d*. Wherefore the triangles GQV, YQR, and VXT, and the rest, which take away solid angles of the Dodecaedron, are equilateral.

Again, forasmuch as in every Pentagon are described five equal right lines, coupling the middle sections of the sides, as are the lines QV, VT, TS, SR, and RQ, they describe a Pentagon in the plain of the Pentagon of the Dodecaedron. And the said Pentagon is contained in a circle, to wit, whose center is the center of a Pentagon of the Dodecaedron. For the lines drawn from that center to the angles of this Pentagon are equal, for that they are perpendiculars upon the bases cut.


R
cut e. V
equiangle



the re
bases
like.

Wh
numb
triang
lid an
laid
When
edron
edron
the
Penta
each

cut e. Wherefore the Pentagon QRSTV, is e 12. 4. equiangled f. And by the same reason may f 11. 4.



the rest of the Pentagons described in the bases of the Dodecaedron, be proved equal and like.

Wherefore those Pentagons are twelve in number: And forasmuch as the equal and like triangles do subtend and take away twenty solid angles of the Dodecaedron; therefore the said triangles shall be twenty in number. Wherefore we have described an Icosidodecaedron by the Definition, which Icosidodecaedron is equilateral; for that all the sides of the triangles are equal and common with the Pentagons; and it is also equiangled. For each of the solid angles is made of two superficial

perificial angles of an equilateral Pentagon, and of two superficial angles of an equilateral triangle.

Now let us prove that it is contained in the given sphere whose diameter is NL. Forasmuch as perpendiculars drawn from the centers of the Dodecaedron, to the middle sections of his sides, are the halves of the lines which couple the opposite middle sections of the sides of the Dodecaedron *g*; which lines also *h* do in the center divide one another into two equal parts. Therefore right lines drawn from that point to the angles of the Icosidodecaedron (which are set in those middle sections) are equal; which lines are thirty in number, according to the number of the sides of the Dodecaedron; for each of the angles of the Icosidodecaedron are set in the middle sections of each of the sides of the Dodecaedron. Wherefore making the center of the Dodecaedron, and the space any one of the lines drawn from the center to the middle sections, describe a sphere, and it shall pass by all the angles of the Icosidodecaedron, and shall contain it.


i 4. cor.
17. 13.

And forasmuch as the diameter of this solid, is that right line whose greater segment is the side of the Cube inscribed in the Dodecaedron *i*, which side is NI by supposition. Wherefore that solid is contained in the sphere given, whose diameter is put to be the line NL.

Now



N
of t
the
ang
ded
BE
AV
don
BE
lin
Cu
of
sep
fid
gi
be



Now let us prove that the greater segment of the diameter is double to QV the side of the solid. Forasmuch as the sides of the triangle AEB, are in the points Q and V divided into two equal parts, the lines QV and BE are parallels *k*. Wherefore as AE is to *k cor. 39.1.* AV, so is EB to VQ *l*. But the line AE is *l 2. 6.* double to the line AV. Wherefore the line BE is double to the line QV *m*. Now the *m 4. 6.* line BE is equal to NI, or to the side of the Cube *n*; which line NI is the greater segment *n 2 cor. of* of the diameter NL. Wherefore the greater *17. 13.* segment of the diameter given is double to the side of the Icosidodecaedron inscribed in the given sphere. Wherefore, We have described, &c. Which was required to be done.

ADVER-

A TREATISE of ADVERTISEMENT.

To the Understanding of the nature of this Icosidodecaedron, you must well conceive the passions and proprieties of both these solids, of whose bases it consisteth, to wit, of the Icosaedron and of the Dodecaedron. And altho' in it the bases are placed oppositely, yet have they to one another one and the same inclination. By reason whereof there lye hidden in it the actions and passions of the other Regular Solids. And I would have thought it not impertinent to the purpose to have set forth the inscriptions and circumscriptions of this Solid, if want of time had not hindred. But to the end the Reader may the better attain to the Understanding thereof, I have here following briefly set forth, how it may in or about every one of the five Regular Solids be inscribed or circumscribed; by the help whereof he may, with small travel or rather none at all, having well poised and considered the Demonstrations appertaining to the foresaid five Regular Solids, demonstrate both the inscription of the said Solids in it, and the inscription of it in the said Solids.

Of the Inscriptions and Circumscriptions of an Icosidodecaedron.

An Icosidodecaedron may contain the other five regular bodies. For it will receive the angles of a Dodecaedron in the centers of the triangles which subtend the solid angles of the Dodecaedron, which solid angles are twenty in number, and are placed in the same order in which the solid angles of the Dodecaedron taken away, or subtended by them, are. And by that reason it shall receive a Cube and a
Pyramis

R
Pyramis
as the a
of the o
An Ico
in the
of the I
Dodeca
And
twelve
centers
It m
bed in
in a P
ses con
after t
Icosae
be insc
bases
of the
in a
ceive
the b
scribe
comp
centr
solid
La
edron
of in
Dode
Con
T
solid
gles
as v
ten
Ico
thi
in
spr

Pyramis contained in the Dodecaedron, when as the angles of the one, are set in the angles of the other.

An Icosidodecaedron receiveth an Octoedron, in the angles cutting the six opposite sections of the Dodecaedron, even as if it were a simple Dodecaedron.

And it containeth an Icosaedron, placing the twelve angles of the Icosaedron in the same centers of the twelve Pentagons.

It may also by the same reason be inscribed in each of the five regular bodies, to wit, in a Pyramis, if you place four triangular bases concentrical with four bases of the Pyramis, after the same manner that you inscribed an Icosaedron in a Pyramis; so likewise may it be inscribed in an Octoedron, if you make eight bases thereof concentrical with the eight bases of the Octoedron. It shall also be inscribed in a Cube, if you place the angles which receive the Octoedron in it, in the centers of the bases of the Cube. Again, you shall inscribe it in an Icosaedron, when the triangles compassed in of the Pentagon bases, are concentrical with the triangles which make a solid angle of the Icosaedron.

Lastly, It shall be inscribed in a Dodecaedron, if you place each of the angles thereof in the middle sections of the sides of the Dodecaedron, according to the order of the Construction thereof.

The opposite plain superficies also of this solid are parallels. For the opposite solid angles are subtended of parallel plain superficies, as well in the angles of the Dodecaedron subtended by triangles, as in the angles of the Icosaedron subtended of Pentagons, which thing may easily be demonstrated. Moreover, in this solid are infinite properties and passions, springing of the solids whereof it is composed.

Where-

Wherefore it is manifest, that a Dodecaedron and an Icosaedron mixed, are transformed into one and the self same solid of an Icosidodecaedron. A Cube also and an Octoedron are mixed and altered into another solid, to wit, into one and the same Exoctoedron. But a Pyramis is transformed into a simple and perfect solid, to wit, into an Octoedron.

If we will frame these two solids joined together into one solid, this only must we observe.

In the Pentagon of a Dodecaedron inscribe a like Pentagon, and let its angles be set in the middle sections of the Pentagon circumscribed, and then upon the said Pentagon inscribed, let there be set a solid angle of an Icosaedron, and so observe the same order in each of the bases of the Dodecaedron, and the solid angles of the Icosaedron set upon these Pentagons shall produce a solid consisting of the whole Dodecaedron, and whole Icosaedron. In like sort, if in every base of the Icosaedron, the sides being divided into two equal parts, be inscribed an equilateral triangle, and upon each of those equilateral triangles be set a solid angle of a Dodecaedron, there shall be produced the same solid consisting of the whole Icosaedron, and of the whole Dodecaedron.

And after the same order, if in the bases of a Cube be inscribed squares subtending the solid angles of an Octoedron, or in the bases of an Octoedron be inscribed equilateral triangles subtending the solid angles of a Cube, there shall be produced a solid consisting of either of the whole solids, to wit, of the whole cube, and of the whole Octoedron.

But equilateral triangles inscribed in the bases of a Pyramis, having their angles set in the middle sections of the sides of the Pyramis,
and

R
and the
the said
produce
like pyr

And
you tak
restored
wit, the
decaedr
one, th
the soli
an Oct
shall be
solid a
compos
an Oct

Of

1. A
ded i
square
cut o
perpe
the
ramis

2.
Pyran
take
whic
the
in th

3.
of r
diam

4.
sect
is r

and the solid angles of a Pyramis, set upon the said equilateral triangles, there shall be produced a solid consisting of two equal and like pyramids.

And now if in these solids thus composed, you take away the solid angles, there shall be restored again the first composed solids, to wit, the solid angles taken away from a Dodecaedron and an Icosaedron composed into one, there shall be left an Icosidodecaedron, the solid angles taken away from a Cube and an Octoedron composed into one solid, there shall be left an Exoctoedron. Moreover, the solid angles taken away from two pyramids composed into one solid, there shall be left an Octoedron.

Of the nature of a trilateral and equilateral Pyramis.

1. A trilateral equilateral Pyramis is divided into two equal parts, by three equal squares, which in the center of the Pyramis cut one another into two equal parts, and perpendicularly, and whose angles are set in the middle sections of the sides of the Pyramis.

2. From a Pyramis are taken away four Pyramids like unto the whole, which utterly take away the sides of the Pyramis, and that which is left is an Octoedron inscribed in the Pyramis, in which all the solids inscribed in the Pyramis are contained.

3. A perpendicular drawn from the angle of the Pyramis to the base, is double to the diameter of the Cube inscribed in it.

4. And a right line coupling the middle sections of the opposite sides of the Pyramis is triple to the side of the same Cube.

5, The

5. The side also of a Pyramis is triple to the diameter of the base of the Cube.

6. Wherefore the same side of the Pyramis is in power double to the right line which coupleth the middle sections of the opposite sides.

7. And it is in power sesquialter to the perpendicular which is drawn from the angle to the base.

8. Wherefore the perpendicular is in power sesquitertia to the line which coupleth the middle sections of the opposite sides.

9. A Pyramis and an Octoedron inscribed in it, also an Icosaedron inscribed in the same Octoedron, do contain one and the same sphere.

Of the nature of an Octoedron.

1. Four perpendiculars of an Octoedron, drawn in four bases thereof from two opposite angles of the said Octoedron, and coupled together by those four bases, describe a Rhombus, or Diamond figure; one of whose diameters is in power double to the other diameter.

2. For it hath the same proportion that the diameter of the Octoedron hath to the side of the Octoedron.

3. An Octoedron and an Icosaedron inscribed in it, do contain one and the same sphere.

4. The diameter of the solid of the Octoedron is in power sesquialter to the diameter of the circle which containeth the base, and is in power duple superbipartiens tertias (that is, as 8 to 3,) to the perpendicular or side of the foresaid Rhombus; and moreover is

is in len
the cen

5. Th
of the
inclinat
angles

1. Th
quialte

2. A

3. A
centers
sextupl

4. A
side o
whole

5. U
as the

6. U
power

7.
power

8.

mis,

a ma

caedr

Pyra

1.

a sol

a Pe

ses

tag

in

wh

is in length triple to the line which coupleth the centers of the next bases.

5. The angle of the inclination of the bases of the Octoedron, doth with the angle of the inclination of the bases of the Pyramis, make angles equal to two right angles.

Of the nature of a Cube.

1. The diameter of a Cube is in power sesquialter to the diameter of his base.

2. And is in power triple to his side.

3. And unto the line which coupleth the centers of the next bases, it is in power sextuple.

4. Again, the side of the Cube, is to the side of the Icosaedron inscribed in it, as the whole is to the greater segment.

5. Unto the side of the Dodecaedron, it is as the whole is to the lesser segment.

6. Unto the side of the Octoedron it is in power duple.

7. Unto the side of the Pyramis it is in power subduple.

8. Again, the Cube is triple to the Pyramis, but to the cube the Dodecaedron is in a manner double. Wherefore the same Dodecaedron is in a manner sextuple to the said Pyramis.

Of the nature of the Icosaedron.

1. Five triangles of an Icosaedron; do make a solid angle, the bases of which triangles make a Pentagon. If therefore from the opposite bases of the Icosaedron be taken the other Pentagon by them described, these Pentagons shall in such sort cut the diameter of the Icosaedron which coupleth the foresaid opposite angles;

I i

that

that that part which is contained between the planes of these two Pentagons shall be the greater segment, and the residue which is drawn from the plain to the angle, shall be the lesser segment.

2. If the opposite angles of two bases joined together, be coupled by a right line, the greater segment of that right line is the side of the Icosaedron.

3. A line drawn from the center of the Icosaedron to the angles, is in power quintuple to half that line, which is taken between the Pentagons, or of the half of that line, which is drawn from the center of the circle which containeth the foresaid Pentagon, which two lines are therefore equal.

4. The side of the Icosaedron containeth in power either of them, and also the lesser segment, to wit, the line which falleth from the solid angle to the Pentagon.

5. The diameter of the Icosaedron containeth in power the whole line, which coupleth the opposite angles of the bases joined together, and the greater segment thereof, to wit, the side of the Icosaedron.

6. The diameter also is in power quintuple to the line which was taken between the Pentagons, or to the line which is drawn from the center to the circumference of the circle which containeth the Pentagon composed of the sides of the Icosaedron.

7. The dimetient containeth in power the right line which coupleth the centers of the opposite bases of the Icosaedron, and the diameter of the circle which containeth the base.

8. Again, the said dimetient containeth in power the diameter of the circle which containeth the Pentagon, and also the line which is drawn from the center of the same circle to the circumference; that is, it is quintuple to

to the
cumfer

9. T
the op
line w
bases,
the sid
is the

10.
ctions
side of
11.
and the
line, r
the D

1. T
in pow
so tha
Dodec
of the
ment,
angle
under
Dodec

2. I
caedro
of one
centers
propon
right
next b

3. I
one b
the c
the op
perfect
coupl

to the line drawn from the center to the circumference.

9. The line which coupleth the centers of the opposite bases, containeth in power the line which coupleth the centers of the next bases, and also the rest of that line of which the side of the Cube inscribed in the Icosaedron is the greater segment.

10. The line which coupleth the middle sections of the opposite sides, is triple to the side of the Dodecaedron inscribed in it.

11. Wherefore if the side of the Icosaedron, and the greater segment thereof be made one line, the third part of the whole is the side of the Dodecaedron inscribed in the Icosaedron.

Of the Dodecaedron.

1. The diameter of a Dodecaedron containeth in power the side of the Dodecaedron, and also that right line to which the side of the Dodecaedron is the lesser segment, and the side of the Cube inscribed in it is the greater segment, which line is that which subtendeth the angle of the inclination of the bases, contained under two perpendiculars of the bases of the Dodecaedron.

2. If there be taken two bases of the Dodecaedron, distant from one another by the length of one of the sides, a right line coupling their centers being divided by an extreme and mean proportion, maketh the greater segment the right line which coupleth the centers of the next bases.

3. If by the centers of five bases set upon one base, be drawn a plain superficies, and by the centers of the bases which are set upon the opposite base, be drawn also a plain superficies, and then be drawn a right line, coupling the centers of the opposite bases,

I i 2

that

that right line is so cut, that each of his parts set without the plain superficies, is the greater segment of that part which is contained between the plains.

4. The side of the Dodecaedron is the greater segment of the line which subtendeth the angle of the Pentagon.

5. A perpendicular line drawn from the center of the Dodecaedron to one of the bases, is in power quintuple to half the line which is between the plains.

6. And therefore the whole line which coupleth the centers of the opposite bases, is in power quintuple to the whole line which is between the said plains.

7. The line which subtendeth the angle of the base of the Dodecaedron, together with the side of the base are in power quintuple to the line which is drawn from the center of the circle which containeth the base, to the circumference.

8. A section of a sphere containing three bases of the Dodecaedron, taketh a third part of the diameter of the said sphere.

9. The side of the Dodecaedron and the line which subtendeth the angle of the Pentagon, are equal to the right line which coupleth the middle sections of the opposite sides of the Dodecaedron.

THE

1. T
sphere
cle eq
2. T
any se
or to
assign

THE
THEOREMS
OF
ARCHIMEDES.

Concerning the Sphere and Cylinder, Investigated by the *Method of Indivisibles*, and briefly Demonstrated by the Reverend and Learned Dr. *Isaac Barrow*.

THE main Design of *Archimedes* in his *Treatise of the Sphere and Cylinder*, is to resolve these four Problems.

1. To find the proportion of the superficies of a sphere to any determinate circle; or to find a circle equal to the superficies of a given sphere.

2. To find the proportion of the superficies of any segment of a sphere to any determined circle; or to find a circle equal to the superficies of any assign'd segment.

I & 3

3. To

3. To find the proportion of the sphere it self (or of its solid content) to any determinate Cone or Cylinder; or to find a Cone or Cylinder equal to a given sphere.

4. To find the proportion of a segment of a sphere to any determinate Cone or Cylinder; or to find a Cone or Cylinder equal to a given segment.

These four Problems Archimedes prosecutes separately, and lays down Theorems immediately subservient to their solution; but we reduce them to two: For since an Hemisphere is the segment of a sphere, and the method of finding out its relations, in respect to the superficies and solid content, is comprehended in the general method of investigating the proportion of the segments: And from the superficies and solid content of an Hemisphere already found, the double of them, (that is, the superficies and content of the whole sphere) is at the same time given. And indeed 'tis superfluous and foreign from the Laws of good Method, to investigate their relations distinctly and separately; so that if it were not a crime, I might on this account, blame even Archimedes himself.

The whole matter therefore is reduc'd to these two Problems.

1. To find the proportion of the superficies of any segment of a sphere, to a determinate circle; or to find a circle equal to the superficies of a given segment.

2. To find the proportion of the solidity of any segment of a sphere to any determinate Cone or Cylinder; or to find a Cone or Cylinder equal to an assign'd segment of a sphere.

I shall resolve these Problems by another much easier and shorter method: In which the order being inverted, first, I shall seek the solidity of a segment, and from thence deduce

its

its sup
ment
I know

Fi
segme
know

1. 7
Arithm
whose
nitude
to the
the g
if th
term

z =

Th
appe
verti

For
the
two
equa
serie
repe
i. e.
term

W
fy
whi
gran
the

its superficies; a thing which is in my judgment well worth observing, and perform'd, as I know of, by none.

First therefore, for finding the solidity of a segment, I shall lay down two, commonly known and receiv'd Suppositions, viz.

1. That a series of magnitudes proceeding in Arithmetical Progression from nothing (inclusive) or whose common difference is equal to the least magnitude, is subduple of as many quantities equal to the greatest; (i. e. subduple of the product of the greatest term and number of terms:) So that if the sum of the terms be called z , the greatest term g , and the number of terms n , then will

$$z = \frac{ng}{2}, \text{ or } 2z = ng.$$

The truth of this Proposition will easily appear by expressing the series twice, and inverting the order;

0, 2, 2a, 3a, 4a.

4a, 3a, 2a, a, 0.

For so the difference always being equal to the least quantity, 'twill be evident that each two correspondent terms taken together are equal to the greatest term; and also, that the series taken twice is equal to the greatest term repeated as many times as there are terms, i. e. the last term drawn into the number of terms.

We have in a triangle a very clear and easy example of this most useful Proposition, which is prov'd hence, to be half a parallelogram having the same altitude, and standing on the same base.

I i 4 Suppose




Suppose the altitude AE of the triangle AEZ to be divided into parts indefinitely many and small AB, BC, CD, DE, and parallels BZ, CZ, DZ, EZ, drawn thro' the points of Divisions; all these proceed from o in

an Arithmetical Progression, and consequently the sum of 'em all, that is, the triangle AEZ, is subduple of the greatest EZ drawn into the altitude AE, by which the sum of the terms is express'd, that is, subduple of the Parallelogram EY, whose base is EZ, and altitude AE.

But the illustration of the Rule will conduce more to our design by inferring hence, *That a circle is equal to half of the radius drawn into the circumference*, after this manner. Conceive a circle to consist of as many concentric Peripheries as there are points or equal parts indefinitely many and small in the radius. These Peripheries, as well as their radii proceed from the center or nothing in an Arithmetical progression; and therefore their sum, that is, the whole circle is equal to half the greatest (or extreme circumference) drawn into the number of terms, that is, the radius.

After the same manner we may suppose the sector AEZ to consist of as many concentric Arcs BZ, CZ, DZ, EZ, as there are points (or equal parts indefinitely small) in the radius AE, which Arcs, as their radii, proceeding from a point or nothing in an Arithmetical progression, the sector



also

also will
to the
made ev
suppose
lar to th
and fro
the radi
and term
AE: ra
EZ, th
manner
the tri

AEZ,

for A
lebrate
qual to
and alti
that v
ption,
Area
finitely
method
me) f
and p
planes
lines,
after
by th
cylind
ing t
the f
with
tion
ther
z.
ceed
prog

also will be equal to half the radius drawn in to the extreme Arc EZ. Which may be made evident also after this manner: Let us suppose the right line EY to be perpendicular to the radius AE, draw the right line AY, and from the points B, C, D, of division in the radius, draw BY, CY, DY, parallel to EY, and terminated at AY. Because EY: DY ($:: \text{rad. AE} : \text{rad. AD}$) $:: \text{Arc EZ} : \text{Arc DZ}$. and EY = EZ, then will DY = Arc DZ; and in like manner will CY = CZ, and BY = BZ. whence the triangle AEY will be = to the sector AEZ, that is, $\frac{AE \times EY}{2} \left(\frac{AE \times EZ}{2} \right) = \text{sector AEZ}$.

By this means we collect that celebrated Theorem of Archimedes, That a circle is equal to a triangle, whose base is equal to the radius, and altitude equal to the periphery of the circle; and that without any inscription or circumscription, of figures, by only supposing that the Area or Superficies of the circle consists of infinitely many concentric Peripheries. Which method of indivisibles, (now first of all known to me) seems no less evident (nay more evident) and perhaps less fallacious than that wherein planes are supposed to consist of parallel right lines, and solids of parallel planes; as hereafter shall be evident, when we shall collect, by this method, the proportions of spheric and cylindric superficies to one another, by knowing the solid content; and on the other hand, the solid content, by knowing the superficies, with admirable facility, and most full satisfaction in those things which are rigidly gathered by pure Geometry.

2. Let us suppose a series of quantities to proceed from 0 (inclusive) in a duplicate Arithmetic progression, that is, 0, 1, 4, 9, 16, &c. the squares of

of numbers in a simple Arithmetic progression, 0, 1, 2, 3, 4, &c. And the triple of this series will always exceed the greatest term multiplied by the number of terms; but the number of terms increasing the proportion, continually approximates till at last it comes to an equality, when the number of terms is increased in infinitum.

$$3 \times 0 + 1 = 3.3$$

$$2 \times 1 = 2.2$$

$$3 \times 0 + 1 + 4 = 15.15 = 5$$

$$3 \times 4 = 12.12 = 4$$

$$3 \times 0 + 1 + 4 + 9 = 42.42 = 7$$

$$4 \times 9 = 36.36 = 6$$

$$3 \times 0 + 1 + 4 + 9 + 16 = 90.90 = 9$$

$$5 \times 16 = 80.80 = 8$$

$$3 \times 0 + 1 + 4 + 9 + 16 + 25 = 165.165 = 11$$

$$6 \times 25 = 125.125 = 10$$

As for example, if the terms be two, the triple of the terms will be to the greatest term drawn into the number of terms as 3 to 2; if there be three terms as 5 to 4; if four, as 7 to 6; if five, as 9 to 8, and so continually: So that the antecedents of these proportions always mutually exceed one another by the number 2; and so every antecedent its consequent by 1. Whence it is evident that by how much the greater the number of terms is, by so much the more the proportion tends to equality. So 100 to 99 is less distant from the proportion of equality than 10 to 9. From hence, supposing the number of terms infinite (or infinitely great,) the triple of quantities proceeding thus in a duplicate proportion (or as the squares of the numbers, 0, 1, 2, 3, 4, &c. will be equal to as many quantities equal to the greatest term.

The

The
down
the F
that,
ted by
Howe
ter by
our C
intelli
ctions.
from
of a
the
less
two
the p
terms
It
of th
subtri
and
AE
and
rall
be a
squa
ters
circ



T
of t
men
logr


The same, as to the substance of it, is laid down by *Archimedes* in his *Book of Spirals*, as the Foundation of many Argumentations, in that, and other Books, and is well demonstrated by our Learned Country-man *Dr. Wallis*: However, I thought fit to illustrate the matter by this method, as being not unworthy our Consideration, and very perspicuous and intelligible in this, that 'tis free from Fractions. And by the way 'tis observ'd, that from hence we may easily find the proportion of a series triple to as many terms equal to the greatest, viz. as twice the number of terms less one, to twice the number of terms less two. So that if the number of terms be 6, the proportion of a series triple to as many terms equal to the greatest will be as 11.

It will be a very easy and apt Illustration of this Rule, if we infer hence, That a Cone is subtriple of a Cylinder, having an equal base and altitude. For let us suppose the altitude *AE* of the cone *ZY* to be divided into equal and indefinitely many parts, by as many parallel right lines *ZY*, and the lines *ZY* will be as the numbers 1, 2, 3, 4, &c. and the squares or circles constituted upon the diameters *ZY*, as 1, 4, 9, 16, &c. whence all those circles, or the whole cone *AZY* (made up of the same) will be subtriple of as many circles equal to the greatest, constituted on the greatest diameter *ZEY*, that is, subtriple of a cylinder whose base is *AEY*, and altitude *AE*.



There occur two other most apt examples of this Rule, viz. by inferring, That the complement of a Semiparabola is subtriple of a parallelogram having the same base and height; as also,

also, That the space comprehended by the Spiral and Radius is subtriple of the circle in which the Spiral is generated: But of these in another place. Wherefore to go on with what we began, these two Rules being supposed; let us conceive ZAY to be a segment of a



sphere, X its center, AT its diameter, and ZAYT a great circle passing thro' the vertex, and the part AE of the Axe to be divided into an indefinitely many equal parts; and let us imagine parallel lines to be drawn thro' the points of Division, generating circles in the sphere,

whose Radii let be BZ, CZ, DZ, and diameters ZY. I suppose the segment of a sphere to consist of all these parallel circles, whose number is as great as that of the points, or equal indefinitely many small parts in the Axe AE, according to the known *Method of Indivisibles*.

But now for brevity's sake, let the diameter AT be called d , and the radius of the sphere r (if need be,) and the Axe AE, by which the number of terms is express'd, call n , and one of the equal parts a ; which being settled, 'tis evident, (*by the Elements*) that $BZ^2 = AB \times BT = a \times d - a = ad - a^2$, and in like manner $CZ^2 = AC \times CT = 2a \times d - 2a = 2ad - 4a^2$, and by the same reasoning $DZ^2 = AD \times DT = 3ad - 9a^2$, and $EZ^2 = AE \times ET = 4ad - 16a^2$, &c. that is, that the squares of the radii of the circles ZY are

to one
and, &c.
Progress
 $9a^2$, 16
of the
first R
and, &
equal
nd
=

Mo
square
gethe
terms

is, =

W
the

equ

term

3 n

the

ar

W

C

fi

tu

b

a

to one another as the rectangles ad , $2ad$, $3ad$, $4ad$, &c. (which proceed in an Arithmetical Progression from 0) less by the square a^2 , $4a^2$, $9a^2$, $16a^2$, &c. which go on as the squares of the numbers, 1, 2, 3, 4, &c.) But by our first Rule, all the Rectangles 0, ad , $2ad$, $3ad$, $4ad$, &c. are equal to half as many terms equal to the greatest $AE \times AT$ or nd , that is,

$$= \frac{nd \times n}{2}.$$

Moreover, by our second Rule, all the squares 0, a^2 , $4a^2$, $9a^2$, $16a^2$, &c. taken together, are equal to a third part of as many terms equal to the greatest AE^2 or n^2 , that is,

$$= \frac{n^2 \times n}{3}.$$

Wherefore all the squares described upon the radii BZ , CZ , DZ , EZ , conjunctly, are equal to the difference $\frac{ndn}{2} - \frac{nnn}{3}$, (or the terms being reduc'd to the same denomination,) $\frac{3ndn - 2nnn}{6}$ and their quadruple, that is, all

the squares described upon the diameter ZY , are equal to $\frac{12ndn - 8n^3}{6}$ or $\frac{6ndn - 4n^3}{3}$.

Whence a segment of a sphere is equal to a Cylinder, the diameter of whose base is the side of a square equal to $6nd - 4n^2$, and altitude is $\frac{1}{3}n$; or to a cone having the same base, but the altitude n , or which is all one, having

a base whose radius is $\sqrt{\frac{6nd - 4n^2}{4}}$ or $\sqrt{\frac{3}{2}nd - n^2}$, and altitude n as before. Which

Cone we may change into a Cone upon the same base ZY with the segment ZAY , by saying, as ZE^2 (i. e. $dn - n^2$) to $\frac{3}{2}nd - n^2$ or (both

(both terms being divided by n) as $d - n$ to $\frac{2}{3}d - n$, so reciprocally n to the altitude of the Cone sought: Or in the figure by making, as TE to $TE + KA$, so is EA to ES . For ES will be the altitude of the Cone ZSY equal to the segment of the sphere ZAY . Which is a noted Theorem of *Archimedes*, demonstrated by him with so much labour and prolixity.

Hence, if the given segment be a Hemisphere, and so $n = \frac{1}{2}d$ or r , then d or $2r$ will be the altitude of a Cone, which having a base equal to the base of the Hemisphere, (or to the greatest circle in the sphere,) will be equal to the Hemisphere. And a Cone whose base is double of the greatest circle, and the altitude $2r$, or the Cylinder whose base is $\frac{2}{3}$ of the greatest circle, and altitude $2r$, will be equal to the whole Sphere. Whence the whole Sphere is $\frac{2}{3}$ of a Cylinder the diameter of whose base is $2r$, and the altitude also $2r$. And this is the chief Theorem of *Archimedes*, viz. That a sphere is subsesquialter or $\frac{2}{3}$ of that Cylinder, whose Altitude and Diameter of the base is equal to the Diameter of the Sphere.

Furthermore, not to pass over any thing in our Author which seems to be to our purpose:

If to the sum first found, representing a segment, viz. $6ndn - 4unn$, we add $2d^2n - 6dn^2 + 4n^3$

$$\left(= \frac{4dn - 4n^2}{3} \times \frac{d - 2n}{2} = \frac{4}{3} ZE^2 \times XE \right)$$
 representing the Cone ZXY , the aggregate $\frac{2}{3}ddn$ will represent the Sector of the Sphere $ZX-YA$, which for that reason will be equal to a Cylinder, the diameter of whose base $\sqrt{dn}$, and


and the
meter of
 $2d$, or a
base is
ing rec
to a Co
Line A
circum
 $AZ^2 =$
And th
medes,
of the
is equa
scribed
the ver
segmen
dus of
And
which
its pa
cuity.
soluti
posed
of a
To o
cle to
the
the
least
supp
rical



and the altitude $\frac{1}{2}d$, or to a Cone, the diameter of whose base is $\sqrt{dn}$, and the altitude zd , or also in a Cone, the Radius of whose base is $\sqrt{dn}$, and the altitude $\frac{1}{2}d = r$ (it being reciprocally as $4dn : dn :: zd : \frac{1}{2}d$), that is, to a Cone, the Radius of whose base is the Line AZ, drawn from the vertex to the circumference of the base of the segment, (for $AZ^2 = TA \times AE = dn$), and the altitude r . And this is the next famous Theorem of *Archimedes*, concerning the solidity of the sector of the Sphere, viz. *That the sector of a sphere is equal to a Cone, whose base is a circle described by a Radius equal to a line drawn from the vertex to the circumference of the base of the segment, and whose altitude is equal to the Radius of the sphere.*

And thus I think I have completed that which belongs to the solidity of a sphere, and its parts with sufficient brevity and perspicuity. From hence we shall deduce the Resolution of the other Problem, which I proposed concerning the surface of the segment of a sphere; and then of the whole sphere. To obtain this, as we supposed, before a Circle to consist of concentric Peripheries, and the Sector of a Circle, of concentric Arcs, (in the number of which, the greatest, and the least or a point is reckon'd: So now we suppose spheres to consist of concentric spherical superficies, and the Sectors of Spheres, of

like concentric superficies; as for example, the sector of the sphere ZAE, of the superficies BZ, CZ, DZ, EZ, &c.) which supposition indeed seems so easy and natural, that in my judgment 'tis sufficient



cient only to propose it ; neither is a further explication wanting to gain an assent to it.

2. We suppose these spherical superficies to be in a duplicate Ratio of the Radius of the spheres : This is the common affection of all like superficies, and it seems to agree very well with the superficies of spheres, because they appear to be most uniform and similar. But this Supposition might easily be evinc'd and establish'd by the same sort of arguing, as spheres are proved to be in triplicate proportion to their Diameters or Radii ; Or might have been join'd as a *Corollary* to *Prop.* 17. and 18. *Elem.* 12. where the superficies of like Polygons are suppos'd to be inscribed in spheres, having as well the superficies in a duplicate, as the solidity in a triplicate Ratio of the Diameters of the Spheres. These things being premis'd, let us suppose *AE* a Radius or the side of the Sector of a Sphere *EAZ*, to be divided into equal and indefinitely many small parts, and the sector *AEZ* to consist of these spherical superficies *BZ, CZ, DZ, EZ*, it will be evident that all those superficies in the Progression are as the squares of the Radii, that is, as $\overline{AB}^2, \overline{AC}^2, \overline{AD}^2, \overline{AE}^2$, &c. or as the squares of the numbers 1, 2, 3, 4, &c. whence by our second Rule, the sum of all these superficies, that is, the sector *AEZ*, will be $\frac{1}{3}$ of as many superficies equal to the greatest *FZ*, that is, $\frac{1}{3}$ of the greatest *EZ*, drawn into r the number of terms. Whence a sector is equal to a Cylinder whose base is $\frac{1}{3}$ of the greatest or extreme superficies of the sector, and whose altitude is r : Or to a Cone whose base is equal to the superficies of the sector, and its altitude r , which is the last of *Lib.* 1.) but we just now prov'd that a sector.

sector is equal to a Cone whose altitude is r , and base a circle describ'd by the Radius YE, drawn from the vertex of the segment EYZ to the circumference of the base. Wherefore, a Cone whose altitude is, and base equal to the superficies of the sector, is equal to a Cone of the same altitude, whole base is a circle describ'd by the Radius YE.

And so the superficies of the sector EYZ is equal to a circle describ'd by the Radius YE, Which, certainly is the principal Theorem of all those that occur in the Books of Archimedes, nor is there found a more excellent one in all Geometry; viz. That the superficies of any segment of a sphere is equal to a circle whose Radius is a right line drawn from the vertex of the segment to the circumference of the bases: And hence, That the superficies of an Hemisphere is double of the base, or equal to two great circles of the sphere.

For in this Case $\overline{YE}^2 = \overline{AZ}^2 + \overline{AY}^2 = 2 \overline{AE}^2$ and consequently a circle described by the Radius YE is equal to two circles describ'd by the Radius AE. Whence also, the superficies of the whole sphere is quadruple a circle having the same Radius with the sphere, that is, quadruple




the greatest circle in the sphere; and equal to a circle whose Radius is the diameter of the sphere. From hence it follows, That the superficies of a sphere is equal to the superficies of a Cylinder of the same height and breadth; for the superficies of that Cylinder is quadruple to the base, as we shall shew hereafter. And these are the most noted Theorems of Archimedes. Nay, from hence all those things follow, which he

has written concerning the superficies of spheres, and their segments. So that from these few and easy Suppositions, I have demonstrated whatever seem to be of any Note in the Books *Of the Sphere and Cylinder*.

I will only add, that after by the method of *Archimedes*, (for I think scarce any other can be invented, besides ours, for finding the Solidity) the superficies of segments are found equal to the circle described by the Radii YE; hence it will plainly follow, that the superficies of spheres, and thence of like sectors are in a duplicate ratio of the Radii of the spheres; and consequently from the superficies thus found, the contents of segments, and of whole spheres may be mutually deduced, and that very clearly and expeditiously after this manner. Because in the sector EAZ (fig. Pag. 358.) the superficies BZ, CZ, DZ, EZ, proceed as the squares describ'd upon AB, AC, AD, AE, that is, as 1, 4, 9, 16, &c. the whole sector will be equal to $\frac{1}{3}$ of as many superficies equal to the greatest EZ, or $EZ \times r$, that is, to a Cylinder whose base is $\frac{1}{3} EZ$, and altitude r , or to a Cone whose base is $\frac{1}{3} EZ$, and altitude r . But EZ is supposed equal to a circle whose Radius is YE, wherefore the sector EAZ is equal to a Cone whose base is a circle described by the Radius YZ and altitude r : Which is *Archimedes's* universal Theorem for the contents of Circles. Whence if from this the Cone ZAE standing on the base of the segment EYZ, and having the vertex at the center of the sphere A, be subducted, (you'll have that segment EYZ.) But when the sector EYZ is a Hemisphere, there will be no such Cone to be subducted; and for that reason a Cylinder whose base is $\frac{2}{3} EZ$, and altitude r , or the

the Cone whose base is $2EZ$, and altitude likewise r will be equal to the whole sphere. But the superficies of the Hemisphere EZ , is proved to be equal to two of the greatest circles in the sphere, whence the whole sphere is given. This is *Archimedes's* first and principal Theorem, for the content of a sphere; whence 'tis easily deduced, that a sphere is $\frac{2}{3}$ of a circumscrib'd Cylinder, that is, of a Cylinder whose altitude and diameter of its base is equal to the diameter of the sphere.

The Doctrine of our other [*Archimedes*] seems to make against, and subvert the new and celebrated *Method of Indivisibles*, and is press'd to that end by *Tacquet*, for instance, (*Prop. 2. lib. 2. Cylindr.*) For the usual process of that method seems to exhibit the dimension of the superficies of a Cone, (as also of a sphere, and of other Curves) different enough from what our Author and others have demonstrated:



As for example, let us suppose $ABCD$ a right Cone, whose Axe is AX , and base BCD , and plane $\beta \chi \delta$ drawn, at pleasure, parallel to the base BCD . And since, as $Diam. BD : Periph. BCD :: Diam. \beta \delta : Periph. \beta \chi \delta$, and so every where it will be (according to the *Method*

of *Indivisibles*, and by 12. 5.) as $Diam. BD$, to $Periph. BCD$, so is the triangle ABD , consisting of those parallel Diameters, to the Conic Superficies $ABCD$, consisting of those Peripheries, i. e. $Diam. BD : Periph. BCD :: AX \times BD : AX \times Periph. BCD$

Whence

$$\frac{2}{K k z} \quad \frac{2}{AX}$$

$\frac{AX \times \text{Periph. BCD}}{2}$ will be equal to the superficies of the Cone; which is false and contrary to what was demonstrated just now. For we demonstrated that the superficies of the Cone was $\frac{AB \times \text{Periph. BCD}}{2}$.


In answering this Objection, we say, that the *Method of Indivisibles*, in the speculation of Perimeters, and of Curve Surfaces, proceeds otherwise than in the speculation of plane Surfaces and solid Contents. It does indeed suppose that the Area of plane Figures consist, as it were of parallel right lines, and the contents of solids of parallel Planes, and that their number may be express'd by the altitude of the Figures: But it by no means supposes, that the Perimeters of plane figures consist of points, or the superficies of solids, of lines, the number of which may be express'd by the altitude of the figure. As for example, altho' the triangle ABD (in the last figure) consists of lines parallel to BD, the number of which is expressed by the number of points in the perpendicular AX, that is, by the length of the perpendicular: Yet it would be absurd to suppose that the line AB consists of points, whose number may be express'd by the number of points in a less line AX. For altho' the right line $\beta\epsilon$ drawn thro' each infinitely small part of AX, divide AB into as many infinitely small parts, yet those parts are not of the same Denomination or Quality with the parts of AX, but somewhat greater than them; so that if the parts of AX be look'd upon as points, the parts of AB are not to be called points, but greater than points; and on the contrary, if the parts of AB be called points,

points, the parts of AX are to be look'd upon as less than points, if it be lawful to speak so. For the points which are treated of in the *Method of Indivisibles* are not absolutely points, but indefinitely small parts, which usurp the names of points, because of the Affinity. Since therefore points don't admit of greater and less, the name of points is not at the same time to be attributed to the parts of different magnitudes; consequently tho' the number of the greater parts of AB may be express'd by the number of the lesser parts of AX, yet the number of points in AB can no ways be expressed by the number of points in AX, (that is, by the number of parts in AX, equal to the number of parts in AB, which are called points.) The line AB has as many points as there are in it self alone, or another line equal to it self, nor can it be determin'd by any other measure. After the same manner, this method don't suppose the conic Surface ABCD to consist of as many parallel circumferences perpetually increasing from the vertex A, or decreasing from the base BD, as there are points in the Axe AX; but rather of as many thus increasing or decreasing as there are points in the side AB. For in the Revolution of the line AB about the Axis AX, (whereby the superficies of the Cone is generated) every point in the line AB produces a circumference, and consequently more circumferences are produced than the points contained in the Axis AX. Therefore if you would extend the *Method of Indivisibles* to the superficies of solids, and suppose those superficies to consist of parallel lines, you ought not to compute this by the parallel Areas constituting the solid, that is, not to number those Areas by the altitude of the solid, but by other lines agreeable to the condition of each figure. Which lines, in figures that are not irregular, may easily be determin'd: For instance,



stance, in the equilateral Pyramid ABCD, whose Axe is AX, supposing that the lateral surface of the Pyramid consists of Perimeters of triangles, parallel to the base BCD, these can neither be computed by the altitude

AX, nor by the side AB, (for by the former, the thing requir'd, would be wanting of the true Dimension, and by the latter 'twould exceed it,) but by the line AE drawn from the vertex A perpendicular to the side BC of the base; the reason of which is, that every plane side of a Pyramid as ABC, consists of parallel right lines computed by the altitude AE. After the same manner, supposing that the superficies of the Hemisphere



BAD, consist of peripheries of circles parallel to the base BCD, the number of them is not to be computed by the Axis AX, but by the Quadrantal Arc AB, because that every point of


the Arc AD in revolving produces a circumference. And so any superficies, whether plane or curv'd, which is conceived to consist of equidistant right or curv'd lines, is to be computed by a line cutting those equidistant lines perpendicularly. For since those equidistant lines, in this *Method of Indivisibles*, are not consider'd absolutely as lines having an infinitely small breadth, which is the same with the breadth or thickness of the point describing those equidistant lines in their Circumvolution, and since the same equidistant lines divide the line cutting them perpendicularly into parts measuring its breadth, those parts are to be look'd upon as such sort of points, and consequently the number of equidistant lines, or the sum of those breadths is to be computed by the number of points in the line cutting

cutting them perpendicularly, that is, by the length of that line, and not by a line of any other length, for that will consist of more or less points.

Hence therefore in the speculation of the superficies of solids, the *Method of Indivisibles* is not unuseful, but rather very commodious, provided it be rightly understood, and applied according to the Rule prescrib'd. For by the help of it even these superficies may be found, if so be we have some convenient *Data* presuppos'd, on which the reasoning may be founded. For instance, we might by the help of it, investigate the superficies of a Cone, by reasoning after this manner.

If the superficies of the cone ABC (fig. pag. 362.) be divided into innumerable Peripheries of circles $\beta\chi\delta$ parallel to the base BCD, the breadth of those Peripheries taken together, make up the side AB cutting them perpendicularly, and consequently there will be as many Peripheries as there are points in the line AB, that is, their number may be express'd by the number of points in AB, or by its length. Wherefore, if you draw perpendiculars equal to the Peripheries to every point of AB, a superficies will be made out of those perpendiculars equal to the superficies of the Cone. But that superficies will be a triangle whose height is AB, and base equal to the greatest Periphery BDC, and so the superficies of the Cone will be $= \frac{1}{2} AB \times \text{Periph. BDC}$, which conclusion agrees with the things laid down and demonstrated by *Archimedes*.

After the same manner, if you take any right




line $\alpha\beta$ equal to the quadrantal Arc AB of the Hemisphere (in pag. 364.) and to each of its points μ let the per-

perpendiculars uw be erected equal to the Radii MN of parallel circles MOM passing thro' the corresponding points M of that quadrantal Arc, the greatest of which $\beta\xi$ let be equal to the Radius BX of the base of the Hemisphere: The figure $a\beta\xi$ will contain the Radii of all the circles of whose Peripheries the superficies of the Hemisphere consists. And if the perpendicular uw , $\beta\delta$ be erected equal to the Peripheries MOM, BDB, there will be made the figure $a\beta\delta$ equal to the superficies of the Hemisphere. The dimension of which figure if you can by any means find (as in this case you are to find the Area of the figure $a\beta\xi$) thence you will easily deduce the content of the segment of the sphere, agreeing to what you may gather by any lawful reasoning. Which Observation, I think will not be unuseful in Geometry.

E I N I S.

AT the Hand and Pen in Bar-
bican are Taught, Writing,
Common and Merchants Accounts,
after that well approved Method of
Mr. Normansel, Algebra, Geometry,
Measuring, Surveying, Gauging, Na-
vigation and Dialing, with other
parts of the Mathematicks, also the
Use of the Globes, Maps, and other
Instruments, by me

Robert Arnold.



S
e
v
e
n

f
r
e
r